

PEARS (Eastern U.S.): Scab—Use 2 pints in pre-blossom, petal fall and cover sprays. Repeat cover applications at 7 to 14 day intervals under severe scab conditions. Rusting may be reduced on Bosc pears. Do not use on D'Anjou pears.

PEARS (Western U.S., except Pacific Northwest): Pear scab—Use 2 pints. Apply during early finger stage, petal fall and first cover sprays. Under severe scab infection conditions additional treatments may be necessary. Rusting may be reduced on Bosc pears. Do not use on D'Anjou pears.

PLUMS, PRUNES (Western U.S.): Brown rot—Use 2 pints. Apply at green bud, popcorn, bloom and petal fall stages. Repeat if cover sprays as conditions warrant. Prune russet scab (Italy, west of California)—Apply 2 pints per 100 gallons water. Apply at full bloom.

PLUMS, PRUNES (Eastern U.S.): Brown rot—Use 2 pints in full pink, bloom and petal fall sprays. Repeat applications at 7 to 14 day intervals as needed to maintain cover. Continue applications through harvest if conditions favor brown rot. The addition of a neutral spreader has improved coverage.

NECTARINES: Brown rot, scab—Use 2 pints in full pink, bloom, petal fall, shuck, cover and preharvest sprays. Applications at 3 to 4 day intervals may be necessary during bloom to control blossom blight. Repeat application at 7 to 14 day intervals as needed to maintain cover. Continue applications through harvest if conditions favor brown rot. If powdery mildew is a problem, add 3 pounds sulfur to the petal fall, shuck and early cover spray. If sulfur is added, CAPTAN may be reduced to 1 pint in these sprays.

Coryneum blight (peach blight, shot-hole)—Use 2 pints, apply in pink bud, full bloom, petal fall and cover periods as necessary and as post harvest spray (but before leaves drop).

STRAWBERRIES: Botrytis fruit rot—Use 1-1½ quarts. Apply 200 gallons of spray per acre or an equivalent of 2-3 quarts per acre in sufficient water for even coverage. Make first application when new growth starts in the spring before the fruit starts to form. Repeat at weekly intervals. Under severe infection conditions make continuous applications through harvest period, treating immediately after each picking.

CANTALOUPE, CUCUMBERS: Angular leaf spot, anthracnose—Use 1 quart per 100 gallons water, or apply 2 quarts per acre in sufficient water for good coverage. Apply at first sign of disease and repeat at 5-7 day intervals or as needed to maintain disease control.

TOMATOES: Anthracnose, gray leaf spot, septoria leaf spot—Use 6 pints per acre in sufficient water for thorough coverage. Under severe infection conditions use 6 to 8 pints per acre in sufficient water for thorough coverage. Apply at first signs of infection and repeat at 5 to 7 day intervals or as needed to maintain control.

POSTHARVEST DIP: CAPTAN 4L can be used as a postharvest dip or wash on the following fruits for prevention of various rots and molds (Rhizopus, Botrytis, Gloeosporium) during storage: Apples, cherries, and pears—Dilution Rate: Use 1½ quarts per 100 gallons water. Apply as a spray or in dip tank. When used as a dip, recharge wash solution periodically as volume in tank is depleted. When tank volume is reduced 25%, bring water back to volume by adding ¾ quart CAPTAN 4L for every 25 gallons of water added. At end of every 8-hour shift, empty tank, flush, and charge with fresh dilution. Dispose of spent solution in accordance with Federal, State, and Local Laws. Do not allow dip tank solution to stand overnight. Maintain continuous agitation during dipping operation.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE INSTRUCTIONS: Storage should be under lock and key and secure from access by unauthorized persons and children. Storage should be in a cool dry area away from any heat or ignition source. Avoid storage at high temperatures. Do not stack over 2 pallets high. Move containers by handles or cases. Do not move containers from one area to another unless they are securely sealed. Keep container tightly sealed when not in use. Keep away from any puncture source. Avoid storage near water supplies, food, feed, and fertilizer to avoid contamination. Avoid contamination with acids or alkalis. Store in original containers only. If the contents are leaking or material is spilled follow these steps:

1. Contain spill, absorb with a material such as sawdust, clay granules or dirt.
2. Collect and place in suitable containers for disposal.
3. Wash area with water and soap to remove remaining pesticide.
4. Follow washing with clean water rinse.
5. Place a leaking container in a plastic tub and transfer contents as soon as possible to an empty original container.
6. Do not allow runoff to enter sewer or contaminate water supplies.
7. Dispose of waste as indicated below.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the hazardous waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY—CONDITION OF SALE

OUR RECOMMENDATIONS FOR USE of this product are based upon tests believed reliable. Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the Seller. Buyer assumes all risks of use, storage, and handling of this material not in strict accordance with directions given herewith. In no case shall Orezel or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product when such use and/or handling is not in strict accordance with directions given herewith. The foregoing is a condition of sale by Orezel Chemical Company and is accepted as such by the Buyer.

CAPTAN MIXTURE LIQUID NA9099RQ

BEST AVAILABLE COPY

