

Dimethoate 4EC

Systemic Insecticide - Miticide

(EPA Reg. No. 19713-231)
FOR USE IN SORGHUM (MILO)

READTHE ENTIRE DREXEL DIMETHOATE 4EC LABEL BEFORE PROCEEDING A C C E P T E D

WARNING DIRECTIONS FOR USE

JUL 6 2007

Under the Federal Insecticide, Funglishe, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 15713-231

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This labeling and the EPA registered labeling must be in the possession of the user at time of pesticide application.

All applicable directions, restrictions and use precautions on Drexel DIMETHOATE 4EC LABEL(s) are to be followed. This labeling contains supplemental instructions and restrictions for use of this product which may not appear on the Drexel DIMETHOATE 4EC label. Follow the instructions carefully.

Crop	Pest Controlled	Rate per Acre	PHI (Days)
Sorghum (Milo)	Aphids (Green bugs)	0.5 to 1 pt.	28
	SPECIFIC DIRECTIONS: Ground Application - Apply above rates in 25 to 40 gals. of water per acre. Air Application - Apply above rates in 1 or more gal. of water per acre. Do not feed or graze Milo within 28 days of last application. Make no more than 3 applications as needed per season. Do not apply during the poller-shed period.		
	Grasshoppers, Mites (including Banks grass mites (excluding Trans Pecos area of TX)), Twospotted spider mites	1 pt.	28
	SPECIFIC DIRECTIONS: Ground Application - Apply above rates in 25 to 40 gals. of water per acre. Air Application - Apply above rates in 1 or more gal. of water per acre. Do not feed or graze Milo within 28 days of last application. Make no more than 3 applications as needed per season. Do not apply during the pollen-shed period.		
	Sorghum midge	0.25 to 0.5 pt.	28
	SPECIFIC DIRECTIONS: Ground Application - Apply above rates in 25 to 40 gals. of water per acre. Air Application - Apply above rates in 1 or more gal, of water per acre. Do not feed or graze Milo within 28 days of last application. Make no more than 3 applications as needed per season. Do not apply during the pollen-shed period.		