

19713-238

9-30-1986

jacket

1/4



U.S. ENVIRONMENTAL PROTECTION AGENCY  
Office of Pesticide Programs  
Registration Division (H7505C)  
401 "M" St., S.W.  
Washington, D.C. 20460

EPA Reg. Number:  
  
19713-238

Date of Issuance:  
  
SEP 30 1996

NOTICE OF PESTICIDE:  
     Registration  
  X   Reregistration

(under FIFRA, as amended)

Term of Issuance:  
Unconditional

Name of Pesticide Product:  
Sulfur 90W

Name and Address of Registrant (include ZIP Code):

Ms. Barbara Brown  
Drexel Chemical Company  
2487 Pennsylvania Street  
P.O. Box Box 9306  
Memphis, TN 38109

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

Based on your response to the Reregistration Eligibility Document, EPA has reregistered the product listed above. Enclosed is a copy of your label stamped "Accepted". This action is taken under the authority of section 4(g)(2)(C) of the Federal Insecticide, Fungicide and Rodenticide Act, as amended. Reregistration under this section does not eliminate the need for continual reassessment of pesticides. EPA may require submission of data at any time to maintain registration of your product.

Signature of Approving Official:

Date:

9-30-96

2/4

ACCEPTED

9-30-96

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 19713-238



# Sulfur 90W

A Wettable Sulfur

**ACTIVE INGREDIENT:**  
Elemental Sulfur ..... 90.0%  
**INERT INGREDIENTS:** ..... 10.0%  
**TOTAL:** ..... 100.0%

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

Read Entire Label Before Using This Product

EPA Reg. No. 19713-238  
EPA Est. No. 19713-GA-1

■ 238SP-1006

### STATEMENT OF PRACTICAL TREATMENT

**IF SWALLOWED:** Call a physician or poison control center. Drink 1 or 2 glasses of water and induce vomiting by louching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.  
**IF IN EYES:** Flush eyes with plenty of water. Call a physician if irritation persists.  
**IF ON SKIN:** Wash skin with soap and plenty of water. If irritation develops, get medical attention.

### PRECAUTIONARY STATEMENTS

**Hazards To Humans and Domestic Animals**

**CAUTION:** Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling.

#### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:  
Long-sleeved shirt and long pants  
Waterproof gloves  
Shoes plus socks  
Protective eyewear  
Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### USER SAFETY RECOMMENDATIONS

Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.  
Remove clothing immediately if pesticides get inside. Keep and wash PPE separately from other laundry.

### ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water when disposing of equipment washwaters.

#### PHYSICAL HAZARDS

Keep away from heat, sparks or flame.

#### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

**DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

Personal Protective Equipment required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is: Coveralls, waterproof gloves, shoes plus socks, and protective eyewear.

### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides. (40CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Keep children and domestic animals off treated areas until this material has been washed into the soil.

**IMPORTANT:** Some fruits and other plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur on any crop unless local use has proven that sulfur is safe in that locality. During periods of high temperatures, sulfur may burn foliage and fruit. Do not make sulfur applications at excessively high temperatures. Do not use SULFUR 90W with oil or within two (2) weeks of an oil application to all crops except Citrus, for which a three (3) week interval is required. When SULFUR 90W is used with urea-nitricals, lime should be added to prevent plant injury.

Use of this material in a manner other than in accordance with this label may cause serious plant, crop or personal injury, excessive residues, unsatisfactory control or other unintended consequences.

When growing crops for processing, consult the processor before applying SULFUR 90W. Consult your state Agricultural Experiment Station or State Extension Service Specialist for complete details on the spraying program best suited to your local conditions. As a preventive measure for light infestations, use the lower dosage rates. For heavy infestations, use the higher rates.

#### APPLICATION INSTRUCTIONS

Apply through any type of hand-held, knapsack, mechanical or pneumatic spraying equipment.

**Mixing:** Pour recommended amount into partially filled spray tank. Keep agitator running during filling and spraying operation. Do not allow mixture to stand. Failure to maintain agitation will cause SULFUR 90W to settle and may necessitate manual stirring to re-disperse.

#### DILUTE APPLICATION

**Field:** Specified rate in 20 to 60 gallons of water per acre. **Orchard:** Specified rate in 100 to 800 gallons of water per acre.

#### CONCENTRATED APPLICATION

**Field:** Specified rate in 5 to 10 gallons of water per acre. **Orchard:** Specified rate in 20 to 100 gallons of water per acre.

#### AERIAL APPLICATION

**Field:** Specified rate in 3 to 20 gallons of water per acre. **Orchard:** Specified rate in 10 to 20 gallons of water per acre.

Manufactured By

**Drexel Chemical Company**

P.O. BOX 13327 MEMPHIS, TN 38113-0327

**SINCE 1972**

**FRUIT AND NUT CROPS**

**ALMONDS:** Brownrot, Powdery mildew, Leaf spot, Scab — Apply 8 to 26 lbs per acre. Apply pre-bloom, bloom, petal fall and as a cover spray every 10 to 14 days or after a period of wet weather. Mites [Brown, Almond, European red, Two-spotted, Pacific, Strawberry (Atlantic), Silver, Flat] — Apply 8 to 26 lbs per acre as needed when infestation is noted. Do not apply within two (2) weeks of an oil application.

**APPLES:** Scab, Powdery mildew — Apply 8 to 26 lbs per acre. Apply pre-bloom, bloom, petal fall, shuck and as a cover spray throughout the season. Mites [Blister, European red, Two-spotted] — Apply 8 to 26 lbs per acre as needed when infestation is noted.

**AVOCADOS:** Brown mites — Apply 8 to 26 lbs per acre as necessary.

**BUSHBERRIES** — Blueberries, Currants, Huckleberries: Powdery mildew — Apply 8 to 15 lbs per acre when shoots are 6 inches long and before blossoms open. Repeat at 10 day intervals as necessary.

**CANEFRUITS** — Blackberries, Boysenberries, Dewberries, Gooseberries, Loganberries, Raspberries: Powdery mildew — Apply 8 to 15 lbs per acre when shoots are 6 inches long and before blossoms open. Repeat at 10 day intervals as necessary.

**CHERRIES:** Brownrot, Blossom blight, Powdery mildew, Leaf spot, Rust — Apply 8 to 26 lbs per acre. Apply pre-bloom, bloom, petal fall and as a cover spray at 7 to 14 day intervals and as a post-harvest kufuu treatment for leaf spot. Mites [Silver, Flat] — Apply 8 to 26 lbs per acre as needed.

**CITRUS:** Scale crawlers, Mites [Rust, Six-spotted, Citrus, Flat, Yuma, Silver] — Apply 7 to 17 lbs per acre as needed and repeat application as necessary. Apply when pests first appear. Do not combine in spray mixture containing oil or apply for 3 weeks following an oil application.

**DATES:** Banks grass mites — Apply 8 to 26 lbs per acre as needed.

**FIGS:** Fig rust mites — Apply 8 to 26 lbs per acre as needed.

**GRAPES:** Powdery mildew, Mites [Bud, Blister, Red spider] — Apply 2 to 8 lbs per acre, preferably by ground equipment, beginning when shoots are 6 to 8 inches long and continuing at 7 to 10 day intervals as needed. Concord and other Labrusca type grapes may be injured by sulfur.

**MACADAMIAS:** Mites [Almond, Red spider, Two-spotted, Pacific] — Apply 8 to 26 lbs per acre as needed. Broad mites (Hawaii only) — Apply 8 to 26 lbs per acre as needed.

**MANGOES:** Powdery mildew — Apply 8 to 26 lbs per acre. Begin before bloom. Repeat in bloom, after fruit set and 3 weeks later or as necessary.

**OLIVES:** Olive mites — Apply 8 to 26 lbs per acre as needed.

**PAPAYA (Hawaii only):** Mites (except Carmine mites) — Apply 8 to 26 lbs per acre. Apply when mites first appear and repeat as necessary to maintain control. Do not apply in combination with emulsifiable pesticide formulations since foliar injury is likely to result. May be applied up to 24 hours before harvest. Do not apply to low vigor, moisture stressed plants. Do not apply prior to or during periods of excessively high temperatures, unseasonably low rainfall or within two weeks of an oil spray.

**PEACHES AND NECTARINES:** Brownrot, Blossom blight, Coryneum blight, Rust, Powdery mildew, Scab — Apply 8 to 26 lbs per acre pre-bloom, bloom, petal-fall and in cover sprays at 7 to 14 day intervals.

Peach silver mites — Apply 8 to 26 lbs per acre as needed.

**PEARS:** Scab, Powdery mildew — Apply 8 to 26 lbs per acre pre-bloom, bloom, petal-fall, shuck and as a cover spray throughout the season. Mites [Blister, European red, Two-spotted] — Apply 8 to 26 lbs per acre as needed. Sensitive varieties such as D'Anjou and Comice may be injured by sulfur under certain climatic conditions.

**PECANS:** Mites [Almond, Red spider, Two-spotted, Pacific, Broad] — Apply 8 to 26 lbs per acre as needed.

**PISTACHIOS:** Citrus flat mites — Apply 8 to 26 lbs per acre when mites first appear and repeat as necessary to maintain control.

**PLUMS AND PRUNES:** Brownrot, Blossom blight, Scab, Powdery mildew, Rust, Leaf spot — Apply 8 to 26 lbs per acre pre-bloom, bloom, petal-fall and as a cover spray at 7 to 10 day intervals. Silver mites — Apply 8 to 26 lbs per acre as needed.

**POMEGRANATES:** Mites — Apply 8 to 26 lbs per acre as needed.

**QUINCE:** Brownrot, Powdery mildew, Scab — Apply 8 to 26 lbs per acre. Begin application at early leaf stage and repeat as necessary. Do not use on sulfur sensitive varieties.

**STRAWBERRIES:** Powdery mildew, Red spider mites — Apply 5 to 10 lbs per acre. Begin application at early leaf stage and repeat as necessary. Do not use on sulfur sensitive varieties.

**WALNUTS:** Mites [European red, Pacific, Almond, Two-spotted] — Apply 8 to 26 lbs per acre as needed. Apply with miticide sprays to increase effectiveness. Certain varieties of walnuts are sensitive to sulfur. Do not apply unless the variety is known to be sulfur tolerant.

**VEGETABLE CROPS**

**ARTICHOKES (all varieties):** Leaf spot — Apply 8 to 26 lbs per acre. Begin when disease first appears and repeat at 7 to 10 day intervals as necessary.

**ASPARAGUS:** Rust — Apply 10 to 20 lbs per acre after cutting stops, and 7 to 10 day intervals through the season. During the fern period, apply 2 to 3 week intervals or with regular insect program. Mites [Two-spotted, Brown] — Apply 4 to 8 lbs per acre as necessary.

**BEANS AND PEAS:** Powdery mildew, Rust, Leaf spot — Apply 3 to 6 lbs per acre at first sign of infection and every 7 to 14 days as necessary. Mites [Two-spotted, Pacific, Strawberry (Atlantic)] — Apply 3 to 6 lbs per acre as necessary.

**CARROTS:** Powdery mildew — Apply 5 to 10 lbs per acre. Begin at first sign of disease and repeat at 7 to 10 day intervals as needed.

**CELERY:** Powdery mildew, Rust, Red spider mites — Apply 3 to 10 lbs per acre.

Start in early leaf stage. Repeat application every 10 to 14 days as needed.

**COLE CROPS** — Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Collards, Kale, Mustard Greens: Powdery mildew — Apply 3 to 5 lbs per acre beginning in early leaf stage and repeating every 10 to 14 days as needed, or immediately after rain. Mites [Two-spotted, Pacific, Strawberry (Atlantic)] — Apply 3 to 5 lbs per acre as necessary.

**EGGPLANTS:** Powdery mildew — Apply 3 to 5 lbs per acre. Begin when first true leaves appear. Repeat at weekly intervals as needed.

**MELONS:** Powdery mildew — Apply 15 to 20 lbs per acre. Begin when first true leaves appear and repeat at weekly intervals as needed. Some varieties are sensitive to sulfur, especially at high temperatures.

**ONIONS AND GARLIC:** Powdery mildew — Apply 5 to 10 lbs per acre. Begin before disease is expected to appear and repeat as necessary.

**PEPPERMINT AND SPEARMINT:** Powdery mildew — Apply 3 to 5 lbs per acre when mint is 5 to 6 inches tall or when disease appears. Repeat twice, at 30 day intervals. Do not apply within 30 days of harvest.

**PEPPERS:** Powdery mildew, Leaf spot — Apply 3 to 5 lbs per acre beginning in the early leaf stage and repeating every 10 to 14 days as necessary.

**POTATOES:** Powdery mildew, Leaf spot — Apply 3 to 8 lbs per acre. Apply when disease first appears and repeat at 3 to 4 week intervals as needed. Two-spotted mites — Apply 3 to 8 lbs per acre as necessary.

**RUTABAGAS:** Powdery mildew, Rust, Red spider mites — Apply 3 to 10 lbs per acre. Start in early leaf stage. Repeat application every 10 to 14 days as needed.

**SPINACH:** Powdery mildew, Rust — Apply 3 to 5 lbs per acre. Apply when disease first appears. Repeat at 7 to 10 day intervals as needed.

**TABLE BEETS:** Powdery mildew, Red spider mites — Apply 4 to 8 lbs per acre at 10 to 30 day intervals as soon as the first symptoms appear.

**TOMATOES:** Tomato russet mites — Apply 2 to 8 lbs per acre. Begin when infestation first occurs and repeat as necessary.

**TURNIPS:** Powdery mildew, Red spider mites — Apply 4 to 8 lbs per acre at early leaf stage and repeat every 14 days as necessary.

**FIELD CROPS**

**ALFALFA:** Mites — Apply 3 to 5 lbs per acre as necessary. Repeat at two week intervals as necessary.

**GLOVER:** Powdery mildew, Red spider mites — Apply 2 to 8 lbs per acre as needed.

**CORN AND SORGHUM:** Powdery mildew — Apply 3 to 5 lbs per acre. Apply when disease appears and repeat as necessary. Mites [Two-spotted, Pacific] — Apply 3 to 5 lbs per acre when infestation first appears, repeating as necessary.

**COTTON:** Mites [Strawberry (Atlantic), Red spider] — Apply 6 to 8 lbs per acre beginning when infestation first occurs and every 7 to 14 days as necessary.

**COMMON PEAS:** Rust — Apply 2 to 8 lbs per acre. Begin soon after seedlings emerge. Repeat at 7 to 10 day intervals as needed through the season.

**FLAX:** Powdery mildew — Apply 2 to 8 lbs per acre. Begin at first sign of disease. Repeat at 7 to 10 day intervals or as necessary.

**GRASS SEED CROPS:** Timothy mites — Apply 2 to 8 lbs per acre as needed.

**HOPS:** Mites — Apply 2 to 4 lbs per acre as needed for mite suppression.

**PEANUTS:** Rust, Powdery mildew, Leaf spot — Apply 3 to 5 lbs per acre beginning in early leaf stage and repeating every 10 to 24 days as necessary.

**SMALL GRAINS** — Barley, Oats, Rye, Wheat: Powdery mildew — Apply 3 to 8 lbs per acre. Apply when disease appears and repeat as necessary. Mites [Two-spotted, Pacific, Strawberry (Atlantic)] — Apply 3 to 6 lbs per acre as needed.

**SOYBEANS:** Powdery mildew — Apply 3 to 8 lbs per acre. Apply when disease appears and repeat as necessary. Mites [Two-spotted, Pacific, Strawberry (Atlantic)] — Apply 3 to 8 lbs per acre beginning when infestation is first noted, repeating as necessary.

**SUGAR BEETS:** Powdery mildew, Red spider mites — Apply 3 to 15 lbs per acre beginning when disease first appears, repeating at 3 to 4 week intervals as needed.

**VEGETABLES:** Rust — Apply 2 to 8 lbs per acre. Begin at first sign of disease. Repeat at 7 to 10 day intervals as needed.

**FLOWERS AND ORNAMENTALS**

**APPLICATION:** To control Red spider mites and the diseases listed below, use SULFUR 90W as a spray applying 2 to 8 lbs per acre by ground or 2 to 17 lbs by air. Applications should be made at 5 to 6 day intervals, beginning when the disease or mites first appear and continuing until all danger from the disease or from mites is past. During periods of humid or rainy weather it may be necessary to make applications as often as every 2 to 3 days.

FLOWERS AND ORNAMENTALS	CONTROLS
Aster, Chrysanthemum, Cosmos, Dahlia, Daisy, Delphinium, Dogwood, Eucalyptus, Golden Plectra, Hibiscus, Ironsuckle, Hydrangea, Lady's mantle, Ligustrum, Lilac, Crepe Myrtle, Phlox, Poplar, Silver vine, Rose, Spruce, Sunflower, Verbena, Willow, Zinnia	Powdery mildew
Arbutus, Carnation, Cedar, Chrysanthemum, Clematis, Hydrangea, Rose, Snapdragon, Sweetpeas	Red spider mites
Calendula, Cherry Laurel, Clematis, Columbine, English ivy, Foxglove, Hydrangea, Peltandra, Phlox, Sage, Simulax, Snapdragon	Leaf spot
Hydrangea	Botrytis blight
Rose	Black spot

4/4

**HOME AND GARDEN USE DIRECT**

**APPLICATION:** To control the diseases and mites listed below, use SULFUR 90W as a spray at a rate of 2 to 4 tablespoons per gallon of water or as a dust at a rate of 16 ounces per 1000 square feet. Applications should begin when the disease or mites first appear and continue at 7 to 14 day intervals until all danger from the disease or mites is past. During periods of humid or rainy weather, it may be necessary to make applications as often as every 2 to 3 days. Make full coverage applications on both sides of leaves. Apply to plant foliage, stems, and flowers to leave a 'barely visible' dust. Do not over apply during full sun nor during times when temperatures exceed 90° F.

**PRECAUTIONS:** Do not apply within 2 weeks of an oil application and do not apply oil for two weeks (three weeks for citrus) after an application of SULFUR 90W. Injury may occur under adverse climatic conditions. Some varieties of some vegetables, fruit trees and ornamentals are sensitive to sulfur. Consult State Agricultural Experiment Station or the State Agricultural Extension Service for sensitive plants and additional information as the timing, number, and rate of application needed will vary with local conditions.

FLOWERS AND ORNAMENTALS	CONTROLS
Alder, Azalea, Buttonbush, Catalpa, Cosmos, Dahlia, Daisy, Delphinium, Dogwood, Euonymus, Golden Fleece, Hibiscus, Honeysuckle, Horse Chestnut, Lady's Mantle, Ligustrum, Lilac, Linden, Matrimony Vine, Crepe Myrtle, Poplar, Rhododendron, Staghorn, Sumac, Silver Vine, Spiraea, Sweetpea, Sunflower, Sycamore, Turnpot Vine, Verbena, Willow, Witch Hazel, Zinnia	Powdery mildew
Arborvitae, Cedar, Sweetgum	Red spider mites
Aster	Powdery mildew, Rust
Calendula, Cherry Laurel, Columbine, English Ivy, Foxglove, Petunia, Sage, Smilax	Leaf spot
Carnation	Red spider mites, Rust
Chrysanthemum	Powdery mildew, Red spider mites, Rust
Clematis	Leaf spot, Red spider mites
Hydrangea	Botrytis blight, Leaf spot, Powdery mildew, Red spider mites
Phlox	Leaf spot, Powdery mildew
Rose	Black spot, Powdery mildew, Red spider mites, Rust
Snap Dragon	Leaf spot, Red spider mite, Rust

FRUITS AND NUTS	CONTROLS
Almonds	Brownrot, Leaf spot, Mites, Powdery mildew, Scab
Apples, Pears	Mites, Powdery mildew, Scab
Avocados	Mites
Bushberries (Blueberries, Currants, Huckleberries)	Powdery mildew
Caneberries (Blackberries, Boysenberries, Dewberries, Gooseberries, Loganberries, Raspberries)	Powdery mildew
Cherries	Brownrot, Blossom blight, Leaf spot, Mites, Powdery mildew, Rust
Citrus, Figs, Macadamias, Olives, Pecans, Pistachos, Pome Granates, Walnuts	Mites
Grapes	Mites, Powdery mildew
Mangoes, Strawberries	Powdery mildew
Nectarines, Peaches	Brownrot, Blossom blight, Coryneum blight, Mites, Powdery mildew, Rust, Scab
Plums, Prunes	Brownrot, Blossom blight, Leaf spot, Mites, Powdery mildew, Rust, Scab
Quince	Brownrot, Blossom blight, Powdery mildew

VEGETABLES	CONTROLS
Artichokes	Leaf spot
Asparagus	Rust, Mites
Beans, Peas	Leaf spot, Mites, Powdery mildew, Rust
Carrots, Eggplants, Garlic, Melons, Onions, Peppermint, Spearmint	Powdery mildew
Celery, Rutabagas	Mites, Powdery mildew, Rust
Cole Crops (Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Collards, Kale, Mustard Greens)	Mites, Powdery mildew
Peppers	Leaf spot, Powdery mildew
Potatoes	Leaf spot, Mites, Powdery mildew
Spinach	Powdery mildew, Rust
Table Beets	Mites, Powdery mildew
Tomatoes	Mites
Turnips	Mites, Powdery mildew

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**STORAGE INSTRUCTIONS:** Storage should be under lock and key and secure from access by unauthorized persons and children. Storage should be in a cool dry area, away from any heat or ignition source. Do not stack over 2 pallets high. Move bags carefully so as not to tear or puncture. Do not move containers from one area to another unless they are securely sealed. Keep container tightly sealed when not in use. Do not allow bags to become wet or to be stored in a damp humid area. Keep away from any puncture source. Avoid storage near water supplies, food, feed, and fertilizer to avoid contamination. Store in original container only. If the contents are leaking or material is spilled, follow these steps:

1. Collect and place in suitable containers for disposal.
2. Wash area with water and soap to remove remaining pesticides.
3. Follow washing with clean water rinse.
4. Do not allow run-off to enter sewer or contaminate water supplies.
5. Dispose of waste as indicated below.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment, then dispose of bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**WARRANTY - CONDITION OF SALE**

OUR RECOMMENDATIONS FOR USE of this product are based upon tests believed reliable. Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the Seller. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with the directions given herewith. In no case shall Drexel or the Seller be liable for consequential, special or indirect damages when such use and/or handling is not in strict accordance with directions given herewith. The foregoing is a condition of sale by Drexel Chemical Company and is accepted as such by the Buyer.