

19713-238

11-12-2002

1/8

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0080. Approval expires 2-28-95



United States
Environmental Protection Agency
Washington, DC 20460

Registration
 Amendment
 Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 19713-238	2. EPA Product Manager MARY L. WALLER	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) DREXEL SULFUR 90W	PM# 21/Fungicide Branch	
5. Name and Address of Applicant (Include ZIP Code) Drexel Chemical Company, P.O. Box 13327 MEMPHIS, TN 38113-0327 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input checked="" type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

NOTIFICATION
NOV 12 2002

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

One (1) copy of the label (238MSP-1002++) revised per PR Notice 2001-6. A copy of the required certification statement is also submitted. Thank you.

Section - III

1. Material This Product Will Be Packaged in:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	If "Yes" Unit Packaging wgt. No. per container		<input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
* Certification must be submitted		If "Yes" Package wgt. No. per container			
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name LUZ G PIWONKA	Title REGISTRATION MANAGER	Telephone No. (include Area Code) (901) 774-4370
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature <i>Luza Piwonka</i>	3. Title REGISTRATION MANAGER	
4. Typed Name LUZ G PIWONKA	5. Date November 6, 2002	

2/8



Drexel Chemical Company


November 6, 2002

Submission of Revised Label per PR Notice 2001-6
DREXEL SULFUR 90W (EPA Reg. No. 19713-238)

This notification is consistent with the Provisions of PR Notice 2001-6 and EPA Regulations at 40 CFR 156.10 and CFR 152.46, and no other changes have been made to the labeling or the Confidential Statement of Formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the guidance of PR Notice 2001-6 and the requirements of 40 CFR 152.10 and 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under Sections 12 and 14 of FIFRA.

FOR DREXEL CHEMICAL COMPANY

NOTIFICATION
NOV 12 2002



LUZ G PIWONKA
Registration Manager

[Faint, illegible text, possibly a stamp or bleed-through]

3/8


MASTER LABEL

Note to EPA

Shaded text added for water-soluble package.



Tear Open Outer Bag to Obtain the Water-Soluble Pak.



Sulfur 90W

Micronized Sulfur

Water-Soluble Packaging

ACTIVE INGREDIENT:

Elemental sulfur	90.0%
OTHER INGREDIENTS:	10.0%
TOTAL:	100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See FIRST AID Below

EPA Reg. No. 19713-238
EPA Est. No. 19713-GA-1

Net Contents:
20 Pounds

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water when disposing of equipment washwaters.

PHYSICAL HAZARDS

Keep away from heat, sparks or flame.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

FIRST AID

IF SWALLOWED:

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything by mouth to an unconscious or convulsing person.

IF IN EYES:

- Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15 to 20 minutes.

Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

PRECAUTIONARY STATEMENTS

Hazards To Humans and Domestic Animals

CAUTION: Causes eye and skin irritation. Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing of dust or spray mist. Avoid contact with eyes, skin or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category A on an EPA chemical-resistance category selection chart. **Applicators and other handlers must wear:** Long-sleeved shirt and long pants, chemical-resistant gloves such as polyethylene or polyvinyl chloride, shoes plus socks and protective eyewear. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should: 1) Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. 2) Remove clothing immediately if pesticides get inside. 3) Keep and wash PPE separately from other laundry.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard (WPS), 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval (REI). The requirements in this box only apply to uses of this product that are covered by the WPS. Do not enter or allow worker entry into treated areas during the REI of 24 hours.

PPE required for early entry to treated areas that is permitted under the WPS and that involves contact with anything that has been treated, such as plants, soil or water is: Coveralls, chemical-resistant gloves such as polyethylene or polyvinyl chloride, shoes plus socks and protective eyewear.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the WPS for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses. Keep children and domestic animals off treated areas until this material has been washed into the soil.



Manufactured By:
Drexel Chemical Company

P.O. BOX 13327, MEMPHIS, TN 38113-0327

SINCE 1972

NOTIFICATION

NOV 12 2002

4/8

DIRECTIONS FOR ORIGINAL FORMULATION

Pour needed amount into spray tank under agitation. If necessary, add a few drops of oil to avoid foaming. Application of this material through any type of irrigation equipment is prohibited.

DIRECTIONS FOR WATER-SOLUBLE PACKAGING

Paks containing this product are water-soluble. Do not allow pak to become wet prior to adding to the spray tank. Do not handle paks with wet hands. To prepare the spray mixture, drop the required number of unopened paks into spray tank while filling with the appropriate amount of water. Operate the agitator while mixing. Depending on the water temperature and the degree of agitation, the paks should be completely dissolved in approximately 5 minutes from the time they are added to the water. Use the specified dosage of this product in the amount of water necessary to give complete coverage. Determine the total amount to be added to the spray tank based on the rates in the "Crop" section.

Note: Products containing Boron and/or fertilizers will prevent the water-soluble packet from dissolving. Boron and/or fertilizer products must only be added after packet(s) have completely dissolved.

USE PRECAUTIONS

IMPORTANT: Some fruits and other plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur on any crop unless local use has proven that sulfur is safe in that locality. During periods of high temperatures, sulfur may burn foliage and fruit. Do not make sulfur applications at excessively high temperatures. Do not use this product with oil or within two weeks of an oil application to all crops except Citrus, for which a three-week interval is required. When this product is used with arsenicals, lime should be added to prevent plant injury.

Use of this material in a manner other than in accordance with this label may cause serious plant, crop or personal injury, excessive residues, unsatisfactory control or other unintended consequences. When growing crops for processing, consult the processor before applying this product. Consult your State Agricultural Experiment Station or State Extension Service Specialist for complete details on the spraying program best suited to your local conditions. As a preventive measure for light infestations, use the lower dosage rates. For heavy infestations, use the higher rates.

APPLICATION INSTRUCTIONS

Apply through any type of hand-held, knapsack, mechanical or pneumatic spraying equipment.

Mixing: Pour recommended amount into partially filled spray tank. Keep agitator running during filling and spraying operation. Do not allow mixture to stand. Failure to maintain agitation will cause this product to settle and may necessitate manual stirring to re-disperse.

DILUTE APPLICATION

Field: Specified rate in 20 to 60 gallons of water per acre.

Orchard: Specified rate in 100 to 800 gallons of water per acre.

CONCENTRATED APPLICATION

Field: Specified rate in 5 to 10 gallons of water per acre.

Orchard: Specified rate in 20 to 100 gallons of water per acre.

AERIAL APPLICATION

Field: Specified rate in 3 to 20 gallons of water per acre.

Orchard: Specified rate in 10 to 20 gallons of water per acre.

FRUIT AND NUT CROPS

Crop	Pest/Disease	Rate per Acre	Number of Water-Soluble Paks
Almonds	Brown rot, Leaf spot, Powdery mildew, Scab	8 to 26 lbs.	4 to 13 per 10 acres
	TIMING/SPECIFIC DIRECTIONS: Apply pre-bloom, bloom, petal fall and as a cover spray every 10 to 14 days or after a period of wet weather.		
	Mites (Almond, Brown, European red, Flat, Pacific, Silver, Strawberry [Atlantic], Twospotted)	8 to 26 lbs.	4 to 13 per 10 acres
TIMING/SPECIFIC DIRECTIONS: Apply as needed when infestation is noted. Do not apply within 2 weeks of an oil application.			
Apples	Powdery mildew, Scab	8 to 26 lbs.	4 to 13 per 10 acres
	TIMING/SPECIFIC DIRECTIONS: Apply pre-bloom, bloom, petal fall, shuck and as a cover spray throughout the season.		
	Mites (Blister, European red, Twospotted)	8 to 26 lbs.	4 to 13 per 10 acres
TIMING/SPECIFIC DIRECTIONS: Apply as needed when infestation is noted.			
Avocados	Brown mites	8 to 26 lbs.	4 to 13 per 10 acres
	TIMING/SPECIFIC DIRECTIONS: Apply as needed.		

FRUIT AND NUT CROPS (Cont.)

Crop	Pest/Disease	Rate per Acre	Number of Water-Soluble Paks
Bushberries (Blueberries, Currants, Huckleberries)	Powdery mildew	6 to 15 lbs.	6 to 15 per 20 acres
	TIMING/SPECIFIC DIRECTIONS: Apply when shoots are 6 inches long and before blossoms open. Repeat at 10-day intervals as necessary.		
Canefruits (Blackberries, Boysenberries, Dewberries, Gooseberries, Loganberries, Raspberries)	Powdery mildew	6 to 15 lbs.	6 to 15 per 20 acres
	TIMING/SPECIFIC DIRECTIONS: Apply when shoots are 6 inches long and before blossoms open. Repeat at 10-day intervals as necessary.		
Cherries	Brown rot blossom blight, Leaf spot, Powdery mildew, Rust	8 to 26 lbs.	4 to 13 per 10 acres
	TIMING/SPECIFIC DIRECTIONS: Apply pre-bloom, bloom, petal fall and as a cover spray at 7- to 14-day intervals as a post-harvest foliar treatment for Leaf spot.		
	Mites (Flat, Silver)	8 to 26 lbs.	4 to 13 per 10 acres
TIMING/SPECIFIC DIRECTIONS: Apply as needed.			
Citrus	Mites (Citrus, Flat, Silver, Six-spotted, Yuma), Scale crawlers	7 to 17 lbs.	7 to 17 per 20 acres
	TIMING/SPECIFIC DIRECTIONS: Apply as needed and repeat application as necessary. Apply when pests first appear. Do not combine in spray mixture containing oil or apply for 3 weeks following an oil application.		
Dates	Banks grass mites	8 to 26 lbs.	4 to 13 per 10 acres
	TIMING/SPECIFIC DIRECTIONS: Apply as needed.		
Figs	Fig rust mites	8 to 26 lbs.	4 to 13 per 10 acres
	TIMING/SPECIFIC DIRECTIONS: Apply as needed.		
Grapes	Mites (Blister, Bud, Red spider), Powdery mildew	2 to 8 lbs.	1 to 4 per 10 acres
	TIMING/SPECIFIC DIRECTIONS: Apply, preferably by ground equipment, beginning when shoots are 6 to 8 inches long and continuing at 7- to 10-day intervals as needed. Concord and other Labrusa type Grapes may be injured by sulfur.		
Macadamia nuts	Mites (Almond, Pacific, Red spider, Twospotted)	8 to 26 lbs.	4 to 13 per 10 acres
	Broad mites (HI Only)	TIMING/SPECIFIC DIRECTIONS: Apply as needed.	
Mangoes	Powdery mildew	8 to 26 lbs.	4 to 13 per 10 acres
	TIMING/SPECIFIC DIRECTIONS: Begin before bloom. Repeat in bloom, after fruit set and 3 weeks later or as necessary.		
Olives	Olive mites	8 to 26 lbs.	4 to 13 per 10 acres
	TIMING/SPECIFIC DIRECTIONS: Apply as needed.		
Papaya (HI)	Mites (except Carmine mites)	8 to 26 lbs.	4 to 13 per 10 acres
	TIMING/SPECIFIC DIRECTIONS: Apply when Mites first appear and repeat as necessary to maintain control. Do not apply in combination with emulsifiable pesticide formulations since foliar injury is likely to result. May be applied up to 24 hours before harvest. Do not apply to low vigor, moisture-stressed plants. Do not apply prior to or during periods of excessively high temperature, unseasonably low rainfall or within 2 weeks of an oil spray.		
Peaches, Nectarines	Brown rot blossom blight, Coryneum blight, Powdery mildew, Rust, Scab	8 to 26 lbs.	4 to 13 per 10 acres
	TIMING/SPECIFIC DIRECTIONS: Apply pre-bloom, bloom, petal fall and in cover spray at 7- to 14-day intervals.		
	Peach silver mites	8 to 26 lbs.	4 to 13 per 10 acres
TIMING/SPECIFIC DIRECTIONS: Apply as needed.			

5/8

FRUIT AND NUT CROPS (Cont.)

Crop	Pest/Disease	Rate per Acre	Number of Water-Soluble Paks
Pears	Powdery mildew, Scab	8 to 26 lbs.	4 to 13 per 10 acres
	TIMING/SPECIFIC DIRECTIONS: Apply pre-bloom, bloom, petal fall, shuck and as a cover spray throughout the season.		
	Mites (Blister, European red, Twospotted)	8 to 26 lbs.	4 to 13 per 10 acres
	TIMING/SPECIFIC DIRECTIONS: Apply as needed. Sensitive varieties such as Comice and D'anjou may be injured by sulfur under certain climatic conditions.		
Pecans	Mites (Almond, Broad, Pacific, Red spider, Twospotted)	8 to 26 lbs.	4 to 13 per 10 acres
TIMING/SPECIFIC DIRECTIONS: Apply as needed.			
Pistachios	Citrus flat mites	8 to 26 lbs.	4 to 13 per 10 acres
	TIMING/SPECIFIC DIRECTIONS: Apply when Mites first appear and repeat as necessary to maintain control.		
Plums, Prunes	Brown rot blossom blight, Leaf spot, Powdery mildew, Rust, Scab	8 to 26 lbs.	4 to 13 per 10 acres
	TIMING/SPECIFIC DIRECTIONS: Apply pre-bloom, bloom, petal fall and as a cover spray at 7- to 10-day intervals.		
	Silver mites	8 to 26 lbs.	4 to 13 per 10 acres
TIMING/SPECIFIC DIRECTIONS: Apply as needed.			
Pomegranates	Mites	8 to 26 lbs.	4 to 13 per 10 acres
	TIMING/SPECIFIC DIRECTIONS: Apply as needed.		
Quince	Brown rot, Powdery mildew, Scab	8 to 26 lbs.	4 to 13 per 10 acres
	TIMING/SPECIFIC DIRECTIONS: Begin application at early leaf stage and repeat as necessary. Do not use on sulfur-sensitive varieties.		
Strawberries	Powdery mildew, Red spider mites	5 to 10 lbs.	1 to 2 per 4 acres
	TIMING/SPECIFIC DIRECTIONS: Begin application at early leaf stage and repeat as necessary. Do not use on sulfur-sensitive varieties.		
Walnuts	Mites (Almond, European red, Pacific, Twospotted)	8 to 26 lbs.	4 to 13 per 10 acres
	TIMING/SPECIFIC DIRECTIONS: Apply as needed. Apply with miticide sprays to increase effectiveness. Certain varieties of Walnuts are sensitive to sulfur. Do not apply unless the variety is known to be tolerant to sulfur.		

VEGETABLE CROPS

Crop	Pest/Disease	Rate per Acre	Number of Water-Soluble Paks
Artichokes (All varieties)	Leaf spot	8 to 26 lbs.	4 to 13 per 10 acres
	TIMING/SPECIFIC DIRECTIONS: Begin applications when disease first appears and repeat at 7- to 10-day intervals as necessary.		
Asparagus	Rust	10 to 20 lbs.	1 to 2 per 2 acres
	TIMING/SPECIFIC DIRECTIONS: Apply after cutting stops and at 7- to 10-day intervals throughout the season. During the fern period, apply at 2- to 3-week intervals or with regular insect program.		
	Mites (Brown, Twospotted)	4 to 6 lbs.	2 to 3 per 10 acres
	TIMING/SPECIFIC DIRECTIONS: Apply as necessary.		
Beans, Peas	Leaf spot, Powdery mildew, Rust	3 to 6 lbs.	3 to 6 per 20 acres
	TIMING/SPECIFIC DIRECTIONS: Apply at first sign of infection and every 7 to 14 days as necessary.		
	Mites (Pacific, Strawberry [Atlantic], Twospotted)	3 to 6 lbs.	3 to 6 per 20 acres
TIMING/SPECIFIC DIRECTIONS: Apply as necessary.			
Carrots	Powdery mildew	5 to 10 lbs.	1 to 2 per 4 acres
	TIMING/SPECIFIC DIRECTIONS: Begin applications at first sign of disease and repeat at 7- to 10-day intervals.		

(Continued)

VEGETABLE CROPS (Cont.)

Crop	Pest/Disease	Rate per Acre	Number of Water-Soluble Paks
Celery	Powdery mildew, Red spider mites, Rust	3 to 10 lbs.	3 to 10 per 20 acres
	TIMING/SPECIFIC DIRECTIONS: Begin applications in early leaf stage. Repeat every 10 to 14 days as needed.		
Cole crops (Broccoli, Brussels sprouts, Cabbage, Cauliflower, Collards, Kale, Mustard greens)	Powdery mildew	3 to 5 lbs.	3 to 5 per 20 acres
	TIMING/SPECIFIC DIRECTIONS: Apply beginning in early leaf stage and repeating every 10 to 14 days as needed or immediately after rain.		
	Mites (Pacific, Strawberry [Atlantic], Twospotted)	3 to 5 lbs.	3 to 5 per 20 acres
TIMING/SPECIFIC DIRECTIONS: Apply as necessary.			
Eggplants	Powdery mildew	3 to 5 lbs.	3 to 5 per 20 acres
	TIMING/SPECIFIC DIRECTIONS: Begin applications when first true leaves appear. Repeat at weekly intervals as needed.		
Melons	Powdery mildew	15 to 20 lbs.	3 to 4 per 4 acres
	TIMING/SPECIFIC DIRECTIONS: Begin applications when first true leaves appear and repeat at weekly intervals as needed. Some varieties are sensitive to sulfur, especially at high temperatures.		
Onions, Garlic	Powdery mildew	5 to 10 lbs.	1 to 2 per 4 acres
	TIMING/SPECIFIC DIRECTIONS: Begin applications before disease is expected to appear and repeat as necessary.		
Peppermint, Spearmint	Powdery mildew	3 to 5 lbs.	3 to 5 per 20 acres
	TIMING/SPECIFIC DIRECTIONS: Apply when Mint is 5 to 6 inches tall or when disease appears. Repeat twice, at 30-day intervals. Do not apply within 30 days of harvest.		
Peppers	Leaf spot, Powdery mildew	3 to 5 lbs.	3 to 5 per 20 acres
	TIMING/SPECIFIC DIRECTIONS: Begin applications at early leaf stage and repeat every 10 to 14 days as necessary.		
Potatoes	Leaf spot, Powdery mildew	3 to 8 lbs.	3 to 8 per 20 acres
	TIMING/SPECIFIC DIRECTIONS: Apply when disease first appears and repeat at 3- to 4-week intervals as needed.		
	Twospotted mites	3 to 8 lbs.	3 to 8 per 20 acres
	TIMING/SPECIFIC DIRECTIONS: Apply as necessary.		
Rutabagas	Powdery mildew, Red spider mites, Rust	3 to 10 lbs.	3 to 10 per 20 acres
	TIMING/SPECIFIC DIRECTIONS: Begin applications in early leaf stage. Repeat every 10 to 14 days as needed.		
Spinach	Powdery mildew, Rust	3 to 5 lbs.	3 to 5 per 20 acres
	TIMING/SPECIFIC DIRECTIONS: Apply when disease first appears. Repeat at 7- to 10-day intervals as needed.		
Table beets	Powdery mildew, Red spider mites	4 to 8 lbs.	1 to 2 per 5 acres
	TIMING/SPECIFIC DIRECTIONS: Begin applications at 10- to 30-day intervals as soon as the first symptoms appear.		
Tomatoes	Tomato russet mites	2 to 8 lbs.	1 to 4 per 10 acres
	TIMING/SPECIFIC DIRECTIONS: Begin applications when infestation first occurs and repeat as necessary.		
Turnips	Powdery mildew, Red spider mites	4 to 8 lbs.	1 to 2 per 5 acres
	TIMING/SPECIFIC DIRECTIONS: Apply at early leaf stage and repeat every 14 days as necessary.		

6/8

FIELD CROPS

Crop	Pest/Disease	Rate per Acre	Number of Water-Soluble Paks
Alfalfa	Mites	3 to 5 lbs.	3 to 5 per 20 acres
	TIMING/SPECIFIC DIRECTIONS: Apply as necessary. Repeat at 2-week intervals as necessary.		
Clover	Powdery mildew, Red spider mites	2 to 8 lbs.	1 to 4 per 10 acres
	TIMING/SPECIFIC DIRECTIONS: Apply as necessary.		
Corn, Sorghum	Powdery mildew	3 to 5 lbs.	3 to 5 per 20 acres
	TIMING/SPECIFIC DIRECTIONS: Apply when disease appears and repeat as necessary.		
	Mites (Pacific, Twospotted)	3 to 5 lbs.	3 to 5 per 20 acres
	TIMING/SPECIFIC DIRECTIONS: Apply when infestation first appears, repeating as necessary.		
Cotton	Mites (Red spider, Strawberry (Atlantic))	6 to 9 lbs.	6 to 9 per 20 acres
	TIMING/SPECIFIC DIRECTIONS: Apply beginning when infestation first occurs and every 7 to 14 days as necessary.		
Cowpeas	Rust	2 to 8 lbs.	1 to 4 per 10 acres
	TIMING/SPECIFIC DIRECTIONS: Begin applications soon after seedlings emerge. Repeat at 7- to 10-day intervals as needed throughout the season.		
Flax	Powdery mildew	2 to 8 lbs.	1 to 4 per 10 acres
	TIMING/SPECIFIC DIRECTIONS: Begin applications at first sign of disease. Repeat at 7- to 10-day intervals or as necessary.		
Grass seed crops	Timothy mites	2 to 8 lbs.	1 to 4 per 10 acres
	TIMING/SPECIFIC DIRECTIONS: Apply as needed.		
Hops	Mites	2 to 4 lbs.	1 to 2 per 10 acres
	TIMING/SPECIFIC DIRECTIONS: Apply as needed for Mite suppression.		
Peanuts	Leaf spot, Powdery mildew, Rust	3 to 5 lbs.	3 to 5 per 20 acres
	TIMING/SPECIFIC DIRECTIONS: Apply beginning in early leaf stage and repeating every 10 to 24 days as necessary.		
Small grains (Barley, Oats, Rye, Wheat)	Powdery mildew	3 to 8 lbs.	3 to 8 per 20 acres
	TIMING/SPECIFIC DIRECTIONS: Apply when disease appears and repeat as necessary.		
	Mites (Pacific, Strawberry (Atlantic))	3 to 8 lbs.	3 to 8 per 20 acres
	TIMING/SPECIFIC DIRECTIONS: Apply as needed.		
Soybeans	Powdery mildew	3 to 8 lbs.	3 to 8 per 20 acres
	TIMING/SPECIFIC DIRECTIONS: Apply when disease appears and repeat as necessary.		
	Mites (Pacific, Strawberry (Atlantic), Twospotted)	3 to 8 lbs.	3 to 8 per 20 acres
	TIMING/SPECIFIC DIRECTIONS: Apply beginning when infestation is first noted, repeating as necessary.		
Sugar beets	Powdery mildew, Red spider mites	3 to 15 lbs.	3 to 15 per 20 acres
	TIMING/SPECIFIC DIRECTIONS: Apply beginning when disease first appears, repeating at 3- to 4-week intervals as necessary.		
Vetch	Rust	2 to 8 lbs.	1 to 4 per 10 acres
	TIMING/SPECIFIC DIRECTIONS: Begin at first sign of disease. Repeat at 7- to 10-day intervals as needed.		

SPECIALTY CROPS

Crop	Pest/Disease	Rate per Acre	Number of Water-Soluble Paks
Kava (Awa, Kawa)	Carmine spider mites	8 to 20 lbs.	2 to 5 per 5 acres
	TIMING/SPECIFIC DIRECTIONS: Apply in sufficient amount of water for airblast sprayers to thoroughly cover all leaf surfaces. As a broadcast or directed spray, apply in 100 gals. of water per acre making sure to achieve complete coverage of upper and lower leaf surfaces. For severe infestations, apply at 7- to 10-day intervals. For control and prevention, apply at 14- to 21-day intervals.		

FLOWERS AND ORNAMENTALS

APPLICATION: To control Red spider mites and the diseases listed below, use this product as a spray, applying 2 to 8 pounds per acre by ground (1 to 4 water-soluble paks per 10 acres) or 2 to 17 pounds by air (2 to 17 water-soluble paks per 20 acres). Applications should be made at 5- to 6-day intervals, beginning when the disease or Mites first appear and continuing until all danger from the disease or from Mites has passed. During periods of humid or rainy weather it may be necessary to make applications as often as every 2 to 3 days.

FLOWERS AND ORNAMENTALS	CONTROLS
Aster, Chrysanthemum, Cosmos, Dahlia, Daisy, Delphinium, Dogwood, Euonymus, Golden fleece, Hibiscus, Honeysuckle, Hydrangea, Lady's mantle, Ligustrum, Lilac, Crepe myrtle, Phlox, Poplar, Rose, Silver vine, Spiraea, Sunflower, Verbena, Willow, Zinnia	Powdery mildew
Arborvitae, Camellia, Cedars, Chrysanthemums, Clematis, Hydrangea, Rose, Snapdragon, Sweetpeas	Red spider mites
Calendula, Cherry laurel, Clematis, Columbine, English ivy, Foxglove, Hydrangea, Petunia, Phlox, Sage, Similax, Snapdragon	Leaf spot
Hydrangea	Botrytis blight
Rose	Black spot

SPECIAL NOTE: THE FOLLOWING HOME AND GARDEN USES WILL NOT APPEAR ON WATER-SOLUBLE PACKAGING DUE TO PACKAGE SIZE.

HOME AND GARDEN USE DIRECTIONS

APPLICATION: To control the diseases and Mites listed below, use this product as a spray at a rate of 2 to 4 tablespoons per gallon of water or as a dust at a rate of 16 ounces per 1,000 square feet. Applications should begin when the disease or Mites first appear and continue at 7- to 14-day intervals until all danger from the disease or Mites has passed. During periods of humid or rainy weather, it may be necessary to make applications as often as every 2 to 3 days. Make full coverage applications on both sides of leaves. Apply to plant foliage, stems and flowers to leave a "barely visible" dust. Do not over-apply during full sun nor during times when temperatures exceed 90°F.

USE PRECAUTIONS: Do not apply within 2 weeks of an oil application and do not apply oil for two weeks (three weeks for Citrus) after an application of this product. Injury may occur under adverse climatic conditions. Some varieties of some vegetables, fruit trees and ornamentals are sensitive to sulfur. Consult State Agricultural Experiment Station or the State Agricultural Extension Service for sensitive plants and additional information as the timing, number and rate of application needed will vary with local conditions.

FLOWERS AND ORNAMENTALS	CONTROLS
Alder, Azalea, Buttonbush, Catalpa, Cosmos, Dahlia, Daisy, Delphinium, Dogwood, Euonymus, Golden fleece, Hibiscus, Honeysuckle, Horse chestnut, Lady's mantle, Ligustrum, Lilac, Linden, Matrimony vine, Crepe myrtle, Poplar, Rhododendron, Staghorn sumac, Silver vine, Spiraea, Sweetpea, Sunflower, Sycamore, Trumpetvine, Verbena, Willow, Witch hazel, Zinnia	Powdery mildew
Arborvitae, Cedar, Sweetpea	Red spider mites
Aster	Powdery mildew, Rust
Calendula, Cherry laurel, Columbine, English ivy, Foxglove, Petunia, Sage, Similax	Leaf spot
Camellia	Red spider mites, Rust
Chrysanthemum	Powdery mildew, Red spider mites, Rust
Clematis	Leaf spot, Red spider mites

(Continued)

(Continued)

FLOWERS AND ORNAMENTALS	CONTROLS
Hydrangea	Botrytis blight, Leaf spot, Powdery mildew, Red spider mites
Phlox	Leaf spot, Powdery mildew
Rose	Black spot, Powdery mildew, Red spider mites, Rust
Snapdragon	Leaf spot, Red spider mites, Rust

FRUITS AND NUTS	CONTROLS
Almonds	Brown rot, Leaf spot, Mites, Powdery mildew, Scab
Apples, Pears	Mites, Powdery mildew, Scab
Avocados	Mites
Bushberries (Blueberries, Currants, Huckleberries)	Powdery mildew
Canefruits (Blackberries, Boysenberries, Dewberries, Gooseberries, Loganberries, Raspberries)	Powdery mildew
Cherries	Brown rot blossom blight, Leaf spot, Mites, Powdery mildew, Rust
Citrus, Figs, Macadamias, Olives, Pecans, Pistachios, Pomegranates, Walnuts	Mites
Grapes	Mites, Powdery mildew
Mangoes, Strawberries	Powdery mildew
Nectarines, Peaches	Brown rot blossom blight, Coryneum blight, Mites, Powdery mildew, Rust, Scab
Plums, Prunes	Brown rot blossom blight, Leaf spot, Mites, Powdery mildew, Rust, Scab
Quince	Brown rot blossom blight, Powdery mildew

VEGETABLES	CONTROLS
Artichokes	Leaf spot
Asparagus	Rust, Mites
Beans, Peas	Leaf spot, Mites, Powdery mildew, Rust
Carrots, Eggplants, Garlic, Melons, Onions, Peppermint, Spearmint	Powdery mildew
Celery, Rutabagas	Mites, Powdery mildew, Rust
Cole crops such as (Broccoli, Brussels sprouts, Cabbage, Cauliflower, Collards, Kale, Mustard greens)	Mites, Powdery mildew
Peppers	Leaf spot, Powdery mildew
Potatoes	Leaf spot, Mites, Powdery mildew
Spinach	Powdery mildew, Rust
Table beets	Mites, Powdery mildew
Tomatoes	Mites
Turnips	Mites, Powdery mildew

GREENHOUSE USE

CROP/SITE: To be applied to the listed crops grown in greenhouse. See crop list below.

Arugula, African yam bean, Allspice, Aloe, Amaranth, Asparagus, Bambara groundnut, Banana, Banana pink jumbo squash, Barley, Beans, Beets, Benzo leaf, Bottle gourd, Broad-bean, Broccoli, Brussels sprouts, Buffalo gourd, Butter bean, Cabbage, Cacao, Cantaloupe, Caraway, Carrots, Cassava, Cauliflower, Celery, Chaya, Chayote, Chervil, Chicory, Chinese cabbage, Chives, Cilantro, Coconut palm, Cocoyam, Coffee, Coriander, Corn, Cotton, Cowpeas, Cucumber, Date palm, Dill, Eggplant, Endive, Fava beans, Flax, Gamma grass, Garden bean, Garlic, Globe artichoke, Golden barrel cactus, Gopher plant, Grape, Grapefruit tree, Green bean, Groundnut, Guar, Guayule, Halophytes, Jerusalem artichoke, Jicama, Jojoba, Kale, Kenaf, Kohlrabi, Lablab bean, Lavendar, Leek, Lemon balm, Lemon tree, Lentils, Lettuce, Leucaena, Lima bean, Lime tree, Luffa gourd, Lupine, Malabar spinach, Marama bean, Marigold, Mesquite, Millet, Mint, Moth bean, Mung bean, Mushrooms, Mustard greens, Nasturtium, Netted melon, Oats,

(Continued)

(Continued)

Ocotillo, Okra, Onion, Orange tree, Oregano, Pak-Choi, Palmer's grass, Panic grass, Papaya, Parsley, Passion fruit, Peach palm, Peanuts, Peas, Pepper, Peppermint, Pickleweed, Pigeon peas, Pineapple, Potatoes, Prickly Pear, Psoralea, Pumpkin, Quinoa, Radicchio, Radish, Rape, Rice, Rice bean, Rosemary, Rye, Safflower, Saffron, Sage, Saghuro, Sago, Scarlet runner bean, Snake gourd, Snap bean, Sorghum, Soybean, Spinach, Squash, Strawberry, Sugar cane, Summer savory, Summer squash, Sunflower, Sweet basil, Sweet Marjoram, Sweet potato, Tangerine, Taro, Tarragon, Tarwi, Tepary bean, Thyme, Tobacco, Tomato, Triticale, Vanilla, Vigna, Water hyacinth, Watercress, Watermelon, Wheat, Winged bean, Winter melons, Winter savory, Yam, Yam bean, Yard-long bean and Ye-eb.

CONTROLS: Mites (Broad, Brown, Citrus, Fist, Pacific, Russet, Rust, Silver, Sixspotted, Strawberry [Atlantic], Tomato), Powdery mildew, Rust.

APPLICATION RATES - DRY

Mites (Brown, Twospotted) 4 to 6 pounds per acre
 Mites (Pacific, Strawberry [Atlantic]) 3 to 6 pounds per acre
 Tomato russet mite 2 to 5 pounds per acre
 Mites (Citrus, Fist, Rust, Silver, Sixspotted, Yuma)..... 2 to 4 pounds per 100 gallons
 Scale crawlers 2 to 4 pounds per 100 gallons
 Powdery mildew 15 to 20 pounds per acre
 Rust 10 to 20 pounds per acre

APPLICATION FREQUENCY

Repeat sprays as necessary.

APPLICATION RATES - NO. OF WATER-SOLUBLE PAKS

Mites (Brown, Twospotted) 2 to 3 per 10 acres
 Mites (Pacific, Strawberry [Atlantic]) 3 to 6 per 20 acres
 Tomato russet mite 2 to 5 per 20 acres
 Mites (Citrus, Fist, Rust, Silver, Sixspotted, Yuma)..... 1 to 2 per 1,000 gallons
 Scale crawlers 1 to 2 per 1,000 gallons
 Powdery mildew 3 to 4 per 4 acres
 Rust 1 to 2 per 2 acres

APPLICATION FREQUENCY

Repeat sprays as necessary.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

PESTICIDE STORAGE: Storage should be under lock and key and secure from access by unauthorized persons and children. Storage should be in a cool, dry area away from any heat or ignition source. Do not stack over 2 pallets high. Move bags carefully so as not to tear or puncture. Do not move containers from one area to another unless they are securely sealed. Keep container tightly sealed when not in use. Do not allow bags to become wet or to be stored in a damp, humid area. Keep away from any puncture source. Avoid storage near water supplies, food, feed and fertilizer to avoid contamination. Store in original container only. If the contents are leaking or material is spilled, follow these steps:
 1. Collect and place in suitable containers for disposal.
 2. Wash area with soap and water to remove remaining pesticides.
 3. Follow washing with clean water rinse.
 4. Do not allow runoff to enter sewer or contaminate water supplies.
 5. Dispose of waste as indicated below.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

WATER-SOLUBLE PAKS: Carefully open outer bag. Never open the water-soluble pouch.

CONTAINER DISPOSAL: Completely empty bag into application equipment, then dispose of bag in a sanitary landfill or by incineration or, if allowed by State and Local authorities, by burning. If burned, stay out of smoke.

CONTAINER DISPOSAL: (For Residential/Household Use):

If empty: Do not reuse this container. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

WARRANTY— CONDITIONS OF SALE

OUR RECOMMENDATIONS FOR USE of this product are based upon tests believed reliable. Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the Seller. Buyer assumes all risks of use, storage and handling of this material in strict accordance with directions given herewith. In no case shall the Manufacturer or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product when such use and/or handling is not in strict accordance with directions given herewith. The foregoing is a condition of sale by the Seller and is accepted as such by the Buyer.

8/8



Drexel Chemical Company

November 6, 2002

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504C)
U.S. Environmental Protection Agency
Ariel Rios Bldg
1200 Pennsylvania Avenue, N.W.
Washington, D.C. 20460-0001

**Re: Notification Submission of a Revised Label per PR Notice 2001-6
DREXEL SULFUR 90W (EPA Reg. No. 19713-238)**

In support of the above, herewith, please find:

- 1. Completed EPA Form 8570-1
(238MSP-1002++)
- 2. One (1) copy of the label with the following revision:

The storage and disposal statements per PR Notice 2001-6 were incorporated in the Storage and Disposal box.

- 3. Certification statement per PR Notice 2001-6

If you have questions/clarification regarding this submission, I can be reached at (901) 774-4370. My e-mail address is lpiwonka@drexchem.com.

Thank you.

Respectfully,
DREXEL CHEMICAL COMPANY

Luz G Piwonka
Luz G Piwonka
Registration Manager

