

P1113 19713-263



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (H7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number:

19713-263

Date of Issuance:

APR 26 1995

Term of Issuance:

Conditional

NOTICE OF PESTICIDE:

Registration
 Reregistration

(under FIFRA, as amended)

Name of Pesticide Product:

DREXEL Diazinon 5G

Name and Address of Registrant (include ZIP Code):

Drexel Chemical Company
P.O. Box 13327
Memphis, TN 38113-0327

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
2. Make the labeling changes listed below before you release the product for shipment:
 - a. Replace the phrase "EPA Reg. No. 19713-EAG" with "EPA Reg. No. 19713-263".
 - b. Revise the first sentence of the "IF SWALLOWED" advice to read: Call a physician or Poison Control Center immediately.
 - c. Revise the first sentence if the IF ON SKIN advice to read: Wash promptly with plenty of soap and water.
 - d. In the sixth sentence of the HAZARDS TO HUMANS AND DOMESTIC ANIMALS, revise "tablespoons" to "teaspoons" or "spoons", since the small use rates on the label are in teaspoons.

Signature of Approving Official

Date

APR 26 1995

19917

- e. Revise the second sentence of the ENVIRONMENTAL HAZARDS to read: Birds, especially waterfowl (ducks or geese), feeding or drinking on treated area may be killed. Also, you may delete "and waterfowl" from the end of the fifth sentence.
- f. In the listing of pests controlled under the heading IN GARDENS DIAZINON 5G KILLS, delete Armyworms, Carrot Rust Flies, Corn Borers, and Rootworms since these pests are not included in specific directions for use or any crop.
- g. The heading and text (i.e, through item 7) of the Special Precautions to Reduce Risk to Birds should appear within an outline box.
- h. Under the heading VEGETABLES - INSECT CONTROL, delete "Unless directions under Timing/Special Directions state otherwise," and capitalize the "A" of "Applications", since it will be the first word in the sentence that remains.
- i. In the Note under the crop Beans, delete ", etc." and add "or" before "navy beans."
- j. In the crop listing for Melons, relocate the closing parenthesis mark to appear after "these" and then add comma after the closing parenthesis marks so that the end of the crop listing reads "...& Hybrids of these), and Watermelons. In the Rate column for these same crops, revise the entry for cutworms to read: 7-14 ozs. per 500 sq. ft. Also, is the omission of directions for control of wireworms intentional? If not, add this pest with the associated Rate of 10 1/2-14 ozs. per 500 sq. ft.
- k. For the crop Onions, in the Timing/Special Directions delete "by disking, harrowing, or other suitable means", since this is just a carryover from the agricultural use label
- l. For the crop Peas, add "(succulent only)" under the crop name.
- m. For the crop Radishes, delete the pest Root Maggots and the associated directions for use. In-furrow application of granular diazinon is not supported for reregistration
- n. For the crop Sweet Corn, revise the Timing/Special Directions for control of Seed Corn Maggots to read: Broadcast and mix into soil 2 inches just before planting.
- o. Be sure to meet type size requirements for elements of the label with minimum type size requirements.

3. Submit three (3) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.
4. Within eighteen months, submit the results of a one-year storage stability study for this product in the container to be marketed under normal storage conditions. Storage stability studies for diazinon products are still required because of the regulatory history for this active ingredient.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Enclosures

**DREXEL
DIAZINON 5G
LAWN AND GARDEN INSECT CONTROL**
**THIS PRODUCT MUST NOT BE USED ON GOLF COURSES AND SOD FARMS.
FOR OUTDOOR USE ON THE HOME LAWN AND IN THE HOME GARDEN ONLY.**
**READY-TO-USE GRANULES
TREATS UP TO 5000 SQUARE FEET**

DIAZINON KILLS:

- | | |
|--------------------|-----------------|
| Ants | Fire Ants* |
| Armyworms | Fleas |
| Bermudagrass Mites | Onion Maggots |
| Billbugs | Root Maggots |
| Brown Dog Ticks | Sod Webworms |
| Chiggers | White Grubs of: |
| Chinch Bugs | Japanese Beetle |
| Crickets | European Chafer |
| Cutworms | Southern Chafer |
| Earwigs | Wireworms |
| | and others |

* Aids in control.
See Directions for Use

ACTIVE INGREDIENT:

Diazinon: O,O-Diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate.....5.0%

INERT INGREDIENTS:.....95.0%

TOTAL:.....100.0%

Contains 5% Diazinon Insecticide

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

EPA Reg. No. 19713-EAG
EPA Est. No. 19713-GA-1

Manufactured by:
Drexel Chemical Company
P.O. Box 9306
Memphis, TN 38109

09/28/94-REVISED
04/13/95 Label Revision (Agency Ltr 4/12/95)
Net Weight: 10 Pounds

ACCEPTED
with COMMENTS
in EPA Letter Dated

APR 26 1995

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

19713-263

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 to 2 glasses of water and induce vomiting by touching the back of throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing.

IF ON SKIN: Wash with plenty of soap and water. Rinse thoroughly. Get medical attention.

IF IN EYES: Rinse eyes with plenty of water and call a physician immediately.

IF INHALED: Remove victim to fresh air and apply artificial respiration, if indicated.

NOTE TO PHYSICIAN: This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.

PRECAUTIONARY STATEMENTS**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating or smoking. Do not store near food or feed products. Food utensils such as tablespoons or measuring cups must not be used for food purposes after use in measuring pesticides. Do not contaminate ornamental fish ponds. Do not allow children or pets on treated areas until granules have been watered into the soil and the grass or soil is dry..

ENVIRONMENTAL HAZARDS

This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl feeding or drinking on treated areas maybe killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping granules. Granules that are spilled or are visible on soil surfaces should be covered with soil or thoroughly mixed into the soil. For lawn, follow all watering instructions in the directions for use, however, stop watering before puddling occurs. For lawns, do not apply this product to sites occupied by waterfowl, or within 75 ft. of any body of water which may attract waterfowl, such as ponds or lakes. Keep out of lakes, streams, ponds, tidal marshes, and estuaries. Do not apply directly to water. Runoff may be hazardous to fish and other aquatic organisms in neighboring areas.

Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters.

IN LAWNS DIAZINON 5G KILLS:	IN GARDENS DIAZINON 5G KILLS:
Ants Armyworms Bermudagrass Mites Billbugs Brown Dog Ticks Chiggers Chinch Bugs Clover Mites Crickets Cutworms Digger Wasps Earwigs Fleas Fire Ants* Lawn Moths Leafhoppers Millipedes Sod Webworms Sowbugs Springtails White Grubs of .European Chafer .Japanese Beetle .Southern Chafer	Armyworms Carrot Rust Flies Corn Borers Cutworms Mole Crickets Onion Maggots Root Maggots Rootworms Wireworms

*Aids in Control.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

OVERALL LAWN APPLICATION:

Special Precautions To Reduce Risk To Birds

Bird, especially waterfowl (ducks or geese), feeding and drinking on treated areas may be killed. To reduce risks to birds:

1. Do not treat areas larger than 15,000 sq. ft.
2. Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York between November 1 and May 20.

3. Do not exceed maximum permitted label rates, since rates above those recommended significantly increase potential hazards to birds, especially waterfowl.
4. Avoid overlapping treated areas.
5. Follow all watering instructions in the directions for use, however, stop watering before puddling occurs.
6. Do not apply this product to sites occupied by waterfowl or to sites within 75 ft. of any body of water that may attract waterfowl, such as lakes or ponds.
7. If it is necessary to repeat application, wait a minimum of 3 weeks between applications. Make no more than 4 applications per year.

Drexel Diazinon 5G controls the following lawn insects on both grass and dichondra lawns. Apply Diazinon 5G with fertilizer spreader. Refer to the tables that follow for use rates and spreader settings.

PESTS	RATE*	TIMING/SPECIAL DIRECTIONS
Ants, Armyworms, Bermudagrass Mites, Brown Dog Ticks, Chiggers, Chinch Bugs, Clover Mites, Crickets, Cutworms, Earwigs, Fleas, Lawn Moths (Sod Webworms), Leaf- hoppers, Millipedes, Sowbugs, Springtails (Collembola).	2 lbs. per 1,000 sq.ft.	Apply when grass is dry. Water lawn lightly after application. If necessary repeat application at 3 week intervals. Make no more than 4 applications per year.

PESTS	RATE*	TIMING/SPECIAL DIRECTIONS
Billbugs, White Grubs of Japanese Beetle, European Chafer, Southern Chafer, Ataenius Spretulus (Dung Beetle):	2 lbs. per 1,000 sq.ft.	Water grass with at least 1/4 inch of water after application, however, stop watering before puddling occurs. For billbug control, apply when activity is first observed or when chewed grass indicates damage. If necessary, repeat application at 3-week intervals. Make no more than 4 applications per year. For grub control, except dung beetle, treat any time between late July and early October. For dung beetle grub control, apply in Spring or Summer when damage is first noticed. Water grass with at least 1/4 inch of water after application, however, stop watering before puddling occurs. Treat again in late Summer or early Fall, if a second generation occurs.
Hyperodes Weevils	2 lbs. per 1,000 sq.ft.	Water grass with at least 1/4 inch of water after application, however, stop watering before puddling occurs. Treat problem areas in mid-April and again in mid-May.

*1,000 sq. ft. = an area 20 ft. x 50 ft.
 100 sq. ft. = an area 10 ft. x 10 ft.
 2 lbs. per 1,000 sq. ft. = 3.2 oz. per 100 sq. ft.

NOTE: The spreader settings listed below are approximate. Accurately weigh a few pounds into your spreader and apply to a measured area. Then adjust the spreader setting, if necessary, in order to assure correct rate of application.

LAWN SPREADER SETTING NO.**
(To deliver 2 lbs. per 1000 sq. ft.)

Spreader Make and Model	Setting
Central Imperial Mark No. 1662-G	6-1/2
Central Suburbia Lawn Spreader - No. 1662-0	6-1/2
Cyclone Model B1 Rotary	4-1/2
Greenfield Model 105	6-1/2
Ortho Drop Spreader Model 3000	4-1/2
Ortho Broadcast Spreader Model 4000	4-1/2
J.C. Penney Model 6005	7-1/4
J.C. Penney Rotary Spreader	5
Scott Precision Flow Spreader Model PF-4	4
Sears Craftsman Model 671.19198	6

**For application with other types of equipment, apply at the rate of 2 lbs. per 1000 sq. ft.

LAWN SPOT TREATMENT

Ants: Apply 1-1/2 teaspoons of Drexel Diazinon 5G over and around each ant hill.

Fire Ants: To aid in control of fire ants in lawns and other recreational areas, apply 1/2 cup over, and 2 ft. around, each fire ant mound. Immediately after application, water in thoroughly. Apply gently to avoid disturbing ants. Use of high pressure watering equipment may disturb the ants and cause migration, reducing product effectiveness. A minimum of 1 gallon of water should be applied to each mound area. Reapply as necessary.

NOTE: For best results, apply in cool weather, 65-80 degrees F., or in early morning or late evening hours. Works within 24 hours. No waiting for days or weeks. Treat new mounds as they appear.

Digger Wasps: Apply 1 1/2 teaspoons of Drexel Diazinon 5G in and around each nest opening. Apply at dusk when wasps are not active.

BAND TREATMENT AROUND HOUSE FOUNDATION:

The following pests are commonly found around or near foundations, and may enter the house:

- | | |
|-----------------|--------------------|
| Ants | Fleas |
| Brown Dog Ticks | Millipedes |
| Clover Mites | Sowbugs (Pillbugs) |
| Crickets | Springtails |
| Earwigs | (Collembola) |

Apply a 2 foot band of Drexel Diazinon 5G next to the house foundation. Distribute granules uniformly, using a rate of 3.2 oz. for each 100 sq. ft. of area (2' by 50') treated. If necessary, repeat application at 3-week intervals.

VEGETABLES - INSECT CONTROL


Special Precautions To Reduce Risk To Birds
 Birds feeding on treated areas may be killed. To reduce risk to birds:

- a) Thoroughly mix this product into the soil during or immediately after application. Refer to the Soil Incorporation table that follows.
- b) Granules that are spilled or are visible on soil surfaces should be covered with soil or thoroughly mixed into the soil.

Drexel Diazinon 5G controls certain garden insects on crops listed in the following tables. Apply uniformly as directed for each crop. When a range of rates is given, use the higher rate only for heavier infestations.

Do not apply this product to food crops grown in greenhouses.

Unless directions under Timing/Special Directions state otherwise, applications for control of pests listed below should be thoroughly mixed into the soil during or immediately after application.



SOIL INCORPORATION

Insect	Depth of Mixing or Incorporation into Soil
Surface Cutworms*	2-3 inches
Subterranean Cutworms*	3-6 inches
Mole Crickets	1-2 inches
Root Maggots	3-4 inches
Wireworms	4-8 inches

*If type of cutworm cannot be identified, mix into soil 4-6 inches deep.

CROP	PEST	RATE	TIMING/SPECIAL DIRECTIONS
Beans (Pole, Snap or Lima) (succulent only)	Cutworms	7-14 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting.
	Wireworms	10 1/2-14 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting.

NOTE: Do not use on dried bean varieties, such as pintos, dried limas, navy beans, etc.

Beets, Red (Table)	Cutworms	7-14 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting.
	Mole Crickets	3 1/2 ozs. per 500 sq.ft.	Broadcast and mix into soil 1-2 days before planting.
	Wireworms	10 1/2-14 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting.
Broccoli	Root Maggots	7-10 1/2 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting or transplanting.
	Cutworms	7-14 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting.
	Mole Crickets	3 1/2 ozs. per 500 sq.ft.	Broadcast and mix into soil 1-2 days before planting.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.

CROP	PEST	RATE	TIMING/SPECIAL DIRECTIONS
Brussels Sprouts	Root Maggots	7-10 1/2 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting or transplanting.
	Cutworms	7-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Mole Crickets	3 1/2 ozs. per 500 sq.ft.	Broadcast and mix into soil 1-2 days before planting.
	Wireworms	10 1/2-14 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting.
Cabbage	Root Maggots	7-10 1/2 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting or transplanting.
	Cutworms	7-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Mole Crickets	3 1/2 ozs. per 500 sq.ft.	Broadcast and mix into soil 1-2 days before planting.
	Wireworms	10 1/2-14 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting.
Carrots	Cutworms	7-14 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting.
	Mole Crickets	3 1/2 ozs. per 500 sq.ft.	Broadcast and mix into soil 1-2 days before planting.
	Wireworms	10 1/2-14 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting.

CROP	PEST	RATE	TIMING/SPECIAL DIRECTIONS
Cauliflower	Root Maggots	7-10 1/2 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting or transplanting.
	Cutworms	7-14 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting.
	Mole Crickets	3 1/2 oz. per 500 sq.ft.	Broadcast and mix into soil 1-2 days before planting.
	Wireworms	10 1/2-14 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting.
Celery	Cutworms	7-14 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting.
	Wireworms	10 1/2-14 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting.
Collards	Cutworms	7-14 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting.
	Mole Crickets	3 1/2 ozs. per 500 sq.ft.	Broadcast and mix into soil 1-2 days before planting.
	Wireworms	10 1/2-14 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting.
Cucumber	Cutworms	7-14 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting.
	Wireworms	10 1/2-14 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting.
Endive (Escarole)	Cutworms	7-14 ozs. per 500 sq.ft.	Broadcast and mix into soil just before

CROP	PEST	RATE	planting. TIMING/SPECIAL DIRECTIONS
	Mole Crickets	3 1/2 ozs. per 500 sq.ft.	Broadcast and mix into soil 1-2 days before planting.
	Wireworms	10 1/2-14 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting.
Kale	Cutworms	7-14 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting.
	Mole Crickets	3 1/2 ozs. per 500 sq.ft.	Broadcast and mix into soil 1-2 days before planting.
	Wireworms	10 1/2-14 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting.
Lettuce	Cutworms	7-14 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting.
	Mole Crickets	3 1/2 ozs. per 500 sq.ft.	Broadcast and mix into soil 1-2 days before planting.
	Wireworms	10 1/2-14 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting.
Melons (Cantaloupes, Casabas, Crenshaws, Honeydew, Muskmelons, Persians, & Hybrids of these and Watermelons.)	Cutworms	7-14 ozs. per	Broadcast and mix into soil just before planting.
Mustard	Mole Crickets	3 1/2 ozs. per 500 sq.ft.	Broadcast and mix into soil 1-2 days before planting.

CROP	PEST	RATE	TIMING/SPECIAL DIRECTIONS
Onions (Green and bulb)	Onion Maggots	7-14 ozs. per 500 sq.ft.	Broadcast just before planting and mix into the top 3-4 inches of soil by disking harrowing, or other suitable means. Note: Diazinon may not control organophosphate- resistant onion maggots.
	Wireworms	10 1/2-14 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting.
Parsley	Cutworms	7-14 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting.
	Wireworms	10 1/2-14 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting.
Peas	Cutworms	7-14 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting.
	Wireworms	10 1/2-14 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting.
NOTE: Do not use on dried pea varieties such as dried blackeyed peas.			
Peppers	Cutworms	7-14 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting.
	Mole Crickets	3 1/2 ozs. per 500 sq.ft.	Broadcast and mix into soil 1-2 days before planting.
	Wireworms	10 1/2-14 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting.

CROP	PEST	RATE	TIMING/SPECIAL DIRECTIONS
Potatoes	Cutworms (Surface & Subterranean)	7-14 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting.
	Wireworms	10 1/2-14 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting.
Radishes	Root Maggots	5 1/2-7 ozs. per 500 sq.ft.	Apply in the seed furrow at plant- ing (2-1/2 to 2-4/5 ozs. per 1,000 linear ft. of row).
	Mole Crickets	3 1/2 ozs. per	Broadcast and mix into soil 1-2 days before planting.
Spinach	Cutworms	7-14 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting.
	Wireworms	10 1/2-14 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting.
Squash, Summer & Winter	Cutworms	7-14 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting.
	Wireworms	10 1/2-14 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting.
Sweet Corn	Cutworms	7-14 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting.
	Seed Corn Maggots	7-14 ozs. per 500 sq.ft.	Broadcast just before planting. Mix into soil 2 inches.
	Wireworms	7-14 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting.
Sweet Potatoes	Wireworms, Flea Beetles	10 1/2 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting.

CROP	PEST	RATE	TIMING/SPECIAL DIRECTIONS
Swiss Chard	Cutworms	7-14 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting.
	Wireworms	10 1/2-14 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting.
Tomatoes	Cutworms	7-14 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting.
	Mole Crickets	3 1/2 ozs. per 500 sq.ft.	Broadcast and mix into soil 1-2 days before planting.
	Wireworms	10 1/2-14 ozs. per 500 sq.ft.	Broadcast and mix into soil just before planting.

STORAGE AND DISPOSAL

STORAGE: Store in original container in a cool, dry well-ventilated area. Store product in an area inaccessible to children and domestic animals. Do not store near fertilizers, herbicides, other insecticides, seeds, food or feed.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bag. Discard bag in trash.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes. In the event of a major spill, fire or other emergency, call 1-800-424-9300 day or night.

WARRANTY - CONDITIONS OF SALE

OUR RECOMMENDATIONS FOR USE of this product are based upon tests believed reliable. Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the Seller. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith.

In no case shall Drexel or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product when such use and/or handling is not in strict accordance with directions given herewith. The foregoing is a condition of sale By Drexel Chemical Company and is accepted as such by the Buyer.