

PM 25 19713-265

ACCEPTED
JUN 18 1995
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 19713-265

Page 178
1

**DREXEL
DEFOL 6W**

Defoliant/Desiccant with Fire Retardant

ACTIVE INGREDIENT:	
Sodium Chlorate	47.2%
INERT INGREDIENTS: 52.8%	
TOTAL:	100.0%

Product contains 6.0 pounds of Sodium Chlorate per gallon.

**KEEP OUT REACH OF CHILDREN
WARNING/AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

EPA REG. NO. 19713-265
EPA EST. NO. 19713-MS-1
NET CONTENTS:

MANUFACTURED BY:
DREXEL CHEMICAL COMPANY
P.O. BOX 9306
MEMPHIS, TN 38109-0306

WPS DELETIONS/REVISIONS 02/15/94
REVISED: 03/27/95 - Adding Potatoes

<p>STATEMENT OF PRACTICAL TREATMENT</p> <p>IF SWALLOWED: Drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid Alcohol. Get Medical Attention.</p> <p>IF IN EYES: Flush with plenty of water. Get Medical Attention.</p> <p>IF ON SKIN: Wash with plenty of soap and water. Get Medical Attention if irritation persists.</p>

198

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING. Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Harmful if swallowed.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

Coveralls over short-sleeved shirt and short pants

Waterproof gloves

Chemical-resistant footwear plus socks

Protective eyewear

Chemical-resistant headgear for overhead exposure

Chemical-resistant apron when cleaning equipment, mixing, or loading

Dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C)

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS


For terrestrial uses, do not apply to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.



For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls over short-sleeved shirt and short pants
- Waterproof gloves
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure

APPLICATION

DEFOL 6W may be applied by aircraft or ground equipment. It is essential that the foliage be thoroughly covered. The specified dilution rates should be carefully observed. A nonionic surfactant or an emulsifiable crop oil may enhance the results. DEFOL 6W should not be mixed with insecticides or other organic materials, unless specifically recommended, because a fire or explosion may result.

Precautions should be taken to avoid having spray or spray drift contact adjacent crops or desirable vegetation, or they may be defoliated. Equipment should be washed before and after use and care should be taken so that product does not dry in tank or on spray nozzles. Wash water should not be allowed to contact desirable vegetation.

DEFOLIATION

Ideally, defoliation is favored on clear, calm, sunny days with high temperatures and high humidity. If rain is anticipated within 24 hours, application should be delayed whenever possible. Defoliation may be slowed when daytime temperatures are below about 60 F., and with some crops, especially with vigorous varieties, regrowth can occur.

In order to obtain acceptable defoliation results, the crop must

be fully mature. In most cases, the signs of maturity include:

1. A slight discoloration of the foliage is apparent, often with a yellowish or reddish tinge.
2. Vegetative growth has slowed or ceased with little or no secondary growth.
3. A slight downward pull on a leaf gives a clean break from the stalk.

Adequate fertility, ample soil moisture, good insect control and good weed control also favor defoliation. Plants which have been under stress as from drought or insect pressure give less reliable results because tough, wilted or damaged leaves do not permit the proper uptake of the chemical and tend to inhibit the formation of an abscission layer. Typically, under such circumstances, the leaves are desiccated but do not drop, which, in turn, often causes problems with excessive trash during harvesting.

For crops where there is a heavy canopy of foliage, obtaining thorough coverage is sometimes difficult. Whenever practical, the use of ground equipment should be considered. The higher dilution rates should be used and any other available methods employed which will assure that the plants are completely covered all the way to the bottom leaves. If coverage is incomplete, a second application may be necessary in order to defoliate the lower leaves and to prevent secondary growth.

DESICCATION

When the objective is to reduce moisture of the crop or to desiccate weeds prior to harvest, most of the points discussed under Defoliation are applicable. Incomplete coverage will give erratic results, especially with vines such as morning glory where the slender stems may be sheltered from the spray by the crop canopy, leaving the vines succulent and tough so they will become entangled in the harvesting machinery. Once again, larger spray volume will encourage more complete coverage and more reliable desiccation.

CROPS

CHILI PEPPERS - Apply 1/2 to 1 1/4 gallons per acre in 5-15 gallons of water by air or 20-40 gallons of water by ground equipment, 10 days before harvest. To be used on peppers for processing only. Consult a processor before applying.

CORN - As a harvest aid to desiccate trashy weeds in early maturing corn, use 1 gallon DEFOL 6W in 5-7 gallons of water per acre by air at least 14 days before anticipated harvest date. Use the low (5 gallon) dilution rates when weeds are small and the crop canopy is fairly open. Desiccation of morning-glory and other vines may be erratic.

DO NOT graze treated fields or feed fodder, forage or residual grain within 14 days of application.

COTTON - Apply 1/2 to 3/4 gallon per acre in 5-10 gallons of water by air or 10-20 gallons of water by ground equipment no later than 7 days before harvest. The cotton plants should be mature with the youngest bolls expected to make cotton being at least 30 days old. Thorough coverage is essential.
 DO NOT graze treated fields or feed gin waste to livestock.

CUCURBITS (grown for seed): To facilitate harvest by drying out the vines, apply at a rate of 1 gallon per acre in 5-10 gallons of water by air, or apply 1 gallon per acre in 20-30 gallons of water by ground equipment.
 DO NOT graze treated fields within 14 days of application.


DRY BEANS, - Apply at a rate of 1 gallon per acre in 5-10 gallons of water by air or 10-20 gallons by ground equipment. Thorough coverage is essential. Make application 7-10 days before anticipated harvest, longer if temperatures are cool.
 DO NOT graze treated fields or feed treated fodder or forage to livestock.

FLAX - As a harvest aid to desiccate trashy weeds and to reduce moisture in the mature crop, apply 1 gallon per acre in enough water to give thorough coverage, 7-10 days before anticipated harvest date. For air application, use a minimum of 5 gallons of water and for ground equipment a minimum of 10 gallons of water.

Activity of the spray will be enhanced by the addition of good nonionic surfactant at the rate of 1 quart per 100 gallons of spray.
 DO NOT graze treated fields or feed treated flax foliage and fodder for 14 days after application.

GRAIN SORGHUM - To reduce the moisture content of the grain while the crop is standing apply 3/4 to 1 gallon per acre in sufficient water to give thorough coverage, 7-10 days before harvest. Apply a total of 5-10 gallons of spray per acre by air and 10-20 gallons per acre by ground equipment. Use the lower rates when grain moisture is low and the weather is clear and dry. Use the higher rate when conditions for desiccation are poor.

GUAR BEANS - Apply at a rate of 1 gallon per acre in 5-10 gallons of water by air or 10-20 gallons of water by ground equipment. Make application 7-10 days before anticipated harvest date.
 DO NOT graze treated fields or feed treated fodder or forage to livestock.



POTATOES - To defoliate plants and facilitate harvest, apply 1 gallon of DEFOL 6 in enough water to provide thorough coverage either by aircraft or ground equipment. Apply 10 days before harvest. Do not apply under conditions of extreme heat during the middle of the day.
AIR APPLICATION: Use 5 - 10 gallons of DEFOL 6W per acre.
GROUND APPLICATION: Use 10 - 20 gallons of DEFOL 6W per acre.

RICE - To facilitate harvest and reduce moisture in the fully mature seed heads, and to desiccate immature red rice in mature white rice, apply by air 3/4 - 1 gallon per acre in 5 or more gallons of water. Use the lower rate when crop conditions and weather are right to enhance rapid drying. Apply 7-10 days before anticipated harvest. To avoid drift do not apply during windy weather. Thorough coverage of the crop is essential to achieve good results.

DO NOT graze treated fields or feed treated straw or crop residues within 14 days of application.

SAFFLOWERS - To facilitate harvest and reduce moisture in the fully mature seed heads apply 1 gallon in 5-10 gallons of water by air or 20-30 gallons of water by ground equipment, 7 or more days before normal harvest when safflower heads are fully mature. DO NOT graze treated areas or feed treated forage.

SOUTHERN PEAS - Apply at a rate of 1 gallon per acre in 5-10 gallons of water by air or 10-20 gallons by ground equipment. Thorough coverage is essential. Make application 7-10 days before anticipated harvest, longer if temperatures are cool. DO NOT graze treated fields or feed treated fodder or forage to livestock.

SOYBEANS - As a soybean harvest aid to dry weeds in early maturing soybeans and to facilitate the harvest, apply 1 gallon per acre. Make application 7-10 days before anticipated harvest date when beans are mature and ready for harvest. For air application use a minimum of 5 gallons of water per acre; for ground equipment apply a minimum of 20 gallons of water per acre. DO NOT graze treated fields or feed treated soybean foliage.

SUNFLOWER - To facilitate harvest and reduce moisture in the fully mature seed heads apply 1 gallon in sufficient water to give good coverage, 7 or more days before normal harvest when sunflower heads are fully mature. Apply in 5-10 gallons of water per acre by air or in 20-30 gallons of water per acre by ground with a high clearance sprayer. Do not apply later than 7 days before harvest.
 DO NOT graze treated fields or feed treated forage within 14 days of application.

STORAGE AND DISPOSAL

6

Do not contaminate water, food, or feed by storage or disposal.

STORAGE INSTRUCTIONS: Storage should be under lock and key and secure from access by unauthorized persons and children. Storage should be in a cool dry area away from any heat or ignition source. Do not stack plastic containers over 2 pallets high. Move containers by handles or cases. Do not move containers from one area to another unless they are securely sealed. Keep container tightly sealed when not in use. Keep away from any puncture source. Avoid storage near water supplies, food, feed and fertilizer to avoid contamination. Store in original containers only. Unloading storage and work areas must be as clean as possible to prevent contamination if spilled. If concentrate or spray solution is allowed to dry on flammable objects, the chlorate component can act as an oxidizer. If the contents are leaking or material is spilled follow these steps:

1. Contain spill, absorb with a material such as saw dust, clay granules or dirt.
2. Collect and place in suitable containers for disposal.
3. Wash area with water and soap to remove remaining pesticide.
4. Follow washing with clean water rinse.
5. Clothing contaminated with chlorate solution must be removed at once before solution dries. Dried contaminated clothes can be ignited easily. Wash contaminated clothing immediately.
6. Do not allow run off to enter sewer or contaminate water supplies.
7. Dispose of waste as indicated below:

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Plastic containers - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary land-fill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY - CONDITION OF SALE

OUR RECOMMENDATIONS FOR USE of this product are based upon tests believed reliable. Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith.

In no case shall Drexel or the Seller be liable for consequential, special or indirect damages resulting from the use

or handling of this product when such use and/or handling is not in strict accordance with directions given herewith. The foregoing is a condition of sale by Drexel Chemical Company and is accepted as such by the Buyer.

