

19713-295

6/14/2010

1/13



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, DC 20460

OFFICE OF  
PREVENTION, PESTICIDES AND  
TOXIC SUBSTANCES

JUN 14 2010

Luz Chan  
Registration Manager  
Drexel Chemical Company  
1700 Channel Avenue  
Memphis, TN 38113-0327

Subject: Drexel Sulfa 80W  
EPA Reg. No. 19713-295  
Your amendment dated March 15, 2010  
EPA Decision Number 428348

Dear Ms Chan:

The amended label referred to above, submitted March 15, 2010 in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act as amended is acceptable.

One copy of the label stamped "Accepted" is enclosed for your records. This label supersedes all labels previously accepted for this product. Please submit one copy of the final printed label before the product is released for shipment.

If you have any questions, please contact Robert Westin by phone at (703) 305-5721 or via email at [westin.robert@epa.gov](mailto:westin.robert@epa.gov).

Sincerely,

Mary L. Waller  
Product Manager (21)  
Fungicide Branch  
Registration Division (7505P)

Enclosure

2/13

# MASTER LABEL



**ACCEPTED**  
**JUN 14 2010**  
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 19713-295

19713-295

# Sulfa 80W

Micronized Sulfur

**ACTIVE INGREDIENT:**

Sulfur ..... 80.0%

**OTHER INGREDIENTS:** ..... 20.0%

**TOTAL:** ..... 100.0%

**KEEP OUT OF REACH OF CHILDREN**

## CAUTION

See FIRST AID Below

EPA Reg. No. 19713-295  
EPA Est. No. 19713-XX-XXX

**Net Content: 25 Pounds**

FIRST AID	
<p><b>IF ON SKIN OR CLOTHING:</b></p> <ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>	
<p><b>IF IN EYES:</b></p> <ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>	
<p><b>IF SWALLOWED:</b></p> <ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious or convulsing person.</li> </ul>	
<p><b>IF INHALED:</b></p> <ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>	
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378.</p>	

Manufactured By:  
**Drexel Chemical Company**  
 P.O. BOX 13327, MEMPHIS, TN 38113-0327  
**SINCE 1972**

**PRECAUTIONARY STATEMENTS**  
**Hazards to Humans and Domestic Animals**

**CAUTION:** Avoid breathing spray mist. Harmful if inhaled or absorbed through skin. Avoid contact with skin, eyes or clothing.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category A on an EPA chemical resistance category selection chart.

**Applicators and other handlers must wear:** Long-sleeved shirt, long pants, chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride, shoes plus socks and protective eyewear. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**USER SAFETY RECOMMENDATIONS**

**Users should:** 1) Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. 2) Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. 3) Remove PPE immediately after handling this product. Wash the outside of the gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift. Do not contaminate water when disposing of rinsate or equipment washwater.

**PHYSICAL AND CHEMICAL HAZARDS**

Sulfur dust suspended in air ignites easily. Keep away from heat, sparks and flame. Do not smoke while applying this product.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval (REI). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is: Coveralls, chemical-resistant gloves made of any waterproof material, shoes plus socks and protective eyewear.

**MIXING DIRECTIONS**

Pour needed amount into spray tank under agitation. If necessary, add a few drops of oil to avoid foaming. Application of this material through any type of irrigation equipment is prohibited.

**USE PRECAUTIONS**

Some crops are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use on any crop unless local use has proven sulfur to be safe in that locality.

Sulfur may burn foliage or cause fruit injury when temperatures are excessively high. Do not apply to or allow spray drift on sulfur-sensitive plants such as Apricots, Cranberries, D'anjou pears or Spinach.

Do not apply with or following an oil application for 21 to 60 days. Do not apply oil for 14 days following treatment, except for dormant sprays on Pears. Follow all other label directions or precautions as indicated.

When growing crops for processing, consult processor before applying sulfur.

**DIRECTIONS FOR DILUTION**

**DILUTE APPLICATION**

**Ground Application:** Apply specified rate in 20 to 60 gallons of water per acre.

**Orchard Application:** Apply specified rate in 100 to 800 gallons of water per acre.

**CONCENTRATE APPLICATION**

**Ground Application:** Apply specified rate in not less than 5 gallons of water per acre.

**Orchard application:** Apply specified rate in 20 to 100 gallons of water per acre. Special concentrate equipment is necessary for these uses.

**Air Application:** Apply specified rate in 3 to 20 gallons of water per acre. Orchard rates should not be applied in less than 10 gallons of water per acre.

Use lower rates when using concentrate equipment. Use higher rates when using dilute applications.

**NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the WPS for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.  
Keep unprotected persons and domestic animals out of the treated area until sprays have dried.

Crop	Pest/Disease	Rate per Acre
Alfalfa	Atlantic mites, Red spider mites	3 to 5 lbs.
	USE INSTRUCTIONS: Apply as needed.	
Almonds, Cherries, Nectarines, Peaches, Plums, Plums, Prunes	Brown rot blossom blight, Leaf spot, Prune rot, Powdery mildew, Rust, Scab, Leaf spot, Silver mites, Flat mites, Red spider mites	10 to 30 lbs.
	USE INSTRUCTIONS: Apply at bloom or early petal fall. Repeat as necessary, usually 10 to 14 days after a period of wet weather. For brown rot of fruit, apply when fruit starts to ripen.	
Apples, Pears	Blister mites, Powdery mildew, Red spider mites, Scab	30 lbs.
	USE INSTRUCTIONS: Do not apply to D'anjou pears. Apply pre-bloom and petal fall. Apply with cover sprays throughout the season.	
Asparagus	Rust	10 to 30 lbs.
	USE INSTRUCTIONS: Use after cutting stops. Irrigate and cultivate before applying this product. Repeat at 7- to 10-day intervals or as needed through fern growth season.	
Avocados	Brown mite	10 to 30 lbs.
	USE INSTRUCTIONS: Apply as needed.	
Beans, Broccoli, Brussels sprouts, Cabbage, Cauliflower, celery, Collards, Kale, Lettuce, Mustard greens, Peas, Peppers, Potatoes, Rutabagas, Turnips	Powdery mildew, Red spider mites, Rust	3 to 10 lbs.
	USE INSTRUCTIONS: Start applications in early leaf stage. Repeat application at 10- to 14-day intervals or as needed.	
Blackberries, Blueberries, Boysenberries, Currants, Dewberries, Gooseberries, Huckleberries, Loganberries	Powdery mildew	6 to 15 lbs.
	USE INSTRUCTIONS: Begin applications when new shoots are 6 inches long and before blossoms open or when disease first appears. Repeat at 7- to 10-day intervals as necessary.	
Carrots, Garlic, Onions	Petrobia mites, Powdery mildew	3 to 10 lbs.
	USE INSTRUCTIONS: Apply when pest and disease first appears and repeat as needed.	
Cereal grains (Barley, Oats, Rice, Rye, Wheat), Corn, Milo, Sorghum, Soybeans, Sweet corn	For suppression of: Banks grass mites, Pacific mites, Red spider mites, Twospotted mites. For control of: Atlantic mites (Small grain only)	6 lbs.
	Powdery mildew (Soybeans only)	7.5 to 12.5 lbs.
	USE INSTRUCTIONS: Apply when disease first appears and repeat as necessary or with the regularly scheduled insect program.	
Citrus	Broad mites, Bud mites, Flat mites, Red spider mites, Rust mites, Silver mites, Sixspotted mites, Thrips, Yuma spider mites	8 to 20 lbs.
	USE INSTRUCTIONS: Apply in any spray NOT CONTAINING OIL. Do not apply within 21 days of an oil spray application.	
<b>(Continued)</b>		

(Cont.)		
Crop	Pest/Disease	Rate per Acre
Cotton	Atlantic mites, Red spider mites	3 to 5 lbs.
	USE INSTRUCTIONS: Apply as needed.	
Cucurbits	Powdery mildew	2 to 4 lbs.
	USE INSTRUCTIONS: Cantaloupe, and Cucumbers, Squash, and Watermelons are very sensitive to sulfur. Never apply more than 1.25 lbs. of this product per acre per application during periods of high temperatures.	
Grapes	Blister mites, Bud mites, Powdery mildew, Red spotted mites	6 lbs.
	USE INSTRUCTIONS: Apply in late dormant or with Cutworm spray, early shoot period and throughout season on a 14-day schedule or with pesticides.	
Macadamia nuts, Pecans, Walnuts	Almond mites, Broad mites, Pacific mites, Red spider mites, Twospotted mites	10 to 30 lbs.
	USE INSTRUCTIONS: Repeat as necessary. Apply with Mite sprays during season to increase the effectiveness. <b>Note:</b> Certain varieties of Walnuts are sensitive to sulfur sprays under certain conditions. Do not apply unless the varieties are known to be tolerant of sulfur.	
Mangoes	Powdery mildew	10 to 30 lbs.
	USE INSTRUCTIONS: Begin application before bloom. Repeat at bloom, after fruit set and 3 weeks later or as necessary.	
Melons, Squash	Powdery mildew	10 to 15 lbs.
	USE INSTRUCTIONS: Treat when the disease first appears and repeat as necessary. <b>Note:</b> Sulfur can injure plants, especially when the temperature reaches 95°F. Do not use on sulfur-sensitive varieties.	
Peanuts	Leaf spot, Powdery mildew, Red spider mites, Rust	3 to 5 lbs.
	USE INSTRUCTIONS: Start applications in early leaf stage. Repeat applications as necessary, usually 10 to 24 days.	
Peppermint, Spearmint	Powdery mildew	4 to 6 lbs.
	USE INSTRUCTIONS: Apply when Mint is 5 to 6 inches tall or when disease appears. Repeat twice at 30-day intervals. Do not apply within 30 days of harvest.	
Raspberries	Powdery mildew	6 to 15 lbs.
	USE INSTRUCTIONS: Spray each week starting at first bloom color. Continue until all fruit is set, then repeat as weather requires.	
Strawberries	Powdery mildew, Red spider mites	5 to 10 lbs.
	USE INSTRUCTIONS: Start applications in early leaf stage. Repeat applications as needed. For Powdery mildew, begin at first sign of disease and repeat as necessary. Do not use on sulfur-sensitive varieties.	
Sugar beets, Table beets	Powdery mildew, Red spider mites	5 to 10 lbs.
	USE INSTRUCTIONS: Apply at 10- to 30-day intervals with or without other pesticides as soon as the first symptoms appear.	
Tomatoes	Red spider mites, Russet mites	3 to 10 lbs.
	USE INSTRUCTIONS: Begin application as soon as first symptoms appear. Thorough coverage is required.	

Ornamentals	Pest/Disease	Rate per Acre
Asters, Carnations	Powdery mildew	3 to 10 lbs. by ground or 3 to 20 lbs. by air
Calendula, Cherry laurel, Chrysanthemums, Hollyhock, Similax	Leaf spot, Powdery mildew	
Clematis, Columbine, Ivy (English), Laurel, Petunia, Sage	Leaf spot	
Cosmos, Crepe myrtle, Dahlia, Delphinium, Gladiolus, Ligustrum, Sunflowers, Sweet peas, Violets, Zinnias	Powdery mildew	
Roses	Black spot, Powdery mildew	3 to 10 lbs. by ground 3 to 20 lbs. by air
USE INSTRUCTIONS: Apply in an aqueous spray ensuring complete plant coverage. Begin when disease first appears and repeat at 5- to 10-day intervals.		

**NUTRIENT USE**

Applications of this product will provide sulfur for the plants' nutrient requirements and should be considered in the total plant nutrient applications. This product can be used as a micronutrient supplement by either foliar or soil application. This product can be used on the following crops: Alfalfa, Almonds, Apples, Avocados, Barley, all Beans, Broccoli, Cabbage, Carrots, Cauliflower, Celery, Citrus, Corn, Grapes, Lawns, Lettuce, Melons, Milo, Nectarines, Peaches, Peanuts, Pears, Pecans, Peppers, Plums, Potatoes, Prunes, Rice, Sorghum, Soybeans, Sugar beets, Sugarcane, Strawberries, Sweet corn, Tomatoes, Turnips, Walnuts, Watermelons, Wheat and most other crops. Consult your State Agricultural Experimental Station or Extension Specialist for advice in selecting treatment to best fit local conditions.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage and disposal. Must be stored under dry conditions.

**PESTICIDE STORAGE:** Storage should be under lock and key and secure from access by unauthorized persons and children. Storage should be in a cool, dry area away from any heat or ignition source. Avoid storage at high temperatures. Do not stack over 2 pallets high. Move bags carefully so as not to tear or puncture. Do not move containers from one area to another unless they are securely sealed. Keep containers tightly sealed when not in use. Do not allow bags to become wet or store in a damp, humid area. Keep away from any puncture source. Avoid storage near water supplies, food, feed and fertilizer to avoid contamination. Store in original containers only. If the contents are leaking or material is spilled follow these steps:

1. Collect and place in suitable containers for disposal.
2. Wash area with soap and water to remove remaining pesticide.
3. Follow washing with clean water rinse.
4. Do not allow runoff to enter sewer or contaminate water supplies.
5. Dispose of waste as indicated below.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Do not contaminate water when disposing of equipment washwaters.

**CONTAINER DISPOSAL:**

**Nonrefillable Container (flexible-bag-all weights):** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**Nonrefillable Container (rigid-fifty lbs. or less):** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container one-fourth full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**Nonrefillable Container (rigid-greater than fifty lbs.):** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the

(Continued)

## STORAGE AND DISPOSAL (Cont.)

remaining contents into application equipment or a mix tank. Fill the container one-fourth full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.


**Refillable Container:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.


## WARRANTY—CONDITIONS OF SALE

OUR DIRECTIONS FOR USE of this product are based upon tests believed reliable. Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the Seller. To the extent consistent with applicable law, Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith.

To the extent consistent with applicable law, in no case shall the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product when such use and/or handling is not in strict accordance with directions given herewith. The foregoing is a condition of sale by the Seller and is accepted as such by the Buyer.

The Drexel logo is a registered trademark of Drexel Chemical Company.





# Sulfa 80W

Micronized Sulfur

Water-Soluble Packaging

**ACTIVE INGREDIENT:**

Sulfur ..... 80.0%

**OTHER INGREDIENTS:** ..... 20.0%

**TOTAL:** ..... 100.0%

**KEEP OUT OF REACH OF CHILDREN**

## CAUTION

See FIRST AID Below

EPA Reg. No. 19713-295

EPA Est. No. 19713-XX-XXX

Net Content: 25 Pounds

FIRST AID	
<b>IF ON SKIN OR CLOTHING:</b>	
<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>	
<b>IF IN EYES:</b>	
<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>	
<b>IF SWALLOWED:</b>	
<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious or convulsing person.</li> </ul>	
<b>IF INHALED:</b>	
<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>	
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378.</p>	



Manufactured By:  
**Drexel Chemical Company**

P.O. BOX 13327, MEMPHIS, TN 38113-0327

**SINCE 1972**



**PRECAUTIONARY STATEMENTS**  
**Hazards to Humans and Domestic Animals**

**CAUTION:** Avoid breathing spray mist. Harmful if inhaled or absorbed through skin. Avoid contact with skin, eyes or clothing.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category A on an EPA chemical resistance category selection chart.

**Applicators and other handlers must wear:** Long-sleeved shirt, long pants, chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride, shoes plus socks and protective eyewear. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**USER SAFETY RECOMMENDATIONS**

**Users should:** 1) Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. 2) Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. 3) Remove PPE immediately after handling this product. Wash the outside of the gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift. Do not contaminate water when disposing of rinsate or equipment washwater.

**PHYSICAL AND CHEMICAL HAZARDS**

Sulfur dust suspended in air ignites easily. Keep away from heat, sparks and flame. Do not smoke while applying this product.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval (REI). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval of 24 hours.

PPE required for early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is: Coveralls, chemical-resistant gloves made of any waterproof material, shoes plus socks and protective eyewear.

**NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the WPS for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Keep unprotected persons and domestic animals out of the treated area until sprays have dried.

**MIXING DIRECTIONS FOR USE OF WATER-SOLUBLE PACKAGING**

Paks containing this product are water-soluble. Do not allow pak to become wet prior to adding to the spray tank. Do not handle paks with wet hands. To prepare the spray mixture, drop the required number of unopened paks into spray tank while filling with the appropriate amount of water. Carefully open outer bag. Never open the water-soluble pouch. Operate the agitator while mixing. Depending on the water temperature and the degree of agitation, the paks should be completely dissolved in approximately 5 minutes from the time they are added to the water. Use the specified dosage of this product in the amount of water necessary to give complete coverage. Determine the total amount to be added to the spray tank based on the rates in the "Crop" section.

**Note:** Products containing BORON and/or fertilizers will prevent the water-soluble packet from dissolving. BORON and/or fertilizer products must only be added after packet(s) have completely dissolved.

**USE PRECAUTIONS**

Some crops are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use on any crop unless local use has proven sulfur to be safe in that locality.  
 Sulfur may burn foliage or cause fruit injury when temperatures are excessively high. Do not apply to or allow spray drift on sulfur-sensitive plants such as Apricots, Cranberries, D'anjou pears or Spinach.  
 Do not apply with or following an oil application for 21 to 60 days. Do not apply oil for 14 days following treatment, except for dormant sprays on Pears. Follow all other label directions or precautions as indicated.  
 When growing crops for processing, consult processor before applying sulfur.

**DIRECTIONS FOR DILUTION**

**DILUTE APPLICATION**

**Ground Application:** Apply specified rate in 20 to 60 gallons of water per acre.  
**Orchard Application:** Apply specified rate in 100 to 800 gallons of water per acre.

**CONCENTRATE APPLICATION**

**Ground Application:** Apply specified rate in not less than 5 gallons of water per acre.  
**Orchard application:** Apply specified rate in 20 to 100 gallons of water per acre. Special concentrate equipment is necessary for these uses.  
**Air Application:** Apply specified rate in 3 to 20 gallons of water per acre. Orchard rates should not be applied in less than 10 gallons of water per acre.

Use lower rates when using concentrate equipment. Use higher rates when using dilute applications.

Crop	Pest/Disease	Rate per Acre	Number of Water-Soluble Paks
Alfalfa	Atlantic mites, Red spider mites	3 to 5 lbs.	3 to 5 paks per 25 acres
	USE INSTRUCTIONS: Apply as needed.		
Almonds, Cherries, Nectarines, Peaches, Plums, Plums, Prunes	Brown rot blossom blight, Leaf spot, Prune rot, Powdery mildew, Rust, Scab, Leaf spot, Silver mites, Flat mites, Red spider mites	10 to 30 lbs.	2 to 6 paks per 5 acres
	USE INSTRUCTIONS: Apply at bloom or early petal fall. Repeat as necessary, usually 10 to 14 days after a period of wet weather. For brown rot of fruit, apply when fruit starts to ripen.		
Apples, Pears	Blister mites, Powdery mildew, Red spider mites, Scab	10 to 30 lbs.	2 to 6 paks per 5 acres
	USE INSTRUCTIONS: Do not apply to D'anjou pears. Apply pre-bloom and petal fall. Apply with cover sprays throughout the season.		
Asparagus	Rust	10 to 30 lbs.	2 to 6 paks per 5 acres
	USE INSTRUCTIONS: Use after cutting stops. Irrigate and cultivate before applying this product. Repeat at 7- to 10-day intervals or as needed through fern growth season.		
Avocados	Brown mite	10 to 30 lbs.	2 to 6 paks per 5 acres
	USE INSTRUCTIONS: Apply as needed.		
Beans, Broccoli, Brussels sprouts, Cabbage, Cauliflower, celery, Collards, Kale, Lettuce, Mustard greens, Peas, Peppers, Potatoes, Rutabagas, Turnips	Powdery mildew, Red spider mites, Rust	3 to 10 lbs.	3 to 10 paks per 25 acres
	USE INSTRUCTIONS: Start applications in early leaf stage. Repeat application at 10- to 14-day intervals or as needed.		
Blackberries, Blueberries, Boysenberries, Currants, Dewberries, Gooseberries, Huckleberries, Loganberries	Powdery mildew	6 to 15 lbs.	2 to 5 paks per 8 acres
	USE INSTRUCTIONS: Begin applications when new shoots are 6 inches long and before blossoms open or when disease first appears. Repeat at 7- to 10-day intervals as necessary.		

(Continued)

<b>(Cont.)</b>			
<b>Crop</b>	<b>Pest/Disease</b>	<b>Rate per Acre</b>	<b>Number of Water-Soluble Paks</b>
Carrots, Garlic, Onions	Petrobia mites, Powdery mildew	3 to 10 lbs.	3 to 10 paks per 25 acres
	USE INSTRUCTIONS: Apply when pest and disease first appears and repeat as needed.		
Cereal grains (Barley, Oats, Rice, Rye, Wheat), Corn, Milo, Sorghum, Soybeans, Sweet corn	For suppression of: Banks grass mites, Pacific mites, Red spider mites, Twospotted mites. For control of: Atlantic mites (Small grain only)	6 lbs.	3 paks per 12.5 acres
	Powdery mildew (Soybeans only)	7.5 to 12.5 lbs.	3 to 5 paks per 10 acres
	USE INSTRUCTIONS: Apply when disease first appears and repeat as necessary or with the regularly scheduled insect program.		
Citrus	Broad mites, Bud mites, Flat mites, Red spider mites, Rust mites, Silver mites, Sixspotted mites, Thrips, Yuma spider mites	8 to 20 lbs.	2 to 5 paks per 6 acres
	USE INSTRUCTIONS: Apply in any spray NOT CONTAINING OIL. Do not apply within 21 days of an oil spray application.		
Cotton	Atlantic mites, Red spider mites	3 to 5 lbs.	3 to 5 paks per 25 acres
	USE INSTRUCTIONS: Apply as needed.		
Cucurbits	Powdery mildew	2 to 4 lbs.	1 to 2 paks per 12.5 acres
	USE INSTRUCTIONS: Cantaloupe, and Cucumbers, Squash, and Watermelons are very sensitive to sulfur. Never apply more than 1.25 lbs. of this product per acre per application during periods of high temperatures.		
Grapes	Blister mites, Bud mites, Powdery mildew, Red spotted mites	6 lbs.	3 paks per 12.5 acres
	USE INSTRUCTIONS: Apply in late dormant or with Cutworm spray, early shoot period and throughout season on a 14-day schedule or with pesticides.		
Macadamia nuts, Pecans, Walnuts	Almond mites, Broad mites, Pacific mites, Red spider mites, Twospotted mites	10 to 30 lbs.	2 to 6 paks per 5 acres
	USE INSTRUCTIONS: Repeat as necessary. Apply with Mite sprays during season to increase the effectiveness. <b>Note:</b> Certain varieties of Walnuts are sensitive to sulfur sprays under certain conditions. Do not apply unless the varieties are know to be tolerant of sulfur.		
Mangoes	Powdery mildew	10 to 30 lbs.	2 to 6 paks per 5 acres
	USE INSTRUCTIONS: Begin application before bloom. Repeat at bloom, after fruit set and 3 weeks later or as necessary.		
Melons, Squash	Powdery mildew	10 to 15 lbs.	2 to 3 paks per 5 acres
	USE INSTRUCTIONS: Treat when the disease first appears and repeat as necessary. <b>Note:</b> Sulfur can injure plants, especially when the temperature reaches 95°F. Do not use on sulfur-sensitive varieties.		
Peanuts	Leaf spot, Powdery mildew, Red spider mites, Rust	3 to 5 lbs.	3 to 5 paks per 25 acres
	USE INSTRUCTIONS: Start applications in early leaf stage. Repeat applications as necessary, usually 10 to 24 days.		
Peppermint, Spearmint	Powdery mildew	4 to 6 lbs.	2 to 3 paks per 12.5 acres
	USE INSTRUCTIONS: Apply when Mint is 5 to 6 inches tall or when disease appears. Repeat twice at 30-day intervals. Do not apply within 30 days of harvest.		
<b>(Continued)</b>			

<b>(Cont.)</b>			
<b>Crop</b>	<b>Pest/Disease</b>	<b>Rate per Acre</b>	<b>Number of Water-Soluble Paks</b>
Raspberries	Powdery mildew	6 to 15 lbs.	2 to 5 paks per 8 acres
	USE INSTRUCTIONS: Spray each week starting at first bloom color. Continue until all fruit is set, then repeat as weather requires.		
Strawberries	Powdery mildew, Red spider mites	5 to 10 lbs.	1 to 4 paks per 5 acres
	USE INSTRUCTIONS: Start applications in early leaf stage. Repeat applications as needed. For Powdery mildew, begin at first sign of disease and repeat as necessary. Do not use on sulfur-sensitive varieties.		
Sugar beets, Table beets	Powdery mildew, Red spider mites	5 to 10 lbs.	1 to 4 paks per 5 acres
	USE INSTRUCTIONS: Apply at 10- to 30-day intervals with or without other pesticides as soon as the first symptoms appear.		
Tomatoes	Red spider mites, Russet mites	3 to 10 lbs.	3 to 10 paks per 25 acres
	USE INSTRUCTIONS: Begin application as soon as first symptoms appear. Thorough coverage is required.		

<b>Ornamentals</b>	<b>Pest/Disease</b>	<b>Rate per Acre</b>	<b>Number of Water-Soluble Paks</b>
Asters, Carnations	Powdery mildew	3 to 10 lbs. by ground or 3 to 20 lbs. by air	3 to 10 paks per 25 acres or 3 to 20 paks per 25 acres
Calendula, Cherry laurel, Chrysanthemums, Hollyhock, Similax	Leaf spot, Powdery mildew		
Clematis, Columbine, Ivy (English), Laurel, Petunia, Sage	Leaf spot		
Cosmos, Crepe myrtle, Dahlia, Delphinium, Gladiolus, Ligustrum, Sunflowers, Sweet peas, Violets, Zinnias	Powdery mildew		
Roses	Black spot, Powdery mildew	3 to 10 lbs. by ground	3 to 10 paks per 25 acres
		3 to 20 lbs. by air	3 to 20 paks per 25 acres
USE INSTRUCTIONS: Apply in an aqueous spray ensuring complete plant coverage. Begin when disease first appears and repeat at 5- to 10-day intervals.			

**NUTRIENT USE**

Applications of this product will provide sulfur for the plants' nutrient requirements and should be considered in the total plant nutrient applications. This product can be used as a micronutrient supplement by either foliar or soil application. This product can be used on the following crops: Alfalfa, Almonds, Apples, Avocados, Barley, Beans, Broccoli, Cabbage, Carrots, Cauliflower, Celery, Citrus, Corn, Grapes, Lawns, Lettuce, Melons, Milo, Nectarines, Peaches, Peanuts, Pears, Pecans, Peppers, Plums, Potatoes, Prunes, Rice, Sorghum, Soybeans, Sugar beets, Sugarcane, Strawberries, Sweet corn, Tomatoes, Turnips, Walnuts, Watermelons, Wheat and most other crops. Consult your State Agricultural Experimental Station or Extension Specialist for advice in selecting treatment to best fit local conditions.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal. Must be stored under dry conditions.

**PESTICIDE STORAGE:** Storage should be under lock and key and secure from access by unauthorized persons and children. Storage should be in a cool, dry area away from any heat or ignition source. Avoid storage at high temperatures. Do not stack over 2 pallets high. Move bags carefully so as not to tear or puncture. Do not move containers from one area to another unless they are securely sealed. Keep containers tightly sealed when not in use. Do not allow bags to become wet or store in a damp, humid area. Keep away from any puncture source. Avoid storage near water supplies, food, feed and fertilizer to avoid contamination. Store in original containers only. If the contents are leaking or material is spilled follow these steps:

1. Collect and place in suitable containers for disposal.
2. Wash area with soap and water to remove remaining pesticide.
3. Follow washing with clean water rinse.
4. Do not allow runoff to enter sewer or contaminate water supplies.
5. Dispose of waste as indicated below.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Do not contaminate water when disposing of equipment washwaters.

### CONTAINER DISPOSAL:

**Nonrefillable Container (flexible-bag-all weights):** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**Nonrefillable Container (rigid-fifty lbs. or less):** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container one-fourth full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**Nonrefillable Container (rigid-greater than fifty lbs.):** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container one-fourth full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**Refillable Container:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

## WARRANTY—CONDITIONS OF SALE

OUR DIRECTIONS FOR USE of this product are based upon tests believed reliable. Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the Seller. To the extent consistent with applicable law, Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith.

To the extent consistent with applicable law, in no case shall the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product when such use and/or handling is not in strict accordance with directions given herewith. The foregoing is a condition of sale by the Seller and is accepted as such by the Buyer.

The Drexel logo is a registered trademark of Drexel Chemical Company.