DATE: 3/30/93

### DREXEL PCNB 2-E LIQUID

For control of Brown Patch of certain lawns and certain soil borne diseases of ornamentals

## 

(Contains 2 lbs. of Pentachloronitrobenzene per U.S. Gallon @ 68° F.)

# KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid inhaling spray mist. This material may cause skin irritation. Wash hands and face after using and before eating or smoking. Avoid contact with skin by wearing the following protective clothing during mixing/loading operations; gloves, long sleeved shirt, long pants, socks and shoes (in addition, professional applicators must wear an approved respirator). Use with adequate ventilation. Do not contaminate feed or foodstuffs.

STATEMENT OF PRACTICAL TREATMENT


IF IN EYES: Flush with plenty of water. Get medical attention.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

IF SWALLOWED: Drink promptly a large quantity of milk, egg whites, gelatin-solution or, if these are not available, drink large quantities of water. Avoid alcohol. Get medical attention.

EPA REG. NO. 19713-312 EPA EST. NO.

DREXEL CHEMICAL COMPANY P.O. BOX 9306 MEMPHIS, TN 23109



BEST AVAILABLE COPT

#### ENVIRONMENTAL HAZARD

This pesticide is toxic to fish and aquatic organisms. Drift and runoff from treated areas may be hazardous to fish and aquatic organisms in adjacent aquatic sites. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water or wetlands when disposing of equipment washwaters. Do not apply directly to potable water supplies.

PHYSICAL OR CHEMICAL HAZARDS

Do not use, pour, spill or store near heat of open flame.

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not plant root crops in PCNB 2-E LIQUID treated fields within 12 months of broadcast or banding application unless PCNB 2-E LIQUID is registered for use on those crops. Do not apply this product through any type of irrigation system.

BROWN PATCH IN LAWNS AND ORNAMENTAL TURF
For control of Brown Patch (Rhizoctonia solani) of Southern
or warm season lawns such as St. Augustine, Bermuda and
similar grasses: Make one application in the Fall or Spring
at first signs of infection. Mix one quart in sufficient
water (10 to 15 gallons) to cover 1,000 square feet. Apply
with sprinkler can, hand sprayer, or any other convenient
method of obtaining even coverage. Then, lightly water the
treated area to insure movement of the chemical through the
grass to the soil level. Retreat the area 3 - 4 weeks later
if it has been subjected to unusually heavy rainfall, or is
flooded, or if the disease is severe or reappears. Under
certain growing conditions, a temporary discoloration of the
grass may appear occasionally. This causes no harm and
disappears shortly.

#### SNOW MOLD

•

For control of Gray Snow Mold (Typhula spp) and Pink Snow Mold (Fusarium Nivale). Make first application prior to first snowfall. Mix 1 1/2 pints in sufficient water (5 to 10 gallons) to cover 1,000 sq. ft. Treated area should be watered lightly as soon as possible following treatment to wash chemical down to the soil level.

Do not graze treated area or field grass clippings to livestock.

Under certain growing conditions, a temporary discoloration of the grass occasionally may occur. This causes no harm and will disappear in a short time.

BEST AVA'LABLE COPY

#### **DICHONDRA**

Treat as for grass, however, use 1 1/2 quarts per 40 gallons of water per 1,000 sq. ft.

## ORNAMENTALS (AZALEAS AND CAMELLIAS)

For Azelia Petal Blight and Camellia Flower Blight: Apply as a drench treatment to soil beneath the plants at a rate of 1 1/2 quarts in 30 gallons of water per 150 square feet. Begin application prior to opening of buds and repeat every 3 to 4 weeks through the bloom period.

#### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in safe place, away from source of heat or flame, and out of reach of children and domestic animals.

#### PESTICIDE DISPOSAL:

Pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to Federal or approved State procedures under Subtitle C of the Resource Conservation and Recovery Act.

## CONTAINER DISPOSAL:

Do not reuse empty container. Wrap container and put in trash.

### WARRANTY-CONDITION OF SALE:

DIRECTIONS FOR USE of this product are based on field use and tests believed reliable and should be followed carefully. It is however impossible to eliminate all risks associated with use of this product. Because such factors as weather conditions, foreign material and manner of use for application are all beyond the control of the Seller of this product, such things as crop injury, ineffectiveness or other unintended consequences may result. ALL SUCH RISKS ARE ASSUMED BY THE BUYER Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the directions for use. Seller makes no other warranties, expressed or implied, including FITNESS OR MERCHANTABILITY. In no case shall the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. The foregoing is a condition of sale by the Seller and is accepted as such by the Buyer.

