

**DREXEL
GLYPHOSATE 53.8%
HERBICIDE**

For broad-spectrum control of emerged weeds.

ACTIVE INGREDIENT:

*Isopropylamine salt of glyphosate. 53.8%


INERT INGREDIENTS: 46.2%

TOTAL. 100.0%

*Contains 648 grams per litre 5.4 pounds of active ingredient isopropylamine salt of N-(phosphonomethyl) glycine per U.S. gallon. Equivalent to 480 grams per litre or 4 pounds per U.S. gallon of the acid glyphosate.

CAUTION

KEEP OUT OF REACH OF CHILDREN



PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: May cause eye irritation. Avoid contact with eyes, skin or clothing.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush with plenty of water for at least 15 minutes. Call a physician.

IF ON SKIN: Flush with water. Wash clothing before reuse

ENVIRONMENTAL HAZARDS

Do not contaminate water by disposal of waste or cleaning of equipment. In case of spill or leak, soap up and remove to a landfill.

PHYSICAL OR CHEMICAL HAZARDS

Spray solution of this product should be mixed, stored and applied only in stainless steel, aluminum, fiberglass, plastic and plastic-lined steel containers. **DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTION IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS.** Glyphosate or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

GENERAL INFORMATION

AVOID CONTACT WITH FOLIAGE, GREEN STEMS, OR FRUIT OF CPOPS, DESIRABLE PLANTS AND TREES, SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT.

GLYPHOSATE 53.5% is a water soluble liquid, mixes readily with water and nonionic surfactant to be applied as a foliage spray for the control or destruction of most herbaceous plants. It may be applied through most standard industrial or field type sprayers after dilution and through mixing with water and surfactant in accordance with label instructions.

GLYPHOSATE 53.5% moves through the plant from the point of foliage contact to and into the root system. Visible effects on most annual weeds occur within 2 to 4 days but on most perennial weeds may not occur for 7 days or more. Extremely cool or cloudy weather following treatment may slow the activity of GLYPHOSATE 53.5% and delay visual effects of control. Visible effects are a gradual wilting and yellowing of the plant which advances to complete browning of above ground growth and deterioration of underground plant parts.

Unless otherwise directed on this label, delay application until vegetation has emerged and reached the stages described for control of such vegetation under the "Weeds Controlled" section of this label.

Unemerged plants arising from unattached underground rhizomes or root stocks of perennials will not be affected by the spray and will continue to grow. For this reason best control of most perennial weeds is obtained when treatment is made at late growth stages approaching maturity.

Always use the higher rate of GLYPHOSATE 53.5% per acre within the recommended range when weed growth is heavy or dense.

Do not treat weeds under poor growing conditions such as drought stress, disease or insect damage as reduced weed control may result. Reduced results may also occur when treating weeds heavily covered with dust.

Reduced control may result when applications are made to any weed or brush species that have been mowed, grazed, or cut, and have not been allowed to regrow to the recommended stage of treatment.

Rainfall or irrigation occurring within 6 hours after application may reduce effectiveness. Heavy rainfall or irrigation within 2 hours after application may wash the product off the foliage and a repeat treatment may be required.

GLYPHOSATE 53.5% does not provide residual weed control. For subsequent residual weed control, follow a label approved herbicide program. Read and carefully observe the Precautionary Statements and all other information appearing on the labels of all herbicides used.

Buyer and all users are responsible for all loss or damage in connection with the use or handling of mixtures of GLYPHOSATE 53.5% or other materials that are not expressly recommended in this label. Mixing GLYPHOSATE 53.5% with herbicides or other materials not recommended on this label may result in reduced performance.

ATTENTION: AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLY GLYPHOSATE 53.5% TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.

Do not allow the herbicide solution to mist, drip, drift, or splash onto desirable vegetation since minute quantities of GLYPHOSATE 53.5% can cause severe damage or destruction to the crop, plants, or other areas on which treatment was not intended. The likelihood of plant or crop injury occurring from the use of this product is greatest when winds are gusty or in excess of 5 miles per hour or when other conditions, including lesser wind velocities, will allow spray drift to occur. When spraying avoid combinations of pressure and nozzle type that will result in splatter or fine particles (mist) which are likely to drift. **AVOID APPLYING AT EXCESSIVE SPEED OR PRESSURE.**

NOTE: Use of GLYPHOSATE 53.5% in a manner not consistent with this label may result in injury to persons, animals or crops, or other unintended consequences. When not in use, keep container closed to prevent spills and contamination.

Clean sprayer and parts immediately after using this product by thoroughly flushing with water.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

AQUATIC SITES

When applied as directed under conditions described, GLYPHOSHATE 53.5% plus nonionic surfactant will control emerged annual weeds and control or partially control emerged perennial weeds, woody brush and trees listed in this label. GLYPHOSHATE 53.5% does not control plants which are either completely submerged or have a majority of the foliage under water. See the "Weeds Controlled" section of this label for rates and degree of control provided.

GLYPHOSHATE 53.5% may be used in and around aquatic sites, including all bodies of fresh and brackish water, which may be flowing, non-flowing, or transient. This includes lakes, rivers, streams, ponds, seeps, irrigation and drainage ditches, canals, reservoirs, and similar sites. There is no restriction on the use of water for irrigation, recreation, or domestic purposes.

Consult local state fish and game agency and water control authorities before applying GLYPHOSHATE 53.5% to public water. Permits may be required to treat such water.

NOTE: Do not apply GLYPHOSHATE 53.5% within 1/2 mile upstream of potable water intakes.

Do not apply this product on rice levees when flood water is present.

Do not apply in estuaries.

For treatments after drawdown of water or in dry ditches, allow 7 or more days after treatment before reintroduction of water. Apply GLYPHOSHATE 53.5% within one day after drawdown to ensure application to actively growing weeds.

When using GLYPHOSHATE 53.5% in aquatic sites where water is present, add 1 to 2 quarts of an agriculturally approved surfactant per 100 gallons of spray solution (1/4 to 1/2 percent surfactant by total spray volume). The following surfactants are approved for use with GLYPHOSHATE 53.5% in aquatic sites: Agri-Dex M, Induce M, Liqua-wet M, Ortho M X-77, Passage M, R-11 M, Spreader Sticker M, Super Spread M 200, and Widespread M.

When using GLYPHOSHATE 53.5% in sites where water is not present (dry ditches, ditchbanks, dry canals), use 1 to 2 quarts of nonionic surfactant per 100 gallons of spray solution (1/4 to 1/2 percent surfactant by total spray volume).

Floating mats of vegetation may require retreatment. Avoid washoff of sprayed foliage by spray boat or recreation boat backwash or by rainfall within six hours of application. Do not retreat within 24 hours following the initial treatment.

Applications made to moving bodies of water must be made while traveling upstream to prevent concentration of this herbicide in water. When making any bankside applications, do not overlap more than 1 foot into open water. Do not spray across open moving bodies of water, or where weeds do not exist. The maximum application rate of 7 1/2 pints per acre must not be exceeded in any single application.

When emerged infestations require treatment of the total surface area of impounded water, treating the area in strips may avoid oxygen depletion due to decaying vegetation. Oxygen depletion may result in fish kill.

TM Agri-Dex and Induce are trademarks of the Helena Chemical Company

TM Liqua-wet is a trademark of the Woodbury Chemical Company

TM Ortho X-77 is a trademark of Chevron Chemical Company

TM Passage is a trademark of Asgrow Florida Company

TM R-11 and Super Spread 200 are trademarks of the Wilbur Ellis Company

TM Spreader Sticker is a trademark of the Southern Mill Creek Products Company

TM Widespread is a trademark of the FMC Corporation

MIXING AND APPLICATION INSTRUCTIONS

APPLY THESE SPRAY SOLUTIONS IN PROPERLY MAINTAINED AND CALIBRATED EQUIPMENT CAPABLE OF DELIVERING DESIRED VOLUMES. HAND GUN APPLICATIONS SHOULD BE PROPERLY DIRECTED TO AVOID SPRAYING DESIRABLE PLANTS. NOTE: REDUCED RESULTS MAY OCCUR IF WATER CONTAINING SOIL IS USED, SUCH AS WATER FROM PONDS AND UNLINED DITCHES.

MIXING

GLYPHOSATE 53.5% mixes readily with water. Mix spray solutions of GLYPHOSATE 53.5% as follows: Fill the mixing or spray tank with the required amount of water while adding the required amount of GLYPHOSATE 53.5% (see "Directions for Use" and "Weeds Controlled" sections of this label).

Near the end of the filling process, add the required surfactant and mix well. Remove hose from tank immediately after filling to avoid siphoning back into the water source. During mixing and application, foaming of the spray solution may occur.

To prevent or minimize foam, avoid the use of mechanical agitators; place the filling hose below the surface of the spray solution, terminate by-pass and return lines at the bottom of the tank and if needed use an approved antifoam or defoaming agent.

Keep by-pass line on or near bottom of tank to minimize foaming. Screen size in nozzle or line strainers should be no finer than 50 mesh. Carefully select correct nozzle to avoid spraying a fine mist. For best results with conventional ground application equipment, use flat fan nozzles. Check for even distribution of spray droplets.

THE USE OF A NONIONIC SURFACTANT, APPROVED FOR THE SITE OF THE DESIRED APPLICATION IS REQUIRED FOR USE WITH GLYPHOSATE 53.5%. Use 1/4 to 1/2 percent surfactant by total spray volume. Carefully observe all Precautionary Statements and other information appearing on the surfactant label.

APPLICATION EQUIPMENT AND TECHNIQUES

AERIAL EQUIPMENT: DO NOT APPLY THIS PRODUCT BY AIR IN CALIFORNIA

Use the recommended rates of GLYPHOSATE 53.5% and surfactant in 3 to 20 gallons of water per acre as a broadcast spray. See the "Weeds Controlled" section of this label for specific rates. Aerial applications of GLYPHOSATE 53.5% may only be made as specifically recommended on this label.

AVOID DRIFT - DO NOT APPLY DURING INVERSION CONDITIONS, WHEN WINDS ARE GUSTY, OR UNDER ANY OTHER CONDITION WHICH WILL ALLOW DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Coarse sprays are less likely to drift, therefore, do not use nozzles or nozzle configurations which disperse spray as fine spray droplets. Do not angle nozzles forward into the airstream and do not increase spray volume by increasing nozzle pressure.

Drift control additives may be used. When a drift control additive is used, read and carefully observe the Precautionary Statements and all other information appearing on the additive label.

79167

Ensure uniform application - To avoid streaked, uneven or overlapped application, use appropriate marking devices.

Thoroughly wash aircraft, especially landing gears, after each day of spraying to remove residues of GLYPHOSATE 53.5% accumulated during spraying or from spills. PROLONGED EXPOSURE OF GLYPHOSATE 53.5% TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. LANDING GEAR ARE MOST SUSCEPTIBLE. The maintenance of an organic coating (paint) which meets aerospace specification MIL-C-38413 may prevent corrosion.

BOOM EQUIPMENT: For control of weed or brush species listed on this label using conventional boom equipment - Use the recommended rates of GLYPHOSATE 53.5% and surfactant in 3 to 20 gallons of water per acre as a broadcast spray. See the "Weed Controlled" section of this label for specific rates. As density of weeds increases, spray volume should be increased within the recommended range to insure complete coverage. Carefully select correct nozzle to avoid spraying a fine mist. For best results with ground application equipment, use flat fan nozzles. Check for even distribution of spray droplets.

HAND-HELD and HIGH-VOLUME EQUIPMENT: Use coarse spray only. For control of weeds listed on this label using knapsack sprayers or high volume spraying equipment utilizing handguns or other suitable nozzle arrangements - Prepare a 1/4 to 1 1/2 percent solution of GLYPHOSATE 53.5% in water, add surfactant and apply to foliage of vegetation to be controlled. For specific rates of application and instructions for control of various annual and perennial weeds, see the "Weed Controlled" section of this label.

Applications should be made on a spray-to-wet basis. Spray coverage should be uniform and complete. Do not spray to point of runoff.

Prepare the desired volume of spray solution by mixing the amount of GLYPHOSATE 53.5% in water, shown in the following table:

Spray solution

DESIRED VOLUME	AMOUNT OF GLYPHOSATE 53.5%			
	1/4%	1%	1 1/4%	1 1/2%
1 gallon	1 oz	1 1/3 oz	1 2/3 oz	2 oz
25 gallons	1 1/2 pt	1 qt	1 1/4 qt	1 1/2 qt
100 gallons	3 qt	1 gal	1 1/4 gal	1 1/2 gal

2 tablespoons = 1 ounce

For use in knapsack sprayers, it is suggested that the recommended amount of GLYPHOSATE 53.5% be mixed with water in a larger container. Fill sprayer with the mixed solution and add the correct amount of surfactant.

WEEDS CONTROLLED

ANNUAL WEEDS: Apply to actively growing annual grasses and broadleaf weeds.

Allow at least 3 days after application before disturbing treated vegetation. After this period the weeds may be mowed, tilled or burned. See "Directions For Use", "General Information", and "Mixing and Application Instructions" for labeled uses and specific application instructions.

Broadcast Application - Use 1 1/2 pints of GLYPHOSATE 53.5% per acre plus 1/4 to 1/2 percent surfactant by total spray volume if weeds are less than 6 inches tall. If weeds are greater than 6 inches tall, use 2 1/2 pints of GLYPHOSATE 53.5% per acre plus 1/4 to 1/2 percent surfactant by total spray volume.

Hand-Held High-Volume Application - Use a 3/4 percent solution of GLYPHOSATE 53.5% in water plus 1/4 to 1/2 percent surfactant by total spray volume and apply to foliage of vegetation to be controlled.

When applied as directed under the conditions described in this label, GLYPHOSATE 53.5% plus nonionic surfactant will control the following annual weeds:

- | | |
|------------------------|-------------------------|
| Barley | Panicum |
| Hordeum vulgare | Panicum spp. |
| Bluegrass (annual) | Pennycress (field) |
| Poa Annua | Thlaspi arvense |
| Brome (downy) | Pigweed, Redroot |
| Bromus tectorum | Amaranthus retroflexus |
| Cocklebur | Pigweed (smooth) |
| Xanthium pensylvanicum | Amaranthus hybridus |
| Corn (volunteer) | Ragweed (common) |
| Zea mays | Ambrosia artemisiifolia |
| Crabgrass | Ragweed (giant) |
| Digitaria spp. | Ambrosia trifida |
| Falseflax (smallseed) | Rye |
| Camelina microcarpa | Secale cereale |
| Fiddleneck | Ryegrass (Italian)* |

Amsinckia spp.	Lolium multiflorum
Fleabane	Sandbur (field)
Erigeron spp.	Cenchrus spp.

Foxtail	Shatercane
Setaria spp.	Sorghum bicolor
Kocia	Smartweed (Pennsylvania)
Kochia scoparia	Polygonium pensylvanicum
Lambsquarters (common)	Spanishneedles*
Chenopodium album	Bidens bipinnata
Lettuce (prickly)	Sunflower
Lactuca serriola	Helianthus annuus
Mustard (tansy)	Thistle (Russian)
Descurainia pinnata	Salsola kali
Oats (wild)	Velvetleaf
Avena fatua	Abutilon theophrasti

*Apply 3 pints of GLYPHOSATE 53.5% per acre.

Annual weeds will generally continue to germinate from seed throughout the growing season. Repeat treatment will be necessary to control later germinating weeds.

PERENNIAL WEEDS

Apply GLYPHOSATE 53.5% as follows to control or destroy most vigorously growing perennial weeds. Unless otherwise directed, allow at least 7 days after application before disturbing vegetation.

Add 1/4 to 1/2 percent nonionic surfactant by total spray volume to the rates of GLYPHOSATE 53.5% given in this list. See the "General Information", "Directions for Use", "Mixing and Application" sections of this label for specific uses and application instructions.

NOTE: If weeds have been mowed or tilled, do not treat until regrowth has reached the recommended stages. Fall treatments must be applied before a killing frost.

Repeat treatments may be necessary to control weeds regenerating from underground parts or seed.

When applied as recommended under the conditions described, GLYPHOSATE 53.5% plus surfactant will control the following Perennial Weeds:

Alfalfa	Dogbane (hemp)
Medicago sativa	Apocynum cannabinum
Alligator weed*	Fescues
Alternanthera philoxeroids	Festuca spp.
American Lotus	Guineagrass

Nelumbo lutea
 (FOR USE IN
 TEXAS ONLY)
 Artichoke (Jerusalem)
 Helianthus tuberosus

Panicum maximum
 Horsenettle
 Solanum carolinense
 Horseradish
 Armoracia rusticana

Bahiagrass
 Paspalum notatum
Bermudagrass
 Cynodon dactylon
 Bindweed (field)
 Convolvulus arvensis
 Bluegrass (Kentucky)
 Poa spp.
 Brackenfern
 Pteridium aquilinum
 Bromegrass (smooth)
 Bromus inermis
 Cattail
 Typha spp.
 Clover (red)
 Trifolium pratense
 Clover (white)
 Trifolium repens
 Cutgrass (giant)*
 Zizaniopsis miliacea
 Dallisgrass
 Paspalum dilatatum
 Dandelion
 Taraxacum officinale
 Dock (curly)
 Rumex crispus
 Paragrass
 Brachiaria mutica
 Phragmites**
 Phragmites spp.
 Quackgrass
 Agropyron repens
 Reed canarygrass
 Phalaris arundinacea
 Ryegrass (perennial)
 Lolium perenne
 Smartweed (swamp)
 Polygonum coccineum
 Spattledock
 Nuphar luteum

Johnsongrass
 Sorghum halepense
Kikuyugrass
 Pennisetum clandestinum
 Knapweed
 Centaurea repens
 Lantana
 Lantana camara
 Maidencane
 Panicum hematomon
 Milkweed
 Asclepias spp.
 Muhly (wirestem)
 Muhlenbergia frondosa
 Mullein (common)
 Verbascum thapsus
 Napiergrass
 Pennisetum purpureum
 Nightshade (silverleaf)
 Solanum elaeagnifolium
 Nutsedge (purple, yellow)
 Cyperus rotundus
 Cyperus esculentus

Orchardgrass
 Dactylis glomerata
 Texas Blueweed
 Helianthus ciliaris
 Thistle (Canada)
 Cirsium arvense
 Timothy
 Phleum pratense
 Torpedograss*
 Panicum repens
 Vaseygrass
 Paspalum urvillei
 Water Hyacinth
 Eichornia crassipes
 Wheatgrass (western)
 Agropyron smithii

*Partial control
 **Partial control in Southeastern states. See specific
 recommendation below.

Alligatorweed: Apply 6 pints of GLYPHOSHATE 53.5% per acre
 as a broadcast spray or as the 1 1/4 percent solution with

hand-held equipment to provide partial control of alligatorweed.

Apply when most of the target plants are in bloom. Repeat applications will be required to maintain such control.

American Lotus: (FOR USE IN TEXAS ONLY) Apply 4 pints of GLYPHOSATE 53.5% per acre as a broadcast spray or as a 1 percent solution with hand-held equipment. Treat when plants are actively growing at or beyond the bloom stage of growth. Best results are achieved when application is made during summer or fall months. Fall treatments must be applied before a killing frost. Repeat treatment may be necessary to control regrowth from underground parts and seeds.

Bermudagrass: Apply 7 1/2 pints of GLYPHOSATE 53.5% per acre as a broadcast spray or as a 1 1/2 percent solution with hand-held equipment. Apply when target plants are actively growing and when seed heads appear.

Brackenfern: Apply 4 1/2 to 6 pints of GLYPHOSATE 53.5% per acre as a broadcast spray or as a 3/4 to 1 percent solution with hand-held equipment. Apply to fully expanded fronds which are at least 18 inches long.

Canada Thistle: Apply 3 to 4 1/2 pints of GLYPHOSATE 53.5% per acre as a broadcast spray or as a 1 1/2 percent solution with hand-held equipment. Apply when target plants are actively growing and are at or beyond the bud stage of growth. Allow 3 or more days after application before tillage.

Cattail: Apply 4 1/2 to 6 pints of GLYPHOSATE 53.5% per acre as a broadcast spray or as a 3/4 percent solution with hand-held equipment. Apply when target plants are actively growing and are at or beyond the early-to-full bloom stage of growth. Best results are achieved when application is made during the summer or fall months.

Cutgrass (giant): Apply 6 pints of GLYPHOSATE 53.5% per acre as a broadcast spray or as a 1 percent solution with hand-held equipment to provide partial control of giant cutgrass. Repeat applications will be required to maintain such control, especially where vegetation is partially submerged in water. Allow for substantial regrowth to the seven-to-ten leaf stage prior to retreatment.

Field Bindweed/Silverleaf Nightshade/Texas Bluweed: Apply 6 to 7 1/2 pints of GLYPHOSATE 53.5% per acre as a broadcast spray west of the Mississippi River and 4 1/2 to 6 pints of GLYPHOSATE 53.5% per acre east of the Mississippi River. With hand-held equipment, use a 1 1/2 percent solution.

Nutsedge (purple, yellow): Apply 4 1/2 pints of GLYPHOSATE 53.5% per acre as a broadcast spray, or as a 3/4 percent solution with hand-held equipment to control existing nutsedge plants and immature nutlets attached to treated plants.

Apply when target plants are in flower or when new nutlets can be found at rhizome tips. Nutlets which have not germinated will not be controlled and may germinate following treatment. Repeat treatments will be required for long-term control.

Phragmites: In SC, GA, AL, FL, MS, LA, or TX, apply 7 1/2 pints of GLYPHOSATE 53.5% per acre as a broadcast spray or as a 1 1/2 percent solution with hand-held equipment to provide partial control of Phragmites. In all other states apply 6 pints of GLYPHOSATE 53.5% per acre as a broadcast spray or as a 3/4 percent solution with hand-held equipment to provide control. Apply when most of the target plants are in full bloom, or during the fall months. Repeat treatments will be required to maintain control.

Quackgrass/Wirestem Muhly/Kikuyugrass: Apply 3 to 4 1/2 pints of GLYPHOSATE 53.5% per acre as a broadcast spray or as a 3/4 percent solution with hand-held equipment when most quackgrass or wirestem muhly is at least 8 inches in height (3 to 4-leaf stage of growth) and actively growing. Allow 3 or more days after application before tillage.

Spatterdock: Apply 6 pints of GLYPHOSATE 53.5% per acre as a broadcast spray or as a 3/4 percent solution with hand-held equipment. Apply when most plants are in full bloom. For best results, apply during the summer or fall months.

Torpedograss: Apply 6 to 7 1/2 pints of GLYPHOSATE 53.5% per acre as a broadcast spray or as a 3/4 to 1 1/2 percent solution with hand-held equipment to provide partial control of torpedograss. Use the lower rates under terrestrial conditions, and the higher rates under partially submerged or a floating mat condition. Repeat treatments will be required to maintain such control.

Water Hyacinth: Apply 5 to 6 pints of GLYPHOSATE 53.5% per acre as a broadcast spray or as a 1 percent solution with hand-held equipment. Apply when target plants are actively growing and at or beyond the early bloom stage of growth. After application, visual symptoms may require 3 or more weeks to appear with complete necrosis and decomposition usually occurring within 60 to 90 days.

Other perennials listed on this label: Apply 4 1/2 to 7 1/2 pints of GLYPHOSATE 53.5% per acre as a broadcast spray or

as a 3/4 to 1 1/2 percent solution with hand-held equipment. Apply when target plants are actively growing and most have reached early head or early bud stage of growth.

WOODY BRUSH AND TREES

Apply GLYPHOSATE 53.5% as follows to control actively growth woody brush and trees listed below. Unless otherwise specified, allow at least 7 days after application before disturbing treated vegetation.

Add 1/4 to 1/2 percent surfactant by total spray volume to the rates of GLYPHOSATE 53.5% given in this list. See the "General Information", "Directions for Use", and "Mixing and Application" sections of this label for specific uses and applications instructions

When applied as recommended under the conditions described, GLYPHOSATE 53.5% plus surfactant controls the following woody brush plants and trees:

- | | |
|--------------------|------------------|
| Alder | Kudzu |
| Alnus spp. | Pueraria lobata |
| Berries* | Maple** |
| Rubus spp. | Acer spp. |
| Elderberry | Oak*** |
| Sambucus spp. | Quercus spp. |
| Honeysuckle | Multiflora rose |
| Lonicera spp. | Rosa multiflora |
| Poison Ivy | Trumpet creeper |
| Rhus radicans | Campsis radicans |
| Poison Oak | Willow |
| Rhus toxicodendron | Salix spp. |

*Includes blackberry, dewberry and raspberry.

**Includes sugar maple and red maple.

***Includes red oak, white oak and Northern pin oak.

Apply GLYPHOSATE 53.5% when plants are vigorously growing, and unless otherwise directed, after full leaf expansion. Use the higher rate for larger plants and/or dense areas of growth. On vines, use the higher rate for plants that have reached the woody stage of growth. Best results are obtained when application is made in late summer or fall after fruit formation. Ensure thorough coverage when using hand-held equipment. Symptoms may not appear prior to frost or senescence with fall treatments.

Repeat treatments may be necessary to control plants regenerating from underground parts or seed. Some autumn colors on undesirable deciduous species are acceptable provided no major leaf fall has occurred.

Apply GLYPHOSATE 53.5% as follows to control the following listed plants and trees.

Alder/Berries/Elderberry/Honeysuckle: Apply 4 1/2 to 6 pints of GLYPHOSATE 53.5% per acre as a broadcast spray or as a 3/4 to 1 percent solution with hand-held equipment.

Kudzu: Apply 6 pints of GLYPHOSATE 53.5% per acre as a broadcast spray or as a 1 1/2 percent solution with hand-held equipment.

Maples/Oaks: Apply GLYPHOSATE 53.5% as a 3/4 to 1 percent solution with hand-held equipment. Apply when at least 50 percent of the new leaves are fully developed. Use the higher rate for large mature trees.

Multiflora Rose: Apply 3 pints of GLYPHOSATE 53.5% per acre as a broadcast spray or as a 3/4 percent solution with hand-held equipment. Treatments should be made prior to leaf deterioration by leaf-feeding insects.

Poison Ivy/Poison Oak: Apply 6 to 7 1/2 pints of GLYPHOSATE 53.5% per acre as a broadcast spray or as a 1 1/2 percent solution with hand-held equipment. Treatments must be applied before leaves lose green color. Repeat applications may be required to maintain control.

Trumpet Creeper: Apply 3 to 4 1/2 pints of GLYPHOSATE 53.5% per acre as a broadcast spray or as a 3/4 to 1 percent solution with hand-held equipment.

Willow: Apply GLYPHOSATE 53.5% as a 3/4 percent solution with hand-held equipment.

STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, seed or feed by storage and disposal.

STORAGE:

STORE ABOVE 10°F. (-12°C) TO KEEP FROM CRYSTALIZING

Crystals will settle to the bottom. If allowed to crystalize, place in a warm room 68°F. (20°C) for several days to redissolve and mix well before using.

DISPOSAL

Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or buried on site in a safe place away from water supplies. All disposal should be in accordance with applicable Federal, state or local procedures.

Emptied container retains vapor and product residue. Observe all labeled safeguards until container is destroyed. Do not reuse container, destroy when empty.

Triple rinse container, then puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY

DREXEL CHEMICAL COMPANY warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use as modified by the above. Drexel makes no other warranties, expressed or implied, including FITNESS or MERCHANTABILITY. In no case shall Drexel or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product, when such use and/or handling is not in strict accordance with directions given herewith. The foregoing is a condition of sale by DREXEL CHEMICAL COMPANY and is accepted as such by the Buyer.