

PM-21

Reg # 19713-380

49 16



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (H7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number:
19713-380

Date of Issuance:

NOTICE OF PESTICIDE:
 Registration
 Reregistration

Term of Issuance:
Conditional

Name of Pesticide Product:
Drexel Manzi Flowable

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Drexel Chemical Co.
P. O. Box 9306
Memphis, TN 38190-0306

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
2. Make the following labeling change before you release the product for shipment:
 - Add the phrase: "EPA Registration Number 19713-380."
 - Under "Foliar Applications" revise the last phrase in the first paragraph to read: "then the total poundage.... must not exceed any of the specified individual EBDC product maximum seasonal poundage of active ingredient allowed per acre."

Signature of Approving Official:

Cornelia P. Webb

Date:

7/20/95

29 16

-2-

3. Submit five copies of the final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

3916

ACCEPTED
WITH COMMENTS
to EPA Letter Dated

JUL 20 1970
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

DREXEL MANZI FLOWABLE
37% MANEB DISPERSION

ACTIVE INGREDIENT:

Maneb (Manganese ethylenebisdithiocarbamate)..... 37.0%
(Equivalent to manganese metallic 7.6%)

INERT INGREDIENTS:..... 63.0%

TOTAL:.....100.0%

This product contains 4 pounds Maneb per gallon.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

IF IN EYES: Flush eyes with plenty of water for at least 15 minutes. Get medical attention if irritation persists.

IF ON SKIN: Immediately flush skin with plenty of water. Get medical attention if irritation persists.

EPA Reg. No.: 19713-
EPA Est. No.: 19713-

Manufactured By:
Drexel Chemical Company
P.O. Box 13327
Memphis, TN 38113-0327

PRECAUTIONARY STATEMENTS

Hazard to Humans and Domestic Animals

CAUTION: May cause irritation of eyes, nose, throat and skin. Avoid breathing spray mist. Do not rub eyes or mouth with hands.

PERSONAL PROTECTIVE EQUIPMENT:

Applicators and other handlers (other than mixers and loaders) must wear:

- Coveralls over long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Mixers and Loaders must wear:

- Coveralls over long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Protective eyewear
- Chemical-resistant apron when mixing or loading

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering controls statement

When handlers use enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS:

Users should:

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. For terrestrial uses do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Cover or incorporate spilled treated seed. Do not contaminate water when disposing of equipment washwaters.

PHYSICAL AND CHEMICAL HAZARDS

Keep away from fire and sparks. Store in a cool dry place.

CHEMIGATION

Apply this product only through sprinkler irrigation systems including center pivot, lateral move, end tow, side (Wheel) roll, traveler, big gun, solid set, or hand move. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.

If you have any questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label prescribed safety devices for public water systems are in place.

SPECIFIC INSTRUCTIONS FOR SPRINKLER IRRIGATION SYSTEMS

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticide and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the areas intended for treatment.

Good agitation is required in the injection tank. In moving systems, apply specified dosage of product as a continuous injection. In nonmoving systems inject this product for 15 to 30 minutes at end of cycle. Use the least amount of water possible with uniform coverage.

Mix the amount of product needed for acreage to be treated into the quantity of water determined during prior calibration. For moving systems inject into the system continuously for one complete revolution of the field. For nonmoving systems inject into system for the time established during calibration.

Stop injection equipment after treatment is completed and continue to operate irrigation equipment until all product is flushed from system.

Specific Instructions for Public Water Systems:

Public water system means a system for the provision of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone backflow preventor (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top of the overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent the water fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injections pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

Coveralls over long-sleeved shirt and long pants
 Waterproof gloves
 Shoes plus socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The Worker Protection Standard applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep unprotected persons, children, livestock and pets away from treated areas or where there is danger of drift.

DIRECTIONS FOR USE AS A SPRAY

Manzi Flowable is approved for application in sufficient water to provide good coverage with available equipment in either dilute sprays or in concentrate ground or aerial sprays. Rates listed are based on 100 gallons of dilute spray unless otherwise noted. Rates of product for concentrate and aerial applications should be equal to the rates for dilute sprays on a per acre basis.

FOR GROUND APPLICATIONS use at the rate indicated in sufficient water for thorough coverage, minimum of 10 gallons. Increased volume of water may be necessary as foliage density increases.

FOR AERIAL APPLICATIONS use at the rate indicated in sufficient water for thorough coverage, a minimum of 3 gallons per acre for field crops and 10 gallons per acre for orchard crops. Apply with properly calibrated aerial equipment, arrange nozzles so that spray delivery is uniform over the entire spray swath. Human flaggers are prohibited unless in totally enclosed vehicles.

Where dosage rates are given, use the higher rate and shorter intervals under severe disease pressure, but do not exceed the maximum rate or apply more frequently than the minimum interval given in the directions for that crop.

FOLIAR APPLICATIONS

Where EBDC products used allow the same maximum poundage of active ingredient per acre per season:

If more than one product containing an EBDC active ingredient (maneb, mancozeb, or metiram) is used on a crop during the same growing season and the EBDC products used allow the same maximum poundage of active ingredient per acre per season, then the total poundage of all such EBDC products used must not exceed any of the specified individual per acre.

10 9 16

Where EBDC products used allow different maximum poundage of active ingredient per acre per season:

If more than one product containing an EBDC active ingredient is used on a crop during the same growing season and the EBDC products used allow different maximum poundage of active ingredient per acre per season, then the total poundage of all such EBDC products used must not exceed the lowest specified individual EBDC product maximum seasonal poundage of active ingredient allowed per acre.

SEED TREATMENT

In addition to the maximum number of foliar applications permitted by the formula stated above, a single application or seed treatment may be made on crops which have registered seed treatment uses.

AGRICULTURAL FRUIT AND NUT CROPS

CROPS	DISEASES	RATE QTS/A	MAX QTS/A PER SEASO	DIRECTIONS
<p>FOR POME FRUITS: Use either the 'Pre-Bloom' or 'Extended Application' schedules DO NOT COMBINE OR INTEGRATE THE TWO TREATMENT SCHEDULES</p>				
Apples	Rusts (Including Cedar Apple), Fabrea Leaf Spot, Flyspeck, Scab, Sooty Blotch	4.8 or (1.2 per 100 gal. in dilute spray)	19.2	PRE-BLOOM: Begin application at ¼ to ½ inch green tip and continue on a 7-10 day application schedule through bloom. Do not graze livestock in treated areas. It is recommended that this product be used in an integrated Pest Management Program.
		2.4 or (0.6 per 100 gal. in dilute spray)	16.8	EXTENDED APPLICATION OR TANK MIX: For implementation of IPM programs, applications based on tree-row volume or for use as a resistance management tool: begin applications at ¼ to ½ inch green tip and continue applications on a 7-10 day application schedule through the second cover spray or to within 77 days of harvest. Do not apply within 77 days of harvest. Do not graze livestock in treated areas. It is recommended that this product be used in an integrated Pest Management Program. Add spreader/sticker to spray mixture for cover sprays and tank mix with an effective systemic protectant/curative fungicide for more effective control of diseases.
Almonds	Brown Rot: (Blossom and Twig Blight), Scab, Shot-hole, fungus (Leaf Blight)	4.8 - 6.4 or (1.2 - 1.6 per 100 gal. in dilute spray)	25.6	Apply at popcorn, full bloom, and petal fall or every 7-10 days if bloom is staggered. Omit petal fall spray if only brown rot is present. Do not apply later than 5 weeks after petal fall or within 145 days of harvest. Do not allow livestock to graze in almond orchards.
Bananas	Sigatoka	1.6 - 2.4	24	Apply when leaves first appear and repeat as needed on a 14-21 day interval. May be applied up to the day of harvest.
Cranberries	Fruit Rot	3.8 - 4.8	14.4	Spray on a 7-10 day interval. Do not apply within 30 days of harvest.
Grapes	Black Rot, Bunch Rot, Deadarm, Downy Mildew	1.2 - 3.2 or (0.3 - 0.8 per 100 gal. in dilute spray)	19.2	East of the Rocky Mountains: Apply in sufficient water to provide thorough coverage starting when new shoots are ½ to 1½ inches long and continue at 7-10 day intervals until fruit is set. Do not apply within 66 days of harvest.
		1.2 - 2.0 or (0.3 - 0.5 per 100 gal. in dilute spray)	6	West of the Rocky Mountains: Apply when shoots are ½ to 1½ inches long and continue at 7-10 day intervals. Do not apply within 66 days of harvest except in California. California - Do not apply after bloom.
Kadota Figs	Surface Mold (Cladosporium), Surface Rot (Alternaria)	0.6 per 100 gals.	2.4	Apply when disease threatens in 100-400 gallons of water per acre. Do not apply within 10 days of harvest.
Papayas	Anthracnose, Phytophthora Fruit Rot	1.6 - 2.0 or (0.4 - 0.5 per 100 gal. in dilute spray)	28	Apply when disease first threatens and repeat at 14-21 day intervals. May be applied up to the day of harvest.

VEGETABLE CROP USES

CROPS	DISEASES	RATE QTS/A	MAX QTS/A PER SEASON	DIRECTIONS
Beans (Dry)	Anthracoese, Downy Mildew, Rust	1.2 - 1.6	9.6	Begin when plants are small. Spray on a 5-7 day interval. Do not apply within 30 days of harvest.
Broccoli Brussel Sprouts Cauliflower	Alternaria Leaf Spot, Downy Mildew	1.2 - 1.6	9.6	Begin when disease threatens. Spray on a 7-10 day interval. Do not apply within 7 days of harvest.
Cabbage Kohlrabi Chinese Cabbage (tight headed only)	Alternaria Leaf Spot, Downy Mildew	1.2 - 1.6	9.6	Plant beds and direct seeded fields: Spray on a 7-10 day interval. Do not apply within 7 days of harvest.
Chinese Cabbage (loose head)	Alternaria Leaf Spot, Downy Mildew	0.8 - 1.2	7.2	Plant beds and direct seeded fields: Spray on a 7-10 day interval. Do not apply later than 7 days before harvest in CA and 10 days in HI.
Corn (sweet corn, popcorn, sweet corn used for seed production)	Common Rust, Helminthosporium Blight	1.2	18	East of the MS - (including AR & LA) Use sufficient water for thorough coverage. Start applications when disease first appears and repeat at 3-10 day intervals. Add a recommended surfactant or spreader/sticker if needed for better coverage. Do not apply within 7 days of harvest. Do not feed treated forage to livestock.
			6	West of the MS - (except AR & LA) Use sufficient water for thorough coverage. Start applications when disease first appears and repeat at 3-10 day intervals. Add a recommended surfactant or spreader/sticker if needed for better coverage. Do not apply within 7 days of harvest. Do not feed treated forage to livestock.
Cucumbers	Alternaria (Macrosporium) Leaf Spot, Angular Leaf Spot, Anthracnose, Downy Mildew, Pythium Fruit Rot	1.2 - 1.6	12.8	Begin when disease threatens or plants begin to run. Spray on a 7-10 day interval. Do not apply within 5 days of harvest.
Eggplant	Anthracoese, Septoria Leaf Spot, Cladosporium Leaf Mold, Early Blight, Late Blight, Gray Leaf Spot (Stemphylium)	1.2 - 1.6	11.2 per crop	Begin at first fruit cluster and repeat at 7-10 day intervals. Do not apply within 5 days of harvest.
Kale	Alternaria Leaf Spot, Downy Mildew	1.2 - 1.6	3.2 per cutting	Begin when disease threatens and apply on a 7-10 day interval. Do not apply within 10 days of harvest.
Lettuce (Head and Leaf), Endive	Downy Mildew	1.2 - 1.6	9.6	Apply when disease appears. Spray on a 7-10 day interval. Do not apply within 10 days of harvest. Remove residues from head lettuce by stripping and trimming.
			6.4	California—Spray on a 7-10 day interval. Do not apply within 14 days of harvest.

13916
6

VEGETABLE CROP USES (CONTINUED)

CROPS	DISEASES	RATE QTS/A	MAX QTS/A PER SEASON	DIRECTIONS
Melons: cantaloupes, casabas, crenshaws, honeydews, watermelons	Alternaria Leaf Spot, Anthracnose, Cercospora Leaf Spot, Downy Mildew, Gummy Stem Blight	1.2 - 1.6	12.8	Apply as soon as plants begin to run or when disease first appears. Spray on a 7-10 day interval. Do not apply within 5 days of harvest.
Onions (Dry Bulb)	Botrytis Leaf Blight, Downy Mildew, Neck Rot, Purple Blotch	1.6 - 2.4	24	Begin applications when diseases are first reported in the area at 7 day intervals throughout the season. Do not apply within 7 days of harvest. Do not apply to exposed bulbs.
Onions (Green)	Botrytis Leaf Blight, Downy Mildew, Neck Rot, Purple Blotch	1.6 - 2.4	16.8	Begin applications when diseases are first reported in the area at 7 day intervals throughout the season. Do not apply within 7 days of harvest. Do not apply to exposed bulbs.
	Smut	2.4	2.4 (Per 29,000 linear ft. of row - 18 in. spacing)	Apply as a furrow drench at time of planting onion seeds. Use 75-100 gallons of water per acre.
Peppers	Anthracnose, Cercospora Leaf Spot (Frogeye Spot), Phytophthora Blight, Ripe Rot	1.2 - 2.4	14.4	East of the MS: Begin when disease threatens. Spray on a 7-10 day interval. Do not apply within 7 days of harvest.
		1.2 - 1.6	9.6	West of the MS: Begin when disease threatens. Spray on a 7-10 day interval. Do not apply within 7 days of harvest.
Potatoes	Early Blight, Late Blight	1.2 - 1.6	11.2	Begin when plants are 2 to 6 inches high. Apply on a 5-10 day interval. Do not apply within 14 days of harvest except for: CT, DE, FL, ME, MI, MA, NH, NY, OH, PA, RI, VT, WI where their PHI is 3 days. It is recommended that this product be used with an Integrated Pest Management Program. Vine kill should occur 14 days before harvest.
Potatoes (Seed Pieces)	Fusarium Seed, Piece Decay	0.8 qts. per 10 gal.		Dip whole or cut tubers. Spread in cool place if held before planting. Seed piece treatment only. Do not use treated seed piece for food, feed, or oil purposes.
Pumpkins	Angular Leaf Spot, Downy Mildew	1.2 - 1.6	12.8	Begin when disease threatens. Spray on a 7-10 day interval. Do not apply within 5 days of harvest.
Squash (Summer and Winter)	Anthracnose, Downy Mildew	1.2 - 1.6	12.8	Begin when plants start to run. Spray on a 7-10 day interval. Do not apply within 5 days of harvest.
Sugar Beets	Cercospora Leaf Spot	1.2 - 1.6	11.2	Apply when disease appears. Spray on a 7 - 10 day interval. Do not apply within 14 days of harvest.
Tomatoes (Greenhouse and Field)	Anthracnose, Cladosporium Leaf Mold, Early Blight, Late Blight, Gray Leaf Spot (Stemphylium), Septoria Leaf Spot	1.2 - 2.4	16.8	East of the MS: Begin at first fruit cluster and repeat on a 7-10 day interval. Do not apply within 5 days of harvest.
		1.2 - 1.6	6.4	West of the MS. Spray on a 7-10 day interval. Begin when first fruit forms. Do not apply within 5 days of harvest.

14916

COMMERCIAL ORNAMENTAL CROP USES (NOT FOR HOME GARDEN USE)

CROPS	DISEASES	RATE/100 GALS.	DIRECTIONS
Carnations	Asterana Leaf Spot, Anthracnose, Botrytis Blight	0.8 - 1.2 qts.	Begin when new growth starts. Repeat weekly.
Chrysanthemums	Ascochyta Ray Blight, Botrytis Petal Spot, Septoria Leaf Spot	12.3 fl. oz. plus 3/4 lbs. Cactan 50W	Apply twice weekly during the blooming season for Petal Spot and Ray Blight. For Septoria Leaf Spot apply weekly throughout the season.
Danias	Asterana Leaf Spot, Botrytis Blight	0.8 - 1.2 qts.	Begin when new growth starts. Repeat weekly.
Dogwood	Anthracnose	0.8 - 1.2 qts.	Begin when buds open. Repeat when bracts fall 4 weeks later and in late summer.
Gladiolus	Botrytis Blight, Curvularia and Stemphylium Leaf Spots	1.2 qts.	Begin when flower spikes are developing. Repeat 2 to 3 times at weekly intervals.
Grasses	Brown Patch	4.8 fl. oz. in sufficient water/1000 sq. ft.	Begin at first sign of disease. Repeat at 7-14 day intervals. Do not use on pasture or range grasses. Do not feed clippings to livestock. Do not use if grass is grown for seed.
	Dollar Spot	9.6 - 12.8 fl. oz. in sufficient water/1000 sq. ft.	
	Melting-Out	4.8 - 6.4 fl. oz. in sufficient water/1000 sq. ft.	
	Leaf, Stem and Strob Rusts	2.4 qts.	
Lilies	Botrytis Blight	0.3 - 1.2	Begin with new growth. Repeat weekly.
Pansies	Anthracnose	0.3 - 1.2	Begin with new growth. Repeat weekly.
Peonies	Asterana Leaf Spot, Botrytis Blight, Phytoconthora Blight	0.3 - 1.2	Apply to foliage and soil in early spring and early fall and 7-10 day intervals during the growing season.
Roses	Black Spot, Cercospora Leaf Spot (TX), Rust (CA)	0.3 - 1.2	Begin when first leaves unfold. Repeat at 7-10 day intervals.
Snapdragons	Rust	0.8 - 1.2	Begin with emergence. Repeat weekly.
Zinnias	Asterana Blight, Leaf Spot, Botrytis Blight	0.3 - 1.2	Begin with emergence. Repeat weekly.

STORAGE AND DISPOSAL

Store in a cool dry place. Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities by burning. If burned, stay out of smoke.

16 9 16

WARRANTY - CONDITION OF SALE:

OUR RECOMMENDATIONS FOR USE of this product are based upon tests believed reliable. Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith.

In no case shall Drexel or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product when such use and/or handling is not in strict accordance with directions given herewith. The foregoing is a condition of sale by Drexel Chemical Company and is accepted as such by the Buyer.