



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

October 01, 2024

Luz Chan
lchan@drexchem.com
DREXEL CHEMICAL COMPANY

Subject: Non-PRIA (Pesticide Registration Improvement Act) Labeling Amendment - Add direction use for soybeans grown for seed
Product Name: DREXEL DEFOL 5
Admin Number: 19713-388
EPA Receipt Date: 08/11/2021
Action Case Number: 00479150

Dear Luz Chan:

The amended labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable.

This approval does not affect any terms or conditions that were previously imposed on this registration. You continue to be subject to existing terms or conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one (1) copy of the final printed labeling before you release this product for shipment with the new labeling. In accordance with 40 CFR § 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR § 152.3.

Should you wish to add/retain a reference to your company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by EPA. If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains statements or claims substantially differing from statements or claims made in connection with obtaining a FIFRA section 3 registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

Your release for shipment of this product constitutes acceptance of these terms. If these terms are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6.

If you have questions, please contact Lydia Crawford via email at crawford.lydia@epa.gov.

Sincerely,

A handwritten signature in black ink that reads "Lydia Crawford". The script is elegant and cursive, with the first letters of each word being capitalized and larger than the rest of the letters.

Lydia Crawford, Biologist
HB, RD
Office of Pesticide Programs

ACCEPTED

10/1/2024

Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered under
EPA Reg. No. 19713-388

Drexel[®]

Defol 5[®]

Defoliant/Desiccant

ACTIVE INGREDIENT:

Sodium chlorate..... 42.3%

OTHER INGREDIENTS: 57.7%

TOTAL:100.0%

This product contains 5 pounds of Sodium chlorate per gallon.
Contains fire retardant.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See FIRST AID Below

[See Side [Back] Panel for FIRST AID]; [See Page ____ for FIRST AID]
[See Attached Booklet for [Full Use Instructions] Complete Directions for Use]
[See Container Labeling for FIRST AID and Full Use Instructions]

EPA Reg. No. 19713-388

EPA Est. No. 19713-XX-XXX

Net Content: _____ Gals. (_____ L)

FIRST AID

IF IN EYES:

- Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

IF SWALLOWED:

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything by mouth to an unconscious or convulsing person.

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15 to 20 minutes.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact CHEMTREC at 800-424-9300 for emergency medical treatment information.

Manufactured By:

Drexel Chemical Company
P.O. BOX 13327, MEMPHIS, TN 38113-0327
SINCE 1972

Defol, Surf-AC, Peptoil and the Drexel logo are trademarks or registered trademarks of Drexel Chemical Company.

388SP-1024*P

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Mixers and loaders for aerial applications must wear: Single layer clothing and waterproof gloves.

Mixers, loaders and applicators for manually-pressurized handwands must wear: Single layer clothing and waterproof gloves.

Mixers, loaders and applicators for mechanically-pressurized handguns must wear: Double layer clothing and waterproof gloves.

All other mixers, loaders, applicators, and handlers must wear: Long-sleeved shirt and long pants, waterproof gloves, shoes, and socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

See engineering controls for additional requirements.

Enclosed Cockpits/Cabs: Applicators must use an enclosed cockpit or enclosed cab that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(6)].

USER SAFETY RECOMMENDATIONS

Users should: 1) Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. 2) Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. 3) Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water when disposing of equipment washwater or rinsate.

NON-TARGET ORGANISM ADVISORY

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product through any type of irrigation system. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard (WPS), 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval (REI). The requirements in this box only apply to uses of this product that are covered by the WPS.

Do not enter or allow worker entry into treated areas during the REI of 12 hours.

PPE required for early entry to treated areas that is permitted under the WPS and that involves contact with anything that has been treated, such as plants, soil or water is: Coveralls, shoes and socks, and waterproof gloves.

APPLICATION

DEFOL 5 may be applied by aircraft or ground equipment. It is essential that the foliage be thoroughly covered. The specified dilution rates should be carefully observed. A non-ionic surfactant such as Surf-Ac® 820 or an emulsifiable crop oil such as Peptoil™ may enhance the results.

Precautions should be taken to avoid having spray or spray drift contact adjacent crops or desirable vegetation or they may be defoliated. Equipment should be washed before and after use and care should be taken so that product does not dry in tank or on spray nozzles. Washwater should not be allowed to contact desirable vegetation.

When using this product under dry land conditions, injury may occur to rotational crops.

Use Restriction:

- Do not mix with organophosphate insecticides.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.

BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size – Aircraft

- Adjust Nozzles - Follow nozzle manufacturers' recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

BOOM HEIGHT – Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

RELEASE HEIGHT - Aircraft

Higher release heights increase the potential for spray drift.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

OTHER STATE AND LOCAL REQUIREMENTS

Applicators must follow all state and local pesticide drift requirements regarding application of Sodium chlorate. Where states have more stringent regulations, they must be observed.

EQUIPMENT

All application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

SPRAY DRIFT ADVISORIES

Boomless Ground Applications:

Setting nozzles at the lowest effective height will help to reduce the potential for spray drift."

Handheld Technology Applications:

Take precautions to minimize spray drift.

MANDATORY SPRAY DRIFT MANAGEMENT

Aerial Applications:

- Do not release spray at a height greater than 10 feet above the ground or vegetative canopy, unless a greater application height is necessary for pilot safety.
- Applicators must select nozzle and pressure that deliver medium or coarser droplets in accordance with American Society of Agricultural & Biological Engineers Standard 641 (ASABE S641). If the windspeed is 10 miles per hour or less, applicators must use $\frac{1}{2}$ swath displacement upwind at the downwind edge of the field. When the windspeed is between 11 to 15 miles per hour, applicators must use $\frac{3}{4}$ swath displacement upwind at the downwind edge of the field.
- Do not apply when wind speeds exceed 15 mph at the application site. If the windspeed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters
- Do not apply during temperature inversions.

Ground Boom Applications:

- User must only apply with the release height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy.
- Applicators must select nozzle and pressure that deliver medium or coarser droplets in accordance with American Society of Agricultural & Biological Engineers Standard 572 (ASABE S572).
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

Boomless Ground Applications:

- Applicators must select nozzle and pressure that deliver medium or coarser droplets in accordance with American Society of Agricultural & Biological Engineers Standard 572 (ASABE S572).
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

DEFOLIATION

Ideally, defoliation is favored on clear, calm, sunny days with high temperatures and high humidity. If rain is anticipated within 24 hours, application should be delayed whenever possible. Defoliation may be slowed when daytime temperatures are below about 60°F and with some crops, especially with vigorous varieties, regrowth can occur.

In order to obtain acceptable defoliation results, the crop must be fully mature. In most cases, the signs of maturity include:

1. A slight discoloration of the foliage is apparent, often with a yellowish or reddish tinge.
2. Vegetative growth has slowed or ceased, with little or no secondary growth.
3. A slight downward pull on a leaf gives a clean break from the stalk.

Adequate fertility, ample soil moisture, good insect control and good weed control also favor defoliation. Plants which have been under stress as from drought or insect pressure give less reliable results because tough, wilted or damaged leaves do not permit the proper uptake of the chemical and tend to inhibit the formation of an abscission layer. Typically, under such circumstances, the leaves are desiccated but do not drop, which in turn, often causes problems with excessive trash during harvesting. For crops where there is a heavy canopy of foliage, obtaining thorough coverage is sometimes difficult. Whenever practical, the use of ground equipment should be considered. The higher dilution rates should be used and any other available methods employed which will assure that the plants are completely covered all the way to the bottom leaves. If coverage is incomplete, a second application may be necessary in order to defoliate the lower leaves and to prevent secondary growth.

DESICCATION

When the objective is to reduce crop moisture or to desiccate weeds prior to harvest, most of the points discussed under “*DEFOLIATION*” are applicable. Incomplete coverage will give erratic results, especially with vines such as Morningglory where the slender stems may be sheltered from the spray by the crop canopy, leaving the vines succulent and tough so they will become entangled in the harvesting machinery. Once again, larger spray volume will encourage more complete coverage and more reliable desiccation.

CROPS

| Crop | Rate Per Acre (qts.) | Water (gals.) |
|---|----------------------|----------------------------------|
| Chili peppers | 4 to 10 | Air: 5 to 15 Ground: 20 to 40 |
| SPECIFIC DIRECTIONS: Apply 10 days before harvest. To be used on Peppers for processing only. Consult a processor before applying. USE RESTRICTIONS: DO NOT apply more than 10 quarts of this product (12.5 lb. a.i.) per acre per application. DO NOT apply more than 10 quarts (12.5 lb. a.i.) per acre per year. Only 1 application is allowed per acre per year. Pre-Harvest Interval: 0 Days | | |
| Corn | 4.8 | Air: 5 to 7 Ground: 10 to 20 |
| SPECIFIC DIRECTIONS: Use as a harvest aid to desiccate trashy weeds in early maturing Corn. Apply at least 14 days before anticipated harvest date. Use the low dilution rates when weeds are small and the crop canopy is fairly open. Desiccation of Morningglory and other vines may be erratic. USE RESTRICTIONS: DO NOT apply more than 4.8 quarts of this product (6 lb. a.i.) per acre per application. DO NOT apply more than 4.8 quarts (6 lb. a.i.) per acre per year. Only 1 application is allowed per acre per year. DO NOT graze treated fields or feed fodder, forage or residual grain within 14 days of application. Pre-Harvest Interval: 0 Days | | |
| Cotton | 2.4 to 4.8 | Air: 5 to 10 Ground: 10 to 20 |
| SPECIFIC DIRECTIONS: Air Application – Apply this product in sufficient water to make 4 to 10 gals. total solution per acre. Use the lower rate when plants and weather conditions are ideal for defoliation. Swath width should not exceed wingspread. Altitude and speed of aircraft should be regulated to give sufficient spray coverage. DO NOT apply during periods of excessive winds as spray may drift and cause defoliation of adjoining Cotton or injury to other crops. DO NOT apply during the very high temperatures found in some areas at midday. Ground Application – Apply this product in sufficient water to make 15 to 30 gals. total solution per acre. Cotton defoliant should be applied when 70% of the bolls are open. Poor defoliation often occurs under unfavorable weather, soil moisture, fertility and plant conditions. Uniform defoliation within a field may sometimes be difficult to obtain because of | | |

irregular plant maturity. Satisfactory defoliation within a field may not be obtained when Cotton plants are suffering from lack of moisture or when plant growth is retarded by cold weather.

Flush out equipment with water after each day's work.

Apply by air or ground at least 7 days before crop harvest.

USE RESTRICTIONS: DO NOT apply more than 4.8 quarts of this product (6 lb. a.i.) per acre per application. DO NOT apply more than 4.8 quarts (6 lb. a.i.) per acre per year except in California. In California, DO NOT apply more than 9.6 quarts of this product (12 lb. a.i.) per acre per year. Only 1 application is allowed per acre per year, except in California, where 2 applications are permitted per acre per year. DO NOT graze treated areas or feed gin waste to livestock. Pre-Harvest Interval: 0 Days

| | | |
|-----------|-----|----------------------------------|
| Dry beans | 4.8 | Air: 5 to 10 Ground: 10 to 20 |
|-----------|-----|----------------------------------|

SPECIFIC DIRECTIONS: Thorough coverage is essential. Make application 7 to 10 days before anticipated harvest, longer if temperatures are cool.

USE RESTRICTIONS: DO NOT apply more than 4.8 quarts of this product (6 lb. a.i.) per acre per application. DO NOT apply more than 4.8 quarts (6 lb. a.i.) per acre per year. Only 1 application is allowed per acre per year. DO NOT graze treated fields or feed treated fodder or forage to livestock. Pre-Harvest Interval: 0 Days

| | | |
|------|-----|------------------------------------|
| Flax | 4.8 | Air: 5 (min.) Ground: 10 (min.) |
|------|-----|------------------------------------|

SPECIFIC DIRECTIONS: Use as a harvest aid to desiccate trashy weeds and to reduce moisture in the mature crop. Apply 7 to 10 days before anticipated harvest date. Thorough coverage is essential. Activity of the spray will be enhanced by the addition of a good, non-ionic surfactant at the rate of 1 quart per 100 gals. of spray.

USE RESTRICTIONS: DO NOT apply more than 4.8 quarts of this product (6 lb. a.i.) per acre per application. DO NOT apply more than 4.8 quarts (6 lb. a.i.) per acre per year. Only 1 application is allowed per acre per year. DO NOT graze treated fields or feed treated foliage and fodder for 14 days after application. Pre-Harvest Interval: 0 Days

| | | |
|---------------|------------|----------------------------------|
| Grain sorghum | 3.6 to 4.8 | Air: 4 to 10 Ground: 10 to 20 |
|---------------|------------|----------------------------------|

SPECIFIC DIRECTIONS: To reduce the moisture content of the grain while the crop is standing, apply 7 to 10 days before harvest. Thorough coverage is essential. Use the lower rates when grain moisture is low and the weather is clear and dry. Use the higher rate when conditions for desiccation are poor.

USE RESTRICTIONS: DO NOT apply more than 4.8 quarts of this product (6 lb. a.i.) per acre per application. DO NOT apply more than 4.8 quarts (6 lb. a.i.) per acre per year. Only 1 application is allowed per acre per year. Pre-Harvest Interval: 0 Days

| | | |
|------------|-----|----------------------------------|
| Guar beans | 4.8 | Air: 5 to 10 Ground: 10 to 20 |
|------------|-----|----------------------------------|

SPECIFIC DIRECTIONS: Make application 7 to 10 days before anticipated harvest date.

USE RESTRICTIONS: DO NOT apply more than 4.8 quarts of this product (6 lb. a.i.) per acre per application. DO NOT apply more than 4.8 quarts (6 lb. a.i.) per acre per year. Only 1 application is allowed per acre per year. DO NOT graze treated fields or feed treated fodder or forage to livestock. Pre-Harvest Interval: 0 Days

| | | |
|----------|-----|----------------------------------|
| Potatoes | 4.8 | Air: 5 to 10 Ground: 10 to 20 |
|----------|-----|----------------------------------|

SPECIFIC DIRECTIONS: To defoliate plants and facilitate harvest, apply 10 days before harvest.

USE RESTRICTIONS: DO NOT apply more than 4.8 quarts of this product (6 lb. a.i.) per acre per application. DO NOT apply more than 4.8 quarts (6 lb. a.i.) per acre per year. Only 1 application is allowed per acre per year. DO NOT apply under conditions of extreme heat during the middle of the day. Pre-Harvest Interval: 0 Days

| | | |
|------|------------|-----------------------------------|
| Rice | 3.6 to 4.8 | Air: 5 (min.) Ground: 10 to 20 |
|------|------------|-----------------------------------|

SPECIFIC DIRECTIONS: Use to facilitate harvest and reduce moisture in the fully mature seed heads and to desiccate immature Red rice in mature White rice. Use the lower rate when crop conditions and weather are right to enhance rapid drying. Apply 7 to 10 days before anticipated harvest. To avoid drift, do not apply during windy weather. Thorough coverage of the crop is essential to achieve good results.

USE RESTRICTIONS: DO NOT apply more than 4.8 quarts of this product (6 lb. a.i.) per acre per application. DO NOT apply more than 4.8 quarts (6 lb. a.i.) per acre per year. Only 1 application is allowed per acre per year. DO NOT graze treated fields or feed treated straw or crop residues within 14 days of application. Pre-Harvest Interval: 0 Days

| | | |
|-----------|-----|----------------------------------|
| Safflower | 4.8 | Air: 5 to 10 Ground: 20 to 30 |
|-----------|-----|----------------------------------|

| | | |
|--|-----|------------------------------------|
| <p>SPECIFIC DIRECTIONS: Use to facilitate harvest and reduce moisture in fully mature seed heads. Apply 7 or more days before normal harvest when Safflower heads are fully mature.</p> <p>USE RESTRICTIONS: DO NOT apply more than 4.8 quarts of this product (6 lb. a.i.) per acre per application. DO NOT apply more than 4.8 quarts (6 lb. a.i.) per acre per year. Only 1 application is allowed per acre per year. DO NOT graze treated fields or feed treated forage. Pre-Harvest Interval: 0 Days</p> | | |
| Southern peas | 4.8 | Air: 5 to 10 Ground: 10 to 20 |
| <p>SPECIFIC DIRECTIONS: Make application 7 to 10 days before anticipated harvest, longer if temperatures are cool. Thorough coverage is essential.</p> <p>USE RESTRICTIONS: DO NOT apply more than 4.8 quarts of this product (6 lb. a.i.) per acre per application. DO NOT apply more than 4.8 quarts (6 lb. a.i.) per acre per year. Only 1 application is allowed per acre per year. DO NOT graze treated fields or feed treated fodder or forage to livestock. Pre-Harvest Interval: 0 Days</p> | | |
| Soybeans | 4.8 | Air: 5 (min.) Ground: 20 (min.) |
| <p>SPECIFIC DIRECTIONS: Use as a harvest aid to dry weeds in early maturing Soybeans and to facilitate the harvest. Make application 7 to 10 days before anticipated harvest date when beans are mature and ready for harvest. For Soybeans grown for seed, apply at least 7 days prior to anticipated harvest date.</p> <p>USE RESTRICTIONS: DO NOT apply more than 4.8 quarts of this product (6 lb. a.i.) per acre per application. DO NOT apply more than 4.8 quarts (6 lb. a.i.) per acre per year. Only 1 application is allowed per acre per year. DO NOT graze treated fields or feed treated forage. Pre-Harvest Interval: 0 Days</p> | | |
| Sunflowers | 4.8 | Air: 5 to 10 Ground: 20 to 30 |
| <p>SPECIFIC DIRECTIONS: Use to facilitate harvest and reduce moisture in fully mature seed heads. When using ground application, apply with a high clearance sprayer. Make application 7 or more days before normal harvest when Sunflower heads are fully mature.</p> <p>USE RESTRICTIONS: DO NOT apply more than 4.8 quarts of this product (6 lb. a.i.) per acre per application. DO NOT apply more than 4.8 quarts (6 lb. a.i.) per acre per year. Only 1 application is allowed per acre per year. Pre-Harvest Interval: 7 Days</p> <p>DO NOT graze treated fields or feed treated forage within 14 days of application.</p> | | |
| Wheat | 4.8 | Air: 5 (min.) Ground: 10 (min.) |
| <p>SPECIFIC DIRECTIONS: Use as desiccant on Wheat.</p> <p>USE RESTRICTIONS: DO NOT apply more than 4.8 quarts of this product (6 lb. a.i.) per acre per application. DO NOT apply more than 4.8 quarts (6 lb. a.i.) per acre per year. Only 1 application is allowed per acre per year. Pre-Harvest Interval: 3 Days</p> | | |

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Storage should be under lock and key and secure from access by unauthorized persons and children. Storage should be in a cool, dry area away from any heat or ignition source. Do not stack over 2 pallets high. Move containers by handles or cases. Do not move containers from one area to another unless they are securely sealed. Keep container tightly sealed when not in use. Keep away from any puncture source. Avoid storage near water supplies, food, feed and fertilizer to avoid contamination. Store in original containers only. Unloading storage and work areas must be as clean as possible to prevent contamination if spilled. If the contents are leaking or material is spilled, follow these steps:

1. Contain spill. Absorb with a material such as sawdust, clay granules or dirt.
2. Collect and place in suitable containers for disposal.
3. Wash area with soap and water to remove remaining pesticide.
4. Follow washing with clean water rinse.
5. Wash contaminated clothing immediately.
6. Do not allow runoff to enter sewer or contaminate water supplies.
7. Dispose of waste as indicated below:

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING:

Nonrefillable Container (rigid material; ≤ 5 gallons): Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container one-fourth full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Nonrefillable Container (rigid material; > 5 gallons up to < 250 gallons): Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container one-fourth full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Dispose of empty container in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Refillable Container (≥ 250 gallons & Bulk): Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

WARRANTY—CONDITIONS OF SALE

OUR DIRECTIONS FOR USE of this product are based upon tests believed reliable. Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically directed and other influencing factors in the use of this product are beyond the control of the Seller. To the extent consistent with applicable law, Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith. To the extent consistent with applicable law, in no case shall the Manufacturer or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product when such use and/or handling is not in strict accordance with directions given herewith. The foregoing is a condition of sale by the Seller and is accepted as such by the Buyer.