



United States
Environmental Protection Agency
Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 19713-405	2. EPA Product Manager Ms. Cynthia Giles-Parker	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Drexel Captan 80W	PM# 21 Fungicide Branch	
5. Name and Address of Applicant (Include ZIP Code) Drexel Chemical Company, 1700 Channel Avenue Memphis, TN 38113 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____	NOTIFICATION MAR 5 1999
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.	
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.	

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

A copy of the printed label (405SP-0299) with the modified respirator statement as per PR Notice 98-9 is submitted, herewith. A copy of the certification statement as per PR Notice 98-10 is also submitted, herewith.

Thank you.

Section - III

1. Material This Product Will Be Packaged in:					
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____		
* Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt	No. per container
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/>	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)

Name Luz G Piwonka	Title Registration Manager	Telephone No. (include Area Code) (901) 774-4370
-----------------------	-------------------------------	---

Certification

I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.

6. Date Application Received
(Stamped)

2. Signature <i>Luz G. Piwonka</i>	3. Title Registration Manager
4. Typed Name Luz G Piwonka	5. Date 2/26/99

NOTIFICATION

MAR 5 1999



Captan 80W

Fungicide

ACTIVE INGREDIENTS:

Captan*	78.3%
Related derivatives	1.8%

OTHER INGREDIENTS: 19.9%

TOTAL: 100.0%

*Trichloromethylthio-4-cyclohexene-1,2-dicarboximide

KEEP OUT OF REACH OF CHILDREN

DANGER / PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See **FIRST AID** Below
Read Entire Label Before Using This Product

EPA Reg. No. 19713-405
EPA Est. No. 19713-GA-1 Net Contents: _____

FIRST AID

IF IN EYES: Flush eyes with plenty of water. Call a physician.

IF SWALLOWED: Contact physician, Poison Control Center or emergency clinic before inducing vomiting. Give water or milk to drink and induce vomiting by touching back of throat with finger, or by administering syrup of ipecac. Do not induce vomiting or give anything by mouth to an unconscious person. If medical advice cannot be obtained, take person and a copy of the label to the nearest emergency treatment facility.

IF ON SKIN: Wash with plenty of soap and water. See a doctor if irritation occurs or rash develops.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals.

DANGER: Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Avoid contact with skin and clothing. Avoid inhalation of dust or spray mist.

PERSONAL PROTECTIVE EQUIPMENT (PPE)
Applicators and other handlers (other than mixers and loaders) must wear: Long-sleeved shirt and long pants, waterproof gloves, shoes plus socks and protective eyewear.

Mixers and loaders must wear: Long-sleeved shirt and long pants, waterproof gloves, shoes plus socks, protective eyewear, chemical resistant apron when cleaning, mixing or loading and dust/mist filtering respirator (MSHA/NIOSH) approval number prefix TC-21C), or a NIOSH approved respirator with any R, P or HE filter.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should: 1) Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. 2) Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. 3) Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not clean equipment or dispose of equipment washwaters in a manner that will contaminate water resources.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), restricted entry interval (REI) and notification to workers. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval of 4 days.

EXCEPTION: For the last 48 hours of the REI, workers may enter the treated area to perform hand labor or other tasks involving contact with anything that has been treated, such as plants, soil or water, without time limit, if they wear the early-entry personal protective equipment listed below.

EXCEPTION: The restricted entry interval for Strawberries is 24 hours. After expiration of the 24 hour period, no personal protective equipment is required. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is: Coveralls, waterproof gloves, shoes plus socks and protective eyewear. Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

Read all precautions and directions for use before using. Use only for claims listed and only as specified on this label. In order that pesticide residues on food and forage crops will not exceed federal tolerances, use only at recommended rates and intervals and do not apply closer to harvest than specified. Do not apply or allow to drift to adjoining food, fiber or pasture crops. Drift of captan onto sensitive crops (e.g., D'Anjou Pears) can cause severe phytotoxicity and crop loss. Consult State Agricultural Experiment Stations or State Agricultural Extension Service for additional information, as the time of applications needed will vary with the local conditions.

Manufactured By:
Drexel Chemical Company
P.O. BOX 13327, N.E. 1st HIS, TN 38138-0327
SINCE 1972

COMPATIBILITY AND PLANT SAFETY: CAPTAN 80W can be combined safely and effectively at recommended dosage rates with most commonly used fungicides and insecticides, with the exception of oil and strongly alkaline materials. Alkaline materials such as spray lime, lime-sulfur and bordeaux mixtures will reduce the fungicidal activity of this product. Do not apply captan in combination with or immediately before or closely following oil sprays. Do not allow oil sprays on adjacent crops to drift onto crops which have been or will shortly be treated with captan. The time factor governing the safe interval between captan and oil sprays varies due to general climatic conditions, therefore, consult local agricultural spray programs and authorities to determine the proper timing. The use of spreaders which cause excessive wetting is not advised. Combinations with solvent formulations of organic phosphates should not be used. Combinations of captan and sulfur should not be used on crops sensitive to sulfur. Used at high rates or in drenching sprays, captan may cause a necrotic spotting of tender, immature leaves of certain varieties of apples, peaches, plums and cherries. This type of injury is most likely to occur in the early cover sprays during long periods of warm, cloudy, humid weather. To avoid the hazard of leaf spotting under such conditions, use captan and other spray materials at lowest recommended rates and avoid drenching trees. Applications can be made by aircraft or ground equipment (including concentrate and semi-concentrate equipment). Pour recommended amount of this material into nearly filled spray tank. Add balance of water. Maintain agitation during filling and spraying operations. Do not allow mixture to stand. Do not combine with emulsifiable liquids or wettable powders unless previous experience has proven them to be physically compatible and safe to plants. (Read "Compatibility and Plant Safety" section above). For aerial or concentrate spray applications, apply the same amount of this product per acre as would normally be applied for dilute spray applications. Apply aerial or concentrate sprays in sufficient water for coverage. Do not apply this product through any type of irrigation system.

GENERAL USE PRECAUTIONS

Except as specified, begin applications before or at first sign of disease and repeat as needed to maintain control, but observe use limitations. Unless otherwise specified, application can be made on the day of harvest. Maximum application is for a crop cycle. Crop cycle is defined as pre-bloom through post-harvest.

Apply the high rate and/or spray at shorter intervals when climatic conditions most favor disease(s). Apply the low rate and/or spray at longer intervals when climatic conditions least favor disease(s). If you are unaware of the climatic conditions favorable for disease(s) claimed for the specific use sites, you must consult with your State Agricultural Extension Service to learn of these conditions.

IMPORTANT: Read label carefully. Although most of the directions on this label may be followed nationwide, a few are limited to either the Eastern or Western U. S. Follow those directions for your growing area where specified.

FRUIT AND NUT CROPS

ALMONDS: Blossom and Twig blight, Brown rot, Leaf blight, Scab, Shot hole — Apply 2½ to 6½ pounds this product per acre in 20 to 300 gallons of water using ground equipment or in 5 to 20 gallons of water by air. Use 3¼ to 5½ pounds per acre when captan is used alone. Use 2½ to 3¼ pounds per acre in tank mixtures with fungicides having a similar spectrum. Apply at popcorn, bloom, petal fall stages and up to 5 weeks after petal fall. Do not apply within 30 days of harvest. Hulls may be fed to livestock. Do not apply more than 32½ pounds of this product per acre per crop cycle. See "General Use Precaution" section for definition of crop cycle.

APPLES (Eastern U.S.): Black rot (Frog-eye), Botrytis blossom end rot, Primary scab — Apply 5 pounds this product per acre in 20 to 400 gallons of water using ground equipment or in 5 to 20 gallons of water by air. Apply at 5 to 7 day intervals as needed to maintain control in pre-bloom, bloom, petal fall and first cover sprays. Blister rot, Black pox, Black rot, Botryosphaeria rot, Brooks fruit spot, Fly speck, Secondary scab, Sooty blotch — Apply 2½ to 5 pounds this product per acre in 20 to 400 gallons of water using ground equipment or in 5 to 20 gallons of water by air. Apply at 10 to 14 day intervals in second and later cover sprays. Do not apply more than 40 pounds of this product per acre per crop cycle. May be applied up to day of harvest. Powdery mildew — If Powdery mildew is a problem add 6 to 12 pounds sulfur per acre.

NOTE: Do not use this product in combination with or closely following or in alternation with wettable sulfur products on sulfur sensitive varieties of apples such as Red Delicious, Staymen, Baldwin, King, etc. as severe injury and defoliation may occur.

APPLES (Western U.S.): Primary scab — Apply 2½ to 5 pounds this product per acre in 20 to 400 gallons of water per acre using ground equipment or in 5 to 20 gallons of water by air. To reduce potential for disease resistance tolerance development to other fungicides having a similar spectrum, the lower rate may be used in tank mixtures.

(Pacific Northwest): Botrytis rot, Bull's eye rot — Apply 3¼ pounds this product per acre in 20 to 400 gallons of water using ground equipment or in 5 to 20 gallons of water by air. Make 1 or 2 applications with late cover sprays and one final spray prior to harvest. Do not apply more than 40 pounds of this product per acre per crop cycle. May be applied up to day of harvest. Secondary scab — In mid-Summer cover sprays, the dosage may be reduced to 2½ pounds per acre.

APRICOTS: Brown rot (Twig blight), Jacket rot — Apply 1½ to 3¼ pounds this product per acre in 20 to 250 gallons of water using ground equipment or in

10 to 20 gallons of water by air. Apply at red bud, bloom, 75% petal fall and in cover sprays. To reduce potential for tolerance development to other fungicides having a similar spectrum, use the lower rate in tank mixtures. Do not apply more than 15½ pounds of this product per acre per crop cycle. May be applied up to day of harvest.

BLUEBERRIES (Western U.S.): Berry rot or Botrytis gray mold, Mummy berry — Apply 1¼ to 3½ pounds this product per acre in 20 to 200 gallons of water by ground or in 5 to 20 gallons of water by air. Begin at mid-bloom, repeat at 7 to 10 day intervals until maturity. Do not apply more than 43¾ pounds of this product per acre per crop cycle. May be applied up to day of harvest.

BLUEBERRIES (Eastern U.S.): Berry rot or Botrytis gray mold, Mummy berry — Apply 3½ pounds this product per acre in sufficient water for thorough coverage or a minimum of 5 gallons of water by air. Start spray program when buds swell and earliest buds have loose scales. Repeat at 7 day intervals through blossom period. Repeat at 7 to 10 day intervals from late bloom. Do not apply more than 43¾ pounds of this product per acre per crop cycle. May be applied up to day of harvest.

CHERRIES (Eastern U.S.): Botrytis rot, Brown rot, Leaf spot — Apply 2½ pounds this product per acre in 20 to 200 gallons of water using ground equipment or in 10 to 20 gallons of water by air. Apply in pre-bloom, bloom, petal fall, shuck, cover and pre-harvest sprays. Application at 3 to 4 day intervals may be necessary during bloom to control Blossom blight. Repeat applications at 7 to 10 day intervals as needed to maintain control up to start of harvest. If Powdery mildew is a problem add 6 pounds sulfur per acre to the petal fall, shuck and early cover sprays. If sulfur is added, this product may be reduced to 1¼ pounds per acre in these sprays. Do not apply more than 17½ pounds of this product per acre per crop cycle. May be applied up to day of harvest.

Post-harvest Sprays: Leaf spot — Apply 2½ pounds this product per acre in 20 to 200 gallons of water using ground equipment. Apply immediately after harvest and repeat application in 10 to 14 days.

CHERRIES (Western U.S.): Blossom blight, Brown rot, Brown rot (Fruit), Leaf spot — Apply 1½ to 2½ pounds this product per acre in 20 to 200 gallons of water using ground equipment or in 10 to 20 gallons of water by air. Apply in pre-bloom, bloom, petal fall, shuck, cover and pre-harvest sprays. Do not apply more than 17½ pounds of this product per acre per crop cycle. May be applied up to day of harvest.

GRAPES (Except CA): Downy mildew, Leaf spot and Phomopsis cane, Suppression of Black rot — Apply 1¼ to 2½ pounds of this product per acre in 20 to 200 gallons of water using ground equipment or in 7 to 20 gallons of water by air; when shoots are ½ to 1½ inches long, when shoots are 3 to 5 inches long and when shoots are 9 to 12 inches long. Repeat just before bloom, immediately after bloom and continue at 10 to 14 day intervals as long as disease conditions persist. Use the lower rate when spraying less susceptible Grape varieties or when conditions are less favorable for disease development. Use the higher rate on susceptible Grape varieties and during periods of weather highly favorable for disease development. Do not apply more than 15 pounds of this product per acre per crop cycle. May be applied up to day of harvest.

GRAPES (CA): Bunch rot (Botrytis) — Apply 2½ pounds of this product per acre in 20 to 200 gallons of water using ground equipment or in 7 to 20 gallons of water by air. Apply 2 applications before bloom and one immediately after bloom. Repeat periodically making 3 cover applications before bunches close. Leaf spot and Phomopsis cane (current season infection) — Apply 2 to 2½ pounds of this product per acre in 20 to 200 gallons of water using ground equipment or apply 2½ pounds of this product per acre in 7 to 20 gallons of water by air. Apply first spray when green tissue begins to show, but before shoots are 1 inch long and repeat application when shoots are 6 to 8 inches long. Do not apply more than 15 pounds of this product per acre per crop cycle. May be applied up to day of harvest.

NECTARINES: Brown rot, Scab — Apply 2½ to 5 pounds of this product per acre in 20 to 250 gallons of water using ground equipment or in 10 to 20 gallons of water by air. To reduce the potential for tolerance development to other fungicides having a similar spectrum, use the lower rate in tank mixtures. Apply in full pink, bloom, petal fall, shuck, cover and pre-harvest sprays. Applications at 3 to 4 day intervals may be necessary during bloom to control Blossom blight. Repeat applications at 7 to 14 day intervals as necessary to maintain control. Continue applications throughout harvest if conditions favor Brown rot. If Powdery mildew is a problem, add 7½ pounds sulfur per acre to the petal fall and shuck sprays and early cover spray. If sulfur is added, this product may be reduced to 1½ pounds per acre in these sprays. Coryneum blight (Peach blight, Shot hole) — Apply 2½ to 5 pounds of this product per acre in 20 to 250 gallons of water using ground equipment or in 10 to 20 gallons of water by air. Apply in pink bud, full bloom, petal fall and cover sprays as necessary and as a post-harvest spray (but before leaves drop). Do not apply more than 30 pounds of this product per acre per crop cycle (including post-harvest sprays). Pre-harvest sprays may be applied up to day of harvest.

PEACHES: Brown rot, Scab — Apply 2½ to 5 pounds of this product per acre in 20 to 400 gallons of water using ground equipment or in 10 to 20 gallons of water by air. To reduce the potential for tolerance development to other fungicides having a similar spectrum, use the lower rate in tank mixtures. Apply at full pink, bloom, petal fall, shuck stages and in cover and pre-harvest sprays. When conditions are favorable make applications at 3 to 4 day intervals during bloom to control Blossom blight. Then repeat

application at 7 to 14 day intervals as needed to maintain control. Continue applications through harvest if conditions favor Brown rot. If Powdery mildew is a problem, add 12 pounds of sulfur per acre to the petal fall, shuck and early cover spray. If sulfur is added, this product may be reduced to 2½ pounds per acre in these sprays. Coryneum blight (Peach blight, Shot hole) — Apply 5 pounds of this product per acre in 20 to 400 gallons of water using ground equipment or in 10 to 20 gallons of water by air. Apply at pink bud, full bloom, petal fall stages and in cover sprays as necessary and as a post-harvest spray (but before leaf drop). Do not apply more than 40 pounds of this product per acre per crop cycle (including post-harvest sprays). Pre-harvest sprays may be applied up to day of harvest.

PLUMS, FRESH PRUNES (Western U.S.): Brown rot — Apply 3¾ pounds of this product per acre in 20 to 300 gallons of water using ground equipment or in 10 to 20 gallons of water by air. Use the lower rate when tank mixes with fungicides of similar spectrum of activity are used. Apply at green bud, popcorn, bloom and petal fall stages. Repeat in cover sprays as conditions warrant. Prune russet scab (Lacy scab) — Apply 2½ to 3¾ pounds of this product per acre in 20 to 300 gallons of water using ground equipment. Apply at full bloom. Do not apply more than 33¾ pounds of this product per acre per crop cycle. May be applied up to day of harvest.

PLUMS, FRESH PRUNES (EASTERN U.S.): Brown rot — Apply 3¾ pounds of this product per acre in 20 to 300 gallons of water using ground equipment or in 10 to 20 gallons of water by air. Apply in full pink, bloom and petal fall sprays. Repeat applications at 7 to 14 day intervals as needed to maintain control. Continue applications through harvest if conditions favor Brown rot. The addition of a neutral spreader has improved coverage. Do not apply more than 33¾ pounds of this product per acre per crop cycle. May be applied up to day of harvest.

STRAWBERRIES: Botrytis (Gray mold), Leaf spot — Apply 1½ to 3¾ pounds of this product per acre in sufficient water for thorough coverage by ground equipment or in 10 to 20 gallons of water by air. Begin applications when new growth starts in the Spring and before fruit starts to form. Repeat at 7 to 14 day intervals. Under conditions favorable to Fruit rot, continue applications through harvest period treating immediately after each picking. Do not apply more than 30 pounds of this product per acre per crop year. May be applied up to day of harvest. Restricted entry interval is 24 hours. If applying as a directed/banded spray, use band rate of this product according to the following formula:

$$\frac{\text{Plant bed width (inches)}}{\text{Row spacing (inches)}} \times \text{Broadcast rate per acre} = \text{Banded rate of this product per acre}$$

SPECIAL USES

PEACH PRE-PLANT ROOT DIP (CA): Preventative pre-plant dip treatment for Crown gall. Use 2½ pounds this product plus 3½ pints diluted sodium hypochlorite (5.25% household bleach) per 100 of gallons water. Wash nursery trees to remove soil from roots. Cut off all dormant buds and suckers in crown area and prune root system if necessary. Submerge the entire dormant tree for 5 minutes. Recharge dip during operation at a rate of 3½ pints diluted sodium hypochlorite per 100 gallons of water.

POST-HARVEST FRUIT APPLICATION: For control of various molds and Storage rots (Botrytis, Gloeosporium, Rhizopus). Use as a post-harvest dip or wash on the following fruits: **Apples, Cherries, Pears** — Use 1½ pounds this product per 100 gallons of water. Apply as a spray or in a dip tank. When used as a dip, recharge wash solution periodically when tank volume is reduced by 25%. Bring water back to volume and add 1½ pounds of this product for each 100 gallons added. At end of every 8 hour shift, empty tank, flush and charge with fresh dilution. Do not allow dip tank solution to stand overnight. Maintain continuous agitation during dipping operation.

DISPOSAL OF LEFTOVER POST-HARVEST TREATMENT MIXTURE: Left-over dip or spray mixtures containing captan may be used as a foliar spray for Apple and Cherry trees, (but not Pear trees) or for ornamental sites listed on the label. Observe all restrictions such as maximum pounds applied per application and season. When calculating application rates, if analytical services are not available to determine the exact quantity of captan remaining in the mixture, assume that the tank still contains 1½ pounds of this product per 100 gallons of water. If the dip or spray mixture contains other pesticides in addition to captan, refer to the product label(s) for information regarding disposal. Captan wastes are acutely hazardous to the eyes. Improper disposal of spray or dip tank mixtures is a violation of Federal Law. If the leftover dip or spray mixture cannot be disposed of in the manner prescribed above, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance concerning the disposal of spent or excess dip tank mixtures.

ORNAMENTALS

Use Precautions

Do not apply spray to ornamental plants listed below beyond the point of drip from the leaf surface. When applying as a drench, apply only sufficient mixture to wet the surface of the soil except when the dose is specified in terms of volume of mixture per square foot of area.

AZALEAS: Damping-off of cuttings — Use 2½ pounds this product per 100 gallons of water. Dip cuttings before bedding. Petal blight — Use 1¾ pounds this product per 100 gallons of water. Apply to soil area around plants and spray flowers just before bloom. Repeat at 7 to 14 day intervals through bloom.

BEGONIAS (Tuberous): Damping-off, Tuber rot — Use 2½ pounds of this product per 100 gallons of water. Dip tubers 30 minutes, drain and plant.

CAMELLIAS: Petal blight — Use ¾ pounds of this product per 100 gallons of water. Apply to drench soil around plants beginning when flowers start to open. Repeat at 7 to 10 day intervals through bloom.

CARNATIONS: Alternaria leaf spot, Rust — Use 1¼ pounds of this product per 100 gallons of water. Begin application at first sign of disease. Repeat at 7 to 10 day intervals. Shorten intervals during frequent rains and heavy dews. Damping-off of cuttings — Use 2½ pounds of this product per 100 gallons of water. Dip cuttings before bedding.

CHRYSANTHEMUMS: Botrytis flower blight, Septoria leaf spot — Use 1½ pounds of this product per 100 gallons of water. Apply at first sign of disease. Repeat at 7 to 10 day intervals. Damping-off of cuttings — Use 2½ pounds of this product per 100 gallons of water. Dip cuttings before bedding.

DICHONDRA (CA Only): White mold (*Sclerotium rolfsii*) — Use 1¼ pounds of this product per 100 gallons of water. Apply 1 gallon spray per 10 square feet. Make 2 to 3 applications at 7 day intervals.

GLADIOLUS (Corms): Corm rot and Decay, Damping-off — Use ½ to 1 pounds of this product per 10 gallons of water. Dip corms 20 to 30 minutes. Drain and plant.

GRASSES (Ornamental in non-pastured areas only): Brown patch, Brown spot on St. Augustine Grass, Damping-off, Leaf spot, Melting out, Seeding blight — Use 1¼ pounds of this product per 100 gallons of water. Apply 10 gallons spray per 1,000 square feet. Begin when growth starts in Spring. Repeat at 7 to 14 day intervals throughout season. Do not graze treated areas or feed clippings to livestock.

GRASSES (Lawn Seedbeds): Damping-off and other soil-borne diseases — Use ¾ to 1¼ pound of this product per 1,000 square feet or 1¼ pounds of this product per 100 gallons of water, using 15 gallons spray per 1,000 square feet. Cultivate into upper 3 to 4 inches before planting.

ROSES: Black spot, Botrytis blossom blight — Use 1¼ pounds of this product per 100 gallons of water. Begin at first growth or first sign of disease. Repeat at 7 to 14 day intervals and more frequently during frequent rains and heavy dews.

SOIL AND GREENHOUSE BENCH TREATMENT: Pre-plant treatment for damping-off, root rot diseases on seedlings or transplants of roses (and other shrubs, trees, flowers) and lawn seedbeds — Use 1¼ pounds of this product per 100 gallons of water at a rate of 15 gallons of spray per 1,000 square feet. Cultivate into upper 3 to 4 inches of soil before planting.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited. Do not reuse empty container.

STORAGE: Keep pesticide in original container. Keep container tightly closed when not in use. Protect from excessive heat. Store in a cool, dry place.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Do not reuse container. Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and Local authorities, by burning. If burned, stay out of smoke.

WARRANTY—CONDITION OF SALE

NOTICE TO BUYER AND USER: Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions or under abnormal use conditions or under conditions not reasonably foreseeable to Seller, and Buyer and User assume the risk of any such use. Seller disclaims all other warranties, expressed or implied, including any warranty of fitness or merchantability. Seller shall not be liable for consequential, special or indirect damages resulting from the use or handling of this product and Seller's sole liability and Buyer's and User's exclusive remedy shall be limited to the refund of the purchase price.