



U.S. ENVIRONMENTAL PROTECTION AGENCY
 Office of Pesticide Programs
 Registration Division (H7505C)
 401 "M" St., S.W.
 Washington, D.C. 20460

EPA Reg. Number: 19713-486
 Date of Issuance: MAY 14 1997

NOTICE OF PESTICIDE:
 Registration
 Reregistration

Term of Issuance: Conditional

(under FIFRA, as amended)

Name of Pesticide Product: Drexel Defol 750

Name and Address of Registrant (include ZIP Code):
 Mr. R. Franklin Handy
 Technical Director
 DREXEL CHEMICAL COMPANY
 P.O. Box 13327
 Memphis, TN 38113-0327

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.
 Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.

2. Make the following label changes:

a. Revise the EPA Registration Number to read, "EPA Reg. No. 19713-486".

3. Submit two copies of the revised final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

John Epler 5/14/97

CONCURRENCES

SYMBOL ▶	7505C						
SURNAME ▶	E.MINOR						
DATE ▶	May 6, 1997						

Labeling OK - no comments 2/2/97

**DREXEL
DEFOL 750**
Defoliant / Desiccant with Fire Retardant

ACTIVE INGREDIENT:

Sodium chlorate 52.0%

INERT INGREDIENTS: 48.0%

TOTAL: 100.0%

This product contains 7.5 pounds of sodium chlorate per gallon.

**KEEP OUT REACH OF CHILDREN
WARNING/AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Read Entire Label Before Using This Product
SHAKE WELL BEFORE USING

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Get medical attention.

IF IN EYES: Flush with plenty of water. Get medical attention.

IF ON SKIN: Wash with plenty of soap and water. Get Medical Attention if irritation persists.

EPA REG. NO. 19713-*UIA*
EPA EST. NO. 19713-MS-1
NET CONTENTS:

MANUFACTURED BY:
DREXEL CHEMICAL COMPANY
P.O. BOX 13327
MEMPHIS, TN 38113-0327

New Submission: 01/28/97
F:\label\bits\Defol75

ACCEPTED
with COMMENTS
In EPA Letter Dated

MAY 14 1997
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
19713-486

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
WARNING. Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Harmful if swallowed.
PERSONAL PROTECTIVE EQUIPMENT
Applicators and other handlers must wear: Coveralls over short-sleeved shirt and short pants, waterproof gloves, chemical-resistant footwear plus socks, protective eyewear, chemical-resistant headgear for overhead exposure, chemical-resistant apron when cleaning equipment, mixing, or loading and Dust / mist filtering respirator (MSHA / NIOSH approval number prefix TC-21C) Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.
 When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS
Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS
 For terrestrial uses, do not apply to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE
 It is a violation of federal law to use this product in a manner inconsistent with its labeling.
 Do not apply this product through any type of irrigation system.
 Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.
 For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval.

The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.


PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

Coveralls over short-sleeved shirt and pants, waterproof gloves, chemical-resistant footwear plus socks, protective eyewear and chemical-resistant headgear for overhead exposure

APPLICATION

DEFOL 750 may be applied by aircraft or ground equipment. It is essential that the foliage be thoroughly covered. The specified dilution rates should be carefully observed. A non-ionic surfactant such as Surf Ac 820 or an emulsifiable crop oil such as Peptoil may enhance the results. DEFOL 750 should not be mixed with insecticides or other organic materials, unless specifically recommended, because a fire or explosion may result.

Precautions should be taken to avoid having spray or spray drift contact adjacent crops or desirable vegetation, or they may be defoliated. Equipment should be washed before and after use and care should be taken so that product does not dry in tank or on spray nozzles. Wash water should not be allowed to contact desirable vegetation. When using DEFOL 750 under dryland conditions, injury may occur to rotational crops.



DEFOLIATION

Ideally, defoliation is favored on clear, calm, sunny days with high temperatures and high humidity. If rain is anticipated within 24 hours, application should be delayed whenever possible. Defoliation may be slowed when daytime temperatures are below about 60° F., and with some crops, especially with vigorous varieties, regrowth can occur.

In order to obtain acceptable defoliation results, the crop must be fully mature. In most cases, the signs of maturity include:

1. A slight discoloration of the foliage is apparent, often with a yellowish or reddish tinge.
2. Vegetative growth has slowed or ceased with little or no secondary growth.
3. A slight downward pull on a leaf gives a clean break from the stalk.

Adequate fertility, ample soil moisture, good insect control and good weed control also favor defoliation. Plants which have been under stress as from drought or insect pressure give less reliable results because tough, wilted or damaged leaves do not permit the proper uptake of the chemical and tend to inhibit the formation of an abscission layer. Typically, under such circumstances, the leaves are desiccated but do not drop, which, in turn, often causes problems with excessive trash during harvesting.

For crops where there is a heavy canopy of foliage, obtaining thorough coverage is sometimes difficult. Whenever practical, the use of ground equipment should be considered. The higher dilution rates should be used and any other available methods employed which will assure that the plants are completely covered all the way to the bottom leaves.

If coverage is incomplete, a second application may be necessary in order to defoliate the lower leaves and to prevent secondary growth.

DESICCATION

When the objective is to reduce crop moisture or to desiccate weeds prior to harvest, most of the points discussed under Defoliation are applicable. Incomplete coverage will give erratic results, especially with vines such as morningglory where the slender stems may be sheltered from the spray by the crop canopy, leaving the vines succulent and tough so they will become entangled in the harvesting machinery. Once again, larger spray volume will encourage more complete coverage and more reliable desiccation.

CROPS

CHILI PEPPERS - Apply 1.6 to 4 qts. per acre in 5 to 15 gallons of water by air or 20 to 40 gals. of water by ground equipment, 10 days before harvest. To be used on peppers for processing only. Consult a processor before applying.

CORN - As a harvest aid to desiccate trashy weeds in early maturing corn, use 3.2 qts. DEFOL 750 in 5 to 7 gallons of water per acre by air at least 14 days before anticipated harvest date. Use the low (5 gallon) dilution rates when weeds are small and the crop canopy is fairly open. Desiccation of morningglory and other vines may be erratic.

Do not graze treated fields or feed fodder, forage or residual grain within 14 days of application.

COTTON - Apply 1.6 to 2.4 gallons per acre in 5 to 10 gallons of water by air or 10-20 gallons of water by ground equipment no later than 7 days before harvest. The cotton plants should be mature with the youngest bolls expected to make cotton being at least 30 days old. Thorough coverage is essential. Do not graze treated fields or feed gin waste to livestock.

CUCURBITS (grown for seed): To facilitate harvest by drying out the vines, apply at a rate of 3.2 qts. per acre in 5 to 10 gallons of water by air, or apply 1 gallon per acre in 20 to 30 gallons of water by ground equipment. Do not graze treated fields within 14 days of application.

DRY BEANS - Apply at a rate of 3.2 qts. per acre in 5 to 10 gallons of water by air or 10 to 20 gallons by ground equipment. Thorough coverage is essential. Make application 7 to 10 days before anticipated harvest, longer if temperatures are cool. Do not graze treated fields or feed treated fodder or forage to livestock.

FLAX - As a harvest aid to desiccate trashy weeds and to reduce moisture in the mature crop, apply 3.2 qts. per acre in enough water to give thorough coverage, 7 to 10 days before anticipated harvest date. For air application, use a minimum of 5 gallons of water and for ground equipment a minimum of 10 gallons of water. Activity of the spray will be enhanced by the addition of good non-ionic surfactant at the rate of 1 quart per 100 gallons of spray. Do not graze treated fields or feed treated flax foliage and fodder for 14 days after application.

GRAIN SORGHUM - To reduce the moisture content of the grain while the crop is standing apply 2.4 to 3.2 qts. per acre in sufficient water to give thorough coverage, 7 to 10 days before harvest. Apply a total of 5 to 10 gallons of spray per acre by air and 10 to 20 gallons per acre by ground equipment. Use the lower rates when grain moisture is low and the weather is clear and dry. Use the higher rate when conditions for desiccation are poor.

GUAR BEANS - Apply at a rate of 3.2 qts. per acre in 5 to 10 gallons of water by air or 10 to 20 gallons of water by ground equipment. Make application 7 to 10 days before anticipated harvest date. Do not graze treated fields or feed treated fodder or forage to livestock.

POTATOES - To defoliate plants and facilitate harvest, apply 3.2 qts. of DEFOL 750 in enough water to provide thorough coverage either by aircraft or ground equipment. Apply 10 days before harvest. Do not apply under conditions of extreme heat during the middle of the day.

AIR APPLICATION: Use 5 to 10 gallons of water per acre.

GROUND APPLICATION: Use 10 to 20 gallons of water per acre.

RICE - To facilitate harvest and reduce moisture in the fully mature seed heads, and to desiccate immature red rice in mature white rice, apply by air 2.4 to 3.2 qts. per acre in 5 or more gallons of water. Use the lower rate when crop conditions and weather are right to enhance rapid drying. Apply 7 to 10 days before anticipated harvest. To avoid drift do not apply during windy weather. Thorough coverage of the crop is essential to achieve good results. Do not graze treated fields or feed treated straw or crop residues within 14 days of application.

SAFFLOWERS - To facilitate harvest and reduce moisture in the fully mature seed heads apply 3.2 qts. in 5 to 10 gallons of water by air or 20 to 30 gallons of water by ground equipment, 7 or more days before normal harvest when safflower heads are fully mature. Do not graze treated fields or feed treated forage.

SOUTHERN PEAS - Apply at a rate of 3.2 qts. per acre in 5 to 10 gallons of water by air or 10 to 20 gallons by ground equipment. Thorough coverage is essential. Make application 7 to 10 days before anticipated harvest, longer if temperatures are cool.

Do not graze treated fields or feed treated fodder or forage to livestock.

SOYBEANS - As a soybean harvest aid to dry weeds in early maturing soybeans and to facilitate the harvest, apply 3.2 qts. per acre. Make application 7 to 10 days before anticipated harvest date when beans are mature and ready for harvest. For air application use a minimum of 5 gallons of water per acre; for ground equipment apply a minimum of 20 gallons of water per acre.

Do not graze treated fields or feed treated soybean foliage.

SUNFLOWER - To facilitate harvest and reduce moisture in the fully mature seed heads apply 3.2 qts. in sufficient water to give good coverage, 7 or more days before normal harvest when sunflower heads are fully mature. Apply in 5 to 10 gallons of water per acre by air or in 20 to 30 gallons of water per acre by ground with a high clearance sprayer. Do not apply later than 7 days before harvest. Do not graze treated fields or feed treated forage within 14 days of application.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE INSTRUCTIONS: Storage should be under lock and key and secure from access by unauthorized persons and children. Storage should be in a cool dry area away from any heat or ignition source. Do not stack plastic containers over 2 pallets high. Move containers by handles or cases. Do not move containers from one area to another unless they are securely sealed. Keep container tightly sealed when not in use. Keep away from any puncture source. Avoid storage near water supplies, food, feed and fertilizer to avoid contamination. Store in original containers only. Unloading, storage and work areas must be as clean as possible to prevent contamination if spilled. If concentrate or spray solution is allowed to dry on flammable objects, the chlorate component can act as an oxidizer. If the contents are leaking or material is spilled follow these steps:

1. Contain spill, absorb with a material such as saw dust, clay granules or dirt.
2. Collect and place in suitable containers for disposal.
3. Wash area with water and soap to remove remaining pesticide.
4. Follow washing with clean water rinse.
5. Clothing contaminated with chlorate solution must be removed at once before solution dries. Dried contaminated clothes can be ignited easily. Wash contaminated clothing immediately.
6. Do not allow run off to enter sewer or contaminate water supplies.
7. Dispose of waste as indicated below:

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Plastic containers - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY - CONDITION OF SALE

OUR RECOMMENDATIONS FOR USE of this product are based upon tests believed reliable. Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith. In no case shall Drexel or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product when such use and/or handling is not in strict accordance with directions given herewith. The foregoing is a condition of sale by Drexel Chemical Company and is accepted as such by the Buyer.