

	U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (7505P) Ariel Rios Building 1200 Pennsylvania Ave., NW Washington, D.C. 20460		EPA Reg. Number: 19713-521	Date of Issuance: JAN 10 2008
	NOTICE OF PESTICIDE: <input type="checkbox"/> Registration <input checked="" type="checkbox"/> Reregistration (under FIFRA, as amended)		Term of Issuance: Unconditional	Name of Pesticide Product: Drexel Chlorpyrifos 15GR Agricultural Insecticide
Name and Address of Registrant (include ZIP Code): Drexel Chemical Company 1700 Channel Avenue Memphis, TN 38113				
<p>Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.</p> <p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>Based on your response to the Reregistration Eligibility Decision, EPA has reregistered the product listed above. This action is taken under the authority of section 4(g)(2)(C) of the Federal Insecticide, Fungicide, and Rodenticide Act, as amended. Reregistration under this section does not eliminate the need for continual reassessment of pesticides. EPA may require submission of data at any time to maintain the registration of your product.</p> <p>Generic data requirements that have been identified in the RED as confirmatory data for the active ingredient contained in the subject product will continue to be tracked and assessed in the Agency's continual reassessment of pesticides.</p> <p>The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable with the following comments:</p> <p>Please see page 2 for comments</p> <p>You must submit one (1) copy of the final printed label before you release the product for shipment. Products shipped after 12 months from the date of this Notice or the next printing of the label whichever occurs first, must bear the new revised label.</p> <p>A stamped copy of the label and acute toxicity review are enclosed for your records. Should you have any questions, please contact Registration Review Manager, Dr. Akiva Abramovitch at 703-308-8328.</p>				
Signature of Approving Official: Meredith F. Laws, Chief <i>Meredith F. Laws</i> Insecticide-Rodenticide Branch Registration Division (7505P)			Date: JAN 10 2008	

Label Comments:

- 1) Delete "Such as" for listed crops under Cole Crops.
- 2) Delete "Our recommendation" and start with "The use" under the Warranty Statement.
- 3) Revise "For control of various insects....." on the top of page 1 to "For control of **listed** insects....."
- 4) Under Citrus Orchards Floors add the disclaimer (except fire ants, pharaoh ants, carpenter ants and harvester ants) in the text and the table heading after ants and delete the footnote and the *.
- 5) The maximum application rate for tobacco is 2.0 lb ai/acre and not 2.025 lb ai/acre.
- 6) Add under Onions and restrictions that the maximum application rate is 1.0 lb ai/acre/season (per acre missing).
- 7) Delete centipede from the Asparagus table since the data is insufficient to support the use on this public health pest.

ACCEPTED
with COMMENTS
In EPA Letter Dated:
JAN 10 2008

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.
19713-521



Chlorpyrifos 15GR

Agricultural Insecticide

GROUP 1B: INSECTICIDE

For control of various insects infesting certain field and vegetable crops.

ACTIVE INGREDIENT:

Chlorpyrifos 15.0%

OTHER INGREDIENTS: 85.0%

TOTAL: 100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See FIRST AID Below

EPA Reg. No. 19713-521

Net Contents:

EPA Est. No. 19713-

50 Pounds

FIRST AID ORGANOPHOSPHATE

IF SWALLOWED:

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything by mouth to an unconscious or convulsing person.

IF IN EYES:

- Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15 to 20 minutes.

IF INHALED:

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.

Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this pesticide product (including health concerns, medical emergencies or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378.

NOTE TO PHYSICIAN: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. If exposed, plasma and red blood cell cholinesterase tests may indicate significance of exposure (baseline data are useful). Atropine, only by injection, is the preferable antidote. Oximes, such as 2-PAM/protopam, may be therapeutic if used early; however, use only in conjunction with atropine. In case of severe, acute poisoning, use antidote immediately after establishing an open airway and respiration.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed. Causes eye irritation. Avoid contact with eyes, skin or clothing. Avoid breathing dust.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are Barrier Laminate and Viton. For more information, following instructions in Supplement Three of PR Notice 93-7. If you want more options, follow the instructions for category G on an EPA chemical-resistance category selection chart.

Loaders, applicators and all other handlers must wear: Coveralls over long-sleeved shirt and long pants, chemical-resistant gloves, chemical-resistant footwear plus socks and a NIOSH-approved dust/mist filtering respirator with MSHA/NIOSH approved number prefix TC-21C or NIOSH-approved respirator with any R, P, or HE filter.

See Engineering Controls for additional requirements.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS

Pilots must use an enclosed cockpit in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(6)].

When applicators use closed cab motorized ground equipment in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should: 1) Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. 2) Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. 3) Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish, aquatic invertebrates, small mammals and birds. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when cleaning equipment or disposing of equipment washwater or rinsate.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read all "DIRECTIONS FOR USE" before applying.

Do not apply this product in a way that will contact workers or other persons either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.



Manufactured By:
Drexel Chemical Company

P.O. BOX 13327, MEMPHIS, TN 38113-0327

SINCE 1972

4 8 8

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard (WPS), 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE), restricted entry interval (REI), and notification of workers. The requirements in this box only apply to uses of this product that are covered by the WPS.

Exception: If this product is injected or incorporated into the soil, the WPS allows workers to enter the treated area if there will be no contact with anything that has been treated under certain circumstances.

Certified crop advisors or persons entering fields under their direct supervision under certain circumstances may be exempt from the early re-entry requirement pursuant to 40 CFR Part 170.

Do not enter or allow entry into treated areas during the REI. The REI for each crop is listed in the directions for use associated with each crop.

PPE required for early entry to treated areas that is permitted under the WPS and that involves contact with anything that has been treated, such as plants, soil or water is: Coveralls over short-sleeved shirt and short pants, chemical-resistant footwear plus socks, chemical-resistant gloves made of any water-proof material, and chemical-resistant headgear for overhead exposure.

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

GROUP 1B INSECTICIDE

INSECTICIDE RESISTANCE MANAGEMENT (IRM)

CHLORPYRIFOS 15GR contains a Group 1B insecticide. Insect and mite biotypes with acquired resistance to Group 1B may eventually dominate the insect/mite population if Group 1B insecticides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by this product or other Group 1B insecticides.

The following practices are recommended to delay development of insecticide resistance:

- Avoid consecutive use of insecticides with the same mode of action (same insecticide group) on the same insect species.
- Use tank mixtures or premix product containing insecticides with different modes of action (different insecticide groups), provided the products are registered for use.
- Base insecticide use on comprehensive Integrated Pest Management (IPM) programs.
- Monitor treated insect populations in the field for loss of effectiveness.
- Contact your Local Agricultural Extension Specialist, certified crop advisor, or Manufacturer for insecticide resistance management and/or IPM recommendations for the specific site and resistant pest problems.

RATES OF APPLICATION

Do not apply this product by aircraft at a rate greater than 1 lb. chlorpyrifos a.i. (6.6 lbs. of this product) per acre.

Rates of Application per 1,000 Feet of Row and Equivalent Acreage at Different Row Spacings								
This Product per 1,000 Feet of Row (ozs.)	Required Amount of This Product per Acre from Various Row Spacing							
	40"	38"	36"	34"	32"	30"	22"	18"
3.7	3.0	3.2	3.4	3.6	3.8	4.0	5.5	6.7
4	3.3	3.4	3.6	3.8	4.1	4.4	5.9	7.3
4.5	3.7	3.9	4.1	4.3	4.6	4.9	6.7	8.2
6	4.9	5.2	5.4	5.8	6.1	6.5	8.9	10.9
6.5	5.3	5.6	5.9	6.2	6.6	7.1	9.7	11.8
7.5	6.1	6.4	6.8	7.2	7.7	8.2	11.1	13.6
8	6.5	6.9	7.3	7.7	8.2	8.7	11.9	14.5
9	7.4	7.7	8.2	8.6	9.2	9.8	13.4	16.3
12	9.8	10.3	10.9	11.5	12.3	13.1	17.8	21.8
15	12.3	12.9	13.9	14.4	15.3	16.3	22.3	27.2
16	13.1	13.8	14.5	15.4	16.3	17.4	23.8	29.0

CALIBRATION OF EQUIPMENT

Note: The chart below lists recommended initial gauge settings for application of this product with one hopper opening per row. Be sure to check the actual application rate under your operating conditions.

1. Fill hopper.
2. Attach a plastic bag to tube opening.
3. Set planter to the initial settings shown on table.
4. Measure off 1,000 feet of row and drive planter the premeasured distance at desired speed.
5. Each bag should contain 6 to 8 ozs. of granules, depending upon desired rate.
6. If the result is over or under the desired rate, adjust settings and repeat the calibration.

Equipment Calibration and Calibration Settings for Different Types of Equipment													
Planter Type	Speed (mph)												
	4			5			6			7		8	
	This Product per 1,000 Feet of Row (ozs.)									8		16	
	8		16		8		16		8		16		
Gauge Setting													
Gandy ¹	21.4	30.2	23.7	32.4	26.0	36.0	27.7	39.0	30.2	41.0			
John Deere ¹ Max-Emerge ²	20	44	26	46	30	49	35	52	40	54			
John Deere ¹ 7000 Max-Emerge (Odd Nos. on Gate)	14	22	16	24	18	26	19	28	21	30			
John Deere ¹ 7000 Max-Emerge (Even Nos. on Gate)	17	30	20	33	24	35	26	36	28	38			
John Deere ² 71 Flexi-Planter and Older Planters	1	2	2	2	2	2	2	2	2	3			
	30	17	5	22	9	27	13	31	16	16			
John Deere ³ Max Emerge Plus	18	-	23	-	29	-	33	-	39	-			
Allis Chalmers ⁴ 70 Series	8	13	8	13	8	13	8	13	8	13			
Allis Chalmers ⁵ 78 & 79 Series	1	3	2	3	2	4	2	5	3	6			
	9.0	3.0	3.3	9.5	6.0	3.0	9.0	4.0	2.5	0			
Noble ¹ (New)	11	19	14	22	16	25	17	28	19	31			
White Planter	11	19	14	22	16	25	17	28	19	31			
Intl Harvester ⁶	1	3	2	3	2	4	2	5	3	6			
	9.0	3.0	3.3	9.5	6.0	3.0	9.0	4.0	2.5	0			
Buffalo All-Flex ⁶ (Fleischer Mfg.)	4.88	10	4.88	10	4.88	10	-	-	-	-			

¹ Gauge setting
² Gauge setting with range 1 & 2 - number is range notch
³ An application rate of 16 ozs. per 1,000 feet of row is not attainable with this equipment.
⁴ Gauge setting is constant regardless of speed.
⁵ Gauge setting shown with stem gates & dial settings - number shown is gauge dial
⁶ Number of turns open on the adjustment nut.

USE SITES

ALFALFA (MO only)

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the REI of 24 hours until PPE required for early entry is worn.

Apply this product at planting as an in-furrow treatment for suppression of target pests during establishment. Direct granules into the planter shoe with the seed, then place the applicator tube directly behind the planter shoe so that granules drop into seed furrow. Alternatively, place the granular band applicator behind planter shoe so granules fall on soil surface and into the open seed furrow and are covered with soil.

Pest(s)	This Product (Lbs./A)
Cutworms Grubs Wireworms	6.6

USE PRECAUTIONS AND RESTRICTIONS: Do not cut or graze treated Alfalfa within 21 days after application. Maximum single application rate is 1 lb. chlorpyrifos a.i. (6.6 lbs. of this product) per acre. Do not apply more than one application of this product per year.

ASPARAGUS (CA only)

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the REI of 24 hours unless PPE required for early entry is worn.

Apply this product as a postharvest ground application for suppression of target pests. Apply as a band over the entire crown area when Asparagus beds have been split (i.e. remove most of the soil from above the crowns). Cover area with soil the day of application. **NOTE:** Control may be reduced in soils with high organic matter content.

Pest(s)	This Product (Lbs./A)
Centipedes Symphylans	10

USE PRECAUTIONS AND RESTRICTIONS: Do not apply within 180 days of harvest. Do not apply more than 3 lbs. chlorpyrifos a.i. (20 lbs. of this product) per acre between harvests.

CITRUS ORCHARD FLOORS

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the REI of 5 days unless PPE required for early entry is worn.

Apply this product as a ground broadcast spray directed to the orchard floor to control foraging ants and suppress mounds. Use a suitable granular applicator, such as a cyclone fertilizer spreader, that uniformly broadcasts the granules over the orchard floor. This product may be custom-blended with granular fertilizers provided that the blended mixture can be uniformly applied to the orchard floor. Do not apply where weeds grow or other obstructions that may impede uniform coverage of the orchard floor. Foliar applications of Chlorpyrifos 4E-AG may be made in addition to orchard floor treatments of this product (see "USE PRECAUTIONS AND RESTRICTIONS").

Pest(s)	This Product (Lbs./A)
Ants*	6.6

USE PRECAUTIONS AND RESTRICTIONS:

- Do not allow meat or dairy animals to graze in treated areas.
- Do not apply within 28 before harvest.
- Do not apply more than 3 lbs. chlorpyrifos a.i. (20 lbs. of this product) per acre per year.
- Do not make more than 3 application of this product or other products containing chlorpyrifos per year (does not include foliar applications to Citrus trees).
- Do not make a second application within 10 days of any chlorpyrifos application to the orchard.
- Do not apply more than 1 lb. active ingredient chlorpyrifos (6.6 lbs. of this product) in a single application.

* Except Fire ants, Carpenter, Harvest and Pharaoh ants.

COLE CROP (Brassica) LEAFY VEGETABLES*, RADISH, RUTABAGA AND TURNIP

(*such as Bok choy, Broccoli, Broccoli raab, Brussels sprout, Cabbage, Cauliflower, Chinese broccoli, Chinese cabbage, Collards, Kale and Kohlrabi)

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the REI of 24 hours (3 days for Cauliflower) unless PPE required for early entry is worn.

For direct-seeded and transplanted crops, apply this product as an at-plant T-band as a 4-inch wide band centered over the row. This application requires a spreader or splitter on the end of the applicator drop tube. Placement behind the planter shoe and in front of the press wheel is recommended.

Crop/(PHI)*	Pest	This Product (ozs./1,000 ft. of row)	Maximum Single Application per Acre per Season** (Ozs. of This Product)		Maximum No. of Applications per Season
			40" row	20" row	
Rutabaga - 7 days; Cauliflower - 21 days; Broccoli, Brussels sprout, Cabbage, Chinese Cabbage (bok choy, napa), Collards, Kale, Kohlrabi and Turnip - 30 days	Root maggot	4.6 to 9.3	7.5	15	1

* Preharvest interval = Number of days between last application and harvest of treated crop.
** Use proportionate amount of this product for other row spacings not to exceed 2.25 lbs. chlorpyrifos a.i. (15 pounds of this product) per acre per season.

Radish

Apply as an at-plant in-furrow treatment. Place granules in seed furrow with seed at time of planting.

Crop/(PHI)*	Pest	This Product (ozs./1,000 ft. of row)	Maximum Single Application per Acre per Season* (Ozs. of This Product)	Maximum No. of Applications per Season
Radish - 7 days	Root maggot	3.3	18.3 (2.75 lbs. a.i./A)	1

USE PRECAUTIONS AND RESTRICTIONS: The maximum single application rate is 0.5 oz. chlorpyrifos a.i. (3.3 ozs. of this product) per 1,000 feet of row.

* Preharvest interval = Number of days between last application and harvest of Radish.

CORN (Field, Sweet, Corn Grown for Seed)

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the REI of 24 hours unless PPE required for early entry is worn.

This product is compatible with all ALS inhibitor herbicides applied in accordance with label recommendations. Refer to product labels for additional precautions and directions for use.

At-Plant T-Band Application

Apply 8 ozs. of this product per 1,000 feet of row as a T-band over an open seed furrow over the row behind the planter shoe, in front of the press wheel. In conventional and minimum-till corn, incorporate into the top one-half to one inch of soil using suitable equipment. A soil applied T-band treatment may be followed by post-applied herbicides such as Callisto. This product has demonstrated suppression of certain soil-borne pathogens that may result in physiological and agronomic advantages to Corn under environmental stress conditions when compared to Corn not treated with this product.

At-Plant In-Furrow Application

Apply 8 ozs. of this product per 1,000 feet of row at planting as an in-furrow treatment in conventional, minimum-till and no-till Corn. Direct granules into the planter shoe with seed, or place applicator tube directly behind planter shoe so that granules drop into seed furrow, or place granular band applicator behind planter shoe so that granules fall on soil surface and into the open seed furrow and are covered with soil.

Postplant Application

To control Corn rootworm larvae, apply 8 ozs. of this product per 1,000 feet of row at cultivation by placing granules at the base of the plant on both sides of the row just ahead of the cultivation shovels and covering the granules with soil. To control European or Southwestern corn borer larvae, apply this product in a band over the row so that granules are directed into the whorl, or use a postplant broadcast treatment. Consult your State Agricultural Experiment Station or Extension Service Specialist for proper time to treat and threshold information. Scouting for insect damage is strongly encouraged.

Postplant Broadcast Application

To control European or Southwestern corn borer, apply this product by uniformly broadcasting granules over corn plants by aerial application. For aerial applications, do not apply within 150 feet of Rivers, Natural ponds, Lakes, Streams, Reservoirs, Marshes, Estuaries, and Commercial fish ponds. Apply 5 lbs. of this product per acre for low to moderate first generation infestations and 6.5 lbs. of this product per acre for severe first generation infestations and all second generation infestations. Apply before larvae have entered stalks. Consult your State Agricultural Experiment Station or Extension Service for local threshold information. Scouting for insect damage is strongly encouraged.

Pest(s)	Banded/In-Furrow Applications (Ozs./1,000 feet of row)			Aerial Broadcast Application (Lbs./A)
	At-Plant		Postplant	
	T-Band	In-Furrow		
Armyworms	-	-	6 to 8	-
Billbugs	8	-	-	-
Chinchbugs ¹	8	8	-	-
Cutworms ¹	8	8	-	-
European and Southwestern corn borers ²				
First Generation	-	-	3.5 to 8	5 to 6.5
Second Generation	-	-	6 to 8	6.5
Ants ¹	8	8	-	-
Grubs	8	8	-	-
Lesser cornstalk borers	8	-	-	-
Northern and Western corn rootworm larvae	8	8	8	-
Seed corn beetles	8	8	-	-
Seed corn maggots	8	8	-	-
Southern corn rootworm larvae	8	8	8	-
Symphylans	8	-	-	-
Wireworms ³	8	8	-	-

SPECIFIC DIRECTIONS:

- ¹ The 8 ozs. application rate provides suppression for in-furrow treatment only of Ants (except Fire ants, Harvester and Pharaoh ants), Chinchbugs and Cutworms.
- ² Use 3.5 to 4 ozs. of this product per 1,000 feet of row for low to moderate first generation infestations of European corn borer before larvae have entered stalks when using postplant banded applications. For severe first generation and all second generation infestations, use 6 to 8 ozs. of this product per 1,000 feet of row. Apply before larvae have entered corn stalks.
- ³ With T-band applications for Wireworm control, consider using a hopper box insecticidal seed treatment. For best control, apply as an in-furrow treatment.

USE PRECAUTIONS AND RESTRICTIONS:

- Do not apply within 21 days of harvest of grain or ears.
- Do not apply by aircraft at a rate greater than 1 lb. chlorpyrifos a.i. (6.6 lbs. of this product) per acre.
- Do not make more than 3 applications of any product containing chlorpyrifos per season, including the maximum allowed of 2 granular applications, at the 1 lb. chlorpyrifos a.i. rate. Retreatment with a second soil application of this product is allowed under replant situations due to loss of crop during establishment only when initially application at the rate of 1 lb. chlorpyrifos a.i.
- Do not apply than 3 lbs. chlorpyrifos a.i. per acre per season.
- Do not make a second application of this product or other products containing chlorpyrifos within 10 days of the first application.
- Maximum single application rate for at-plant applications is 1.3 lbs. chlorpyrifos a.i. per acre (8 ozs. of this product per 1,000 feet of row).
- Maximum single application rate for postplant application is 1 lb. chlorpyrifos a.i. (6.6 lbs. of this product) per acre.
- If more than 1 lb. granular chlorpyrifos a.i. per acre is applied at-plant (for a maximum of 1.3 lbs. chlorpyrifos a.i. per acre per season), only 1 additional application of a liquid product containing chlorpyrifos a.i. per acre is allowed per season, for a total of 2.3 lbs. chlorpyrifos a.i. per acre per season.

ONIONS (Dry Bull)

Worker Restricted Entry Interval: Do not enter or allow worker entry in treated areas during the REI of 24 hours unless PPE required for early entry is worn.

Apply as an at-plant in-furrow treatment.

Pest(s)	This Product (Ozs./1,000 ft. of row)
Onion maggots	3.7

SPECIFIC DIRECTIONS:
CA, ID, WA and OR: To control maggots in onions planted in double rows with rows spaced 2 to 4 inches apart, apply 3.7 ozs. of this product per 1,000 feet of double row. Place granules in a 5- to 7-inch wide band over both rows behind the planter shoe and in front of the press-wheel to achieve shallow incorporation.

USE PRECAUTIONS AND RESTRICTIONS:

- Do not apply within 60 days before harvest.
- Do not apply more than 1 lb. chlorpyrifos a.i. (6.6 lbs. of this product) per season.
- The maximum single application rate is 0.56 oz. chlorpyrifos a.i. (3.7 ozs. of this product) per 1,000 feet of row.
- Do not make more than 1 application of this product or other products containing chlorpyrifos.

PEANUTS

Worker Restricted Entry Interval: Do not enter or allow worker entry in treated areas during the REI of 24 hours unless PPE required for early entry is worn.

At-Plant Preventive Application

Apply this product in a 6- to 12-inch band over the row behind the planter shoe and in front of the press wheel. Incorporate granules to a depth of 1 inch with tines, chains, or other suitable equipment. If 7.5 ozs. is applied at the time of planting, an additional application at postplant of 7.5 ozs. per 1,000 feet of row should be made to extend control.

Postplant Preventive Application

Apply this product at early flowering to pegging stage of growth in a 6- to 8-inch band over the row. For extended insect control and continued suppression of white mold (Southern blight), a second application of this product may be made (see "USE PRECAUTIONS AND RESTRICTIONS").

Band Rescue Application

Apply this product in a 10- to 18-inch band over the fruiting zone.

Pest(s)	Banded Applications (Ozs./1,000 ft. of row)	
	At-Plant	Postplant
Preventive Application: Cutworms Lesser cornstalk borer Southern corn rootworm larvae Wireworms White mold (Southern blight) ¹	7.5 to 15	7.5 to 15
Potato leafhopper	-	15
Rescue Application: Lesser cornstalk borer ²	-	7.5 to 15

SPECIFIC DIRECTIONS:

- ¹ For optimum suppression of White mold (Southern blight), apply 15 ozs. of this product per 1,000 ft. of row. Irrigation or rain following application is required to enhance treatment effectiveness. Under conditions of heavy white mold pressure, a suitable fungicide may also be required and must be applied separately.
- ² For control of Lesser cornstalk borer, use this product as a rescue treatment when the insect first appears, usually just before or at pegging stage of development.

USE PRECAUTIONS AND RESTRICTIONS:

- Do not apply within 21 days before harvest.
- Do not feed treated peanut forage or hay to meat or dairy animals.
- Do not apply more than 15 ozs. of this product per 1,000 ft. of row per crop season or apply more than 4 lbs. chlorpyrifos a.i. (26.6 lbs. of this product) per acre per season.
- Do not make more than 2 applications of this product per year.
- Do not make a second application of this product or any product containing chlorpyrifos within 10 days of first application.
- The combined total of preplant and postplant applications of this product and Chlorpyrifos 4E-AG or other products containing chlorpyrifos must not exceed 4 lbs. chlorpyrifos a.i. per acre per season.
- Aerial application is prohibited.

7 8 8

SORGHUM - GRAIN SORGHUM

Worker Restricted Entry Interval: Do not enter or allow worker entry in treated areas during the REI of 24 hours unless PPE required for early entry is worn.

Use this product as an at-plant T-band treatment in a 6- to 8-inch band over open seed furrow in front of press wheel into the top one-half to 1 inch of soil. Application rates of this product per acre for various row spacings are given in the "Rates of Application per 1,000 Feet of Row and Equivalent Acreage at Different Row Spacings" table.

Pest(s)	At-Plant T-Band Applications (Ozs./1,000 ft. of row)
Lesser cornstalk borer	4 to 8
Ants ¹	8
Chinchbugs ¹	
Corn rootworm	8
Cutworm	

SPECIFIC DIRECTIONS:

¹ For suppression of Chinchbugs and Ants (except Fire ants, Harvester and Pharaoh ants), use 8 ozs. of this product alone.

USE PRECAUTIONS AND RESTRICTIONS:

- Do not make more than one application of this product per season.
- The maximum single application rate per 1,000 feet of row is 1.3 lbs. chlorpyrifos a.i. in 30-inch row spacings (8 ozs. of this product). Proportional amounts for other row spacings should not exceed 1.5 chlorpyrifos a.i. per acre.
- Do not make a foliar application of Chlorpyrifos 4E-AG or any other product containing chlorpyrifos within 10 days of an at-plant application of this product.

SOYBEANS

Worker Restricted Entry Interval: Do not enter or allow worker entry in treated areas during the REI of 24 hours unless PPE required for early entry is worn.

Use this product as an at-plant T-band treatment in a 4- to 10-inch band over an open seed furrow in front of press wheel and incorporate in to the top one-half to 1 inch of soil. For postplant applications, incorporate granules in a 4- to 10-inch band to a depth of one-half to 1 inch using a suitable cultivator. Apply this product with equipment that will provide uniform distribution of granules over the row. Application rates of this product per acre for various row spacings are given in the "Rates of Application per 1,000 Feet of Row and Equivalent Acreage at Different Row Spacings" table.

Pest(s)	Banded Applications (Ozs./1,000 ft. of row)	
	At-plant T-Band	Postplant
Cutworm	8	8
Lesser cornstalk borer		
Ants* (Suppression only)	8	-

USE PRECAUTIONS AND RESTRICTIONS:

- Do not apply within 28 days of harvest.
- Do not apply as an in-furrow treatment.
- Do not make more than one application of this product and two additional applications per year of Chlorpyrifos 4E-AG or other liquid products containing chlorpyrifos.
- The maximum single application per 1,000 feet of row is 1.2 ozs. chlorpyrifos a.i. (8 ozs. of this product).
- Do not make a foliar application of Chlorpyrifos 4E-AG or other products containing chlorpyrifos within an at-plant application of this product.
- Do not apply more than 2 lbs. chlorpyrifos a.i. (13.3 lbs. of this product) per acre for preplant/at-plant incorporation and 1 lb. chlorpyrifos a.i. (6.6 lbs. of this product) per acre for foliar and postharvest applications.
- * Except Fire ants, Harvester and Pharaoh ants.

SUGARBEETS

Worker Restricted Entry Interval: Do not enter or allow worker entry in treated areas during the REI of 24 hours unless PPE required for early entry is worn.

At-Plant T-Band Application

Use this product in a band 4- to 5-inches wide behind the planter shoe, over the drill row, and in front of press wheel. **DO NOT APPLY GRANULES IN DIRECT CONTACT WITH SEEDS.** Apply 4.5 to 9 ozs. of this product per 1,000 feet of row (equivalent to 6.7 to 13.4 lbs. of this product per acre for 22-inch row spacings). Incorporate granules into the top one-half to 1 inch of soil using suitable equipment.

Postemergence Band Application

Apply this product in a 3- to 5-inch wide band over beet row (up to 2- to 4-true leaf stage of development). Apply 6.5 to 9 ozs. of this product per 1,000 feet of row (equivalent to 10 to 13.5 lbs. of this product per acre for 22-inch row spacings). Incorporate into the top one-half to 1 inch of soil using suitable equipment.

Pest(s)	Banded Applications (Ozs./1,000 ft. of row)	
	At-plant T-Band	Postplant
Sugarbeet root maggot ¹	4.5 to 9	6.5 to 9
Cutworm	6.6 to 9	-
Wireworms (Suppression only)	6.5 to 9	-

SPECIFIC DIRECTIONS:

¹ Apply 4.5 ozs. of this product per 1,000 feet of row (equivalent to 6.75 lbs. of this product per acre for 22-inch spacings) when populations of Sugarbeet root maggot are expected to be low. If initial adult fly activity indicates higher than anticipated populations, apply Chlorpyrifos 4E-AG at or near the time of peak adult emergence to augment control.

USE PRECAUTIONS AND RESTRICTIONS:

- Granular insecticides, including this product, may contribute to the stress of Sugarbeet plants under certain environmental conditions. Such stress may reduce plant stand or interfere with normal plant development. Herbicides used during preplant incorporation may interact with insecticides and enhance this stress.
- Do not apply within 30 days of harvest.
- Do not make more than one application of this product per season or apply more than 2 lbs. chlorpyrifos a.i. per acre per season.
- Do not apply more than a total of 3 lbs. chlorpyrifos a.i. (20 lbs. of this product) per acre per year, or make more than 3 applications of products containing chlorpyrifos per season.
- Do not make a foliar application of Chlorpyrifos 4E-AG or other products containing chlorpyrifos within 10 days of an at-plant application of this product.
- The maximum single application rate is 1.35 ozs. chlorpyrifos a.i. (9 ozs. of this product) per 1,000 feet of row or 2 lbs. chlorpyrifos a.i. (13.3 lbs. of this product) per acre for 22-inch spacings.

SUNFLOWER

Worker Restricted Entry Interval: Do not enter or allow worker entry in treated areas during the REI of 24 hours unless PPE required for early entry is worn.

Use this product as an at-plant T-band treatment. Place granules in a band 7 inches wide over the row behind the planter shoe in front of press wheel and incorporate into the top 1 inch of soil using suitable equipment.

Pest(s)	This Product (Ozs./1,000 ft. of row)
Cutworm	8

USE PRECAUTIONS AND RESTRICTIONS:

- Do not apply within 42 days of harvest.
- Do not make more than one application of this product and two additional applications per year of Chlorpyrifos 4E-AG or other liquid products containing chlorpyrifos.
- Do not make a foliar application of Chlorpyrifos 4E-AG or other products containing chlorpyrifos within 10 days of an at-plant application of this product.
- The maximum single application rate is 1.25 ozs. chlorpyrifos a.i. (8.3 ozs. of this product) per 1,000 feet of row or 1.3 lbs. chlorpyrifos a.i. (8.6 lbs. of this product) per acre per 30-inch spacings.
- Do not apply more than 2 lbs. chlorpyrifos a.i. (13.3 lbs. of this product) per acre for preplant/at-plant incorporation and 1 lb. chlorpyrifos a.i. (6.6 lbs. of this product) per acre for foliar and postharvest applications.

8 7 8

SWEET POTATO

Worker Restricted Entry Interval: Do not enter or allow worker entry in treated areas during the REI of 24 hours unless PPE required for early entry is worn.

Use this product as a preplant broadcast treatment by evenly distributing the granules over the soil surface, then incorporate granules into 2 to 4 inches of soil using a rotary hoe, disc cultivator or other suitable equipment. Plant Sweet potatoes no later 14 days after application. Delaying planting will reduce the length of time this product can protect crop against feeding damage. This product will not control false wireworm, whitefringed beetle or other grubs that attack Sweet potatoes.

Pest(s)	This Product (Lbs./A)
Wireworm (<i>Conoderus</i>), Flea beetle (<i>Systema</i>), Sweet potato flea beetle	13.5

USE PRECAUTIONS AND RESTRICTIONS:

- Do not apply within 125 days of harvest.
- Do not make more than one application of this product or other products containing chlorpyrifos per season.
- The maximum single application rate is 2 lbs. chlorpyrifos a.i. (13.3 lbs. of this product) per acre

TOBACCO

Worker Restricted Entry Interval: Do not enter or allow worker entry in treated areas during the REI of 24 hours unless PPE required for early entry is worn.

This product may be used as a preplant broadcast treatment one week before transplanting. Use equipment that evenly distributes granules over treated area. Incorporate granules 2 to 4 inches into the soil using suitable equipment immediately following application. Application of this product also suppresses imported fire ant movement into a treated field.

Pest(s)	This Product (Lbs./A)
Cutworms, Ants (suppression)*, Flea beetles, Mole crickets, Root maggots, Wireworms	13.5

USE PRECAUTIONS AND RESTRICTIONS:

- Do not apply within 7 days of harvest.
- Do not make more than one application of this product or other products containing chlorpyrifos per season.
- The maximum single application rate is 2.025 lbs. chlorpyrifos a.i. per acre.
- * Except Fire ants, Harvester and Pharaoh ants.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal. Open dumping is prohibited.

PESTICIDE STORAGE: Store in a cool, dry place.

PESTICIDE DISPOSAL: Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration or, if allowed by State and Local authorities, by burning. If burned, stay out of smoke.

WARRANTY — CONDITIONS OF SALE

OUR RECOMMENDATIONS FOR USE of this product are based upon tests believed reliable. Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller. To the extent consistent with applicable law, Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith.

To the extent consistent with applicable law, in no case shall the Manufacturer or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product when such use and/or handling is not in strict accordance with directions given herewith. The foregoing is a condition of sale by the Seller and is accepted as such by the Buyer.