

acceptance stamp
on 7/11

19713-556

10/07/2003

1/11



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (H7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number:
19713-556

Date of Issuance:
OCT 7 2003

NOTICE OF PESTICIDE:

Registration
 Reregistration

(under FIFRA, as amended):

Term of Issuance:

Conditional, with
Expiration Date of
March 21, 2007

Name of Pesticide Product:

Drexel Me-Too-Lachlor
III Herbicide

Name and Address of Registrant (include ZIP Code):

Drexel Chemical Company
P.O. Box 13327
Memphis, TN 38113

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Add the phrase "EPA Registration No.19713-556 to the label before you release the product for shipment.

COMMENTS CONTINUED ON PAGES 2 AND 2 OF THIS NOTICE OF REGISTRATION

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product under the enclosed stamped copy of the label constitutes acceptance of these conditions.

Enclosure

Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505C)

Signature of Approving Official:

Joanne I. Miller

Date: OCT 7 2003

Comments Continued:

- 2. Submit the following data required for the registration of this pesticide product within 36 months from March 21, 2002:

<u>EPA Guideline Data Number</u>	<u>Guideline Descriptor</u>
71-4(a)	Avian Reproduction in Bobwhite Quail
72-4(b)	Aquatic Invertebrate Life-Cycle
164-1	Turf Field Dissipation
132-1(a)	Foliar Residue Dissipation
133-3	Dermal Passive Dosimetry Exposure

- 3. Submit the following data required for the registration of this pesticide product within 48 months from March 21, 2002, of this requirement stated in the registration of Drexel Metolachlor Technical Herbicide, EPA Registration No. 19713-539:

<u>EPA Guideline Data Number</u>	<u>Guideline Descriptor</u>
166-1	Small Scale Prospective Ground Water Monitoring Study

- 4. Submit and/or cite all data required for the registration of this product when the Agency requires all registrants of similar products to submit data; and submit acceptable responses required for reregistration of this product under FIFRA, section 4.
- 5. Submit one (1) copy of the final printed labeling before you release this product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Enclosure

Drexel

Me-Too-Lachlor III

Herbicide

For weed control in Field Corn and Popcorn

ACTIVE INGREDIENT:

Metolachlor: 2-chloro- N-(2-ethyl-6-methylphenyl)-N-(2-methoxy-1-methylethyl) acetamide 84.1%

OTHER INGREDIENTS: 15.9%

TOTAL: 100.0%

This product contains 7.8 pounds of metolachlor active ingredient per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See FIRST AID Below
SHAKE WELL BEFORE USING

EPA Reg. No. 19713-

EPA Est. No. 19713-MS-1

Net Contents: _____

FIRST AID

IF INHALED:

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.
- Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15 to 20 minutes.
- Call a poison control center or doctor for treatment advice.

IF IN EYES:

- Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.
- Remove contact lenses, if present, after the first 15 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

IF SWALLOWED:

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything by mouth to an unconscious or convulsing person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

EMERGENCY PHONE NUMBERS:

- (800) 424-9300 CHEMTREC (transportation and spills)
- (800) 900-4044 Poison Control Center (human health)
- (800) 345-4735 ASPCA (animal health)

PRECAUTIONARY STATEMENTS

Hazards To Humans And Domestic Animals

CAUTION: Harmful if swallowed. Causes moderate eye irritation. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Avoid contact with eyes or clothing. Wear protective eye wear. Wear long-sleeved shirt and long pants, socks, shoes and gloves.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category H on an EPA chemical resistance category selection chart. **Applicators and other handlers must wear:**

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves such as barrier laminate or viton
- Chemical-resistant footwear plus socks
- Chemical-resistant headgear for overhead exposure
- Chemical-resistant apron when cleaning equipment

(Continued)

PRECAUTIONARY STATEMENTS (Con't)

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROL STATEMENTS

Mixers and loaders supporting aerial applications are required to use closed systems. The closed system must be used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4)]. When using the closed system, the PPE requirements for mixers and loaders may be reduced or modified as specified in the WPS.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should: 1) Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. 2) Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. 3) Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

Ground Water Advisory

This chemical is known to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

Surface Water Advisory

Metolachlor can contaminate surface water through ground spray drift. Under some conditions, metolachlor may also have a high potential for runoff into surface water (primarily via dissolution in runoff water), for several months post-application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas over-laying extremely shallow ground water, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas over-laying tile drainage systems that drain to surface water.

Mixing/Loading Instructions

Care must be taken when using this product to prevent back-siphoning into wells, spills, or improper disposal of excess pesticide, spray mixtures, or rinsates.

Check-valves or antisiphoning devices must be used on all mixing and/or irrigation equipment.

This product may not be mixed or loaded within 50 feet of perennial or intermittent streams and rivers, natural or impounded lakes and reservoirs. This product may not be mixed/loaded or used within 50 feet of all wells, including abandoned wells, drainage wells, and sink holes. Operations that involve mixing, loading, rinsing, or washing of this product into or from pesticide handling or application equipment or containers within 50 feet of any well are prohibited unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or wash water, and rain water that may fall on the pad. Surface water

Manufactured By:

Drexel Chemical Company

P.O. BOX 13327, MEMPHIS, TN 38113-0327

SINCE 1972

shall not be allowed to either flow over or from the pad, which means the pad must be self-contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times. The above-specified minimum containment capacities do not apply to vehicles when delivering pesticide shipments to the mixing/loading site.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves such as barrier laminate or viton
- Shoes plus socks

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND USE PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

To avoid spray drift, do not apply under windy conditions. Avoid spray overlap, as crop injury may result.

A. GENERAL INFORMATION

Observe all use precautions and limitations on the labels of each product used in tank mixtures. Tank mixtures are permitted only in those states where the tank-mix partner is registered.

Me-Too-Lachlor III Herbicide is a selective herbicide recommended as a preplant surface-applied, preplant incorporated, or preemergence treatment in water or fluid fertilizer for control of most annual grasses and certain broadleaf weeds in Field corn and Popcorn.

Note: Do not use in nurseries, turf, or landscape plantings.

Do not apply under conditions which favor runoff or wind erosion of soil containing this product to non-target areas.

To prevent off-site movement due to runoff or wind erosion:

- 1) Avoid treating powdery dry or light sand soils when conditions are favorable for wind erosion. Under these conditions, the soil surface should first be settled by rainfall or irrigation.
- 2) Do not apply to impervious substrates such as paved or highly compacted surfaces.
- 3) Do not use tailwater from the first flood or furrow irrigation of treated fields to treat non-target crops unless at least one-half inch of rainfall has occurred between application and the first irrigation.

Where directions on this label specify a tank mixture with atrazine follow the rates, recommendations, and limitations on the atrazine product label.

Note: Certain states may have established rate limitations for atrazine within specific geographical areas. Consult your state lead pesticide control agency for additional information. It is a violation of this label to deviate from state use regulations.

If this product is incorporated, any supplemental tillage before planting must not exceed the depth of incorporation.

Dry weather following preemergence application of this product or a tank mixture may reduce effectiveness. Cultivate if weeds develop. Where reference is made to weeds partially controlled, partial control can either mean erratic control from good to poor, or consistent control at a level below that generally considered acceptable for commercial weed control.

1) SOIL TEXTURES AND HERBICIDE RATES

Where rates are based on coarse-, medium-, or fine-textured soils, it is understood that soil textural classes are generally categorized as follows:

<u>Coarse</u>	<u>Medium</u>	<u>Fine</u>
Sand	Loam	Sandy clay loam
Loamy sand	Silt loam	Silty clay loam
Sandy loam	Silt	Clay loam
		Sandy clay
		Silty clay
		Clay

Within rate ranges in the rate tables and elsewhere on this label, use the lower rate on soils relatively coarse-textured or low in organic matter; use the higher rate on soils relatively fine-textured or high in organic matter.

Note: This product may be applied preemergence alone, or in combination with tank-mix partners specified on this label, following preplant incorporated herbicides when used according to their label recommendations, provided that such use is not prohibited on the respective labels.

Thoroughly clean sprayer or other application device before using. Dispose of cleaning solution in a responsible manner. Do not use a sprayer or applicator contaminated with any other materials, or crop damage or clogging of the application device may result.

2) MIXING INSTRUCTIONS

This Product Alone: Mix this product with water or fluid fertilizer and apply as a spray. Fill the spray tank one-half to three-quarters full with water or fluid fertilizer, add the proper amount of this product, then add the rest of the water or fluid fertilizer. Provide sufficient agitation during mixing and application to maintain a uniform emulsion.

Tank Mixtures: Fill the spray tank one-fourth full with water, and start agitation; add 2,4-D, Atrazine, Banvel[®], Lorox[®], Marksman[®], Sim-Trol[®], Prowl[®], or Atrazine + Sim-Trol, and allow it to become dispersed; then add this product, then add Gramoxone[®] Extra, Landmaster[®] BW, or Roundup[®], if these products are being used; and finally the rest of the water. For tank mixtures with Atrazine, Banvel, Lorox, Marksman, Sim-Trol, Prowl, or Atrazine + Sim-Trol, fluid fertilizers may replace all or part of the water as carrier, except in the Atrazine postemergence and the Banvel postemergence tank mixes. For tank mixtures with Atrazine, see additional mixing instructions on the Atrazine label. For each mixture, check compatibility with fluid fertilizer, as described in Appendix A, before mixing in spray tank. For all tank mixtures, agitate during mixing and application to maintain a uniform suspension.

*See Special Mixing Instructions for tank mixtures with Atrazine or Sim-Trol + Prowl under the appropriate tank mixture section.

For directions on how to conduct a compatibility test, see Appendix A.

3) APPLICATION PROCEDURES

Application Timing

This product alone or in some tank mixtures with other labeled herbicides may be applied for weed control in Field corn and Pop corn at various times. Refer to the given crop section of the label to determine if application timings listed below are recommended.

A) Preplant Surface-Applied: For minimum-tillage or no-tillage systems only, this product alone and some tank mixtures of this product may be applied up to 45 days before planting. Use only split applications for treatments made 30 to 45 days before planting, with two-thirds the recommended broadcast rate for the crop and soil texture applied initially and the remaining one-third at planting. Treatments less than 30 days before planting may be made either as a split or a single application. If weeds are present at the time of treatment, apply in a tank-mixture combination with a contact herbicide (for example, Gramoxone Extra or Roundup). Observe directions for use, use precautions, and restrictions on the label of the contact herbicide. To the extent possible, do not move treated soil out of the row or move untreated soil to the surface during planting, or weed control will be diminished.

B) Preplant Incorporated: Apply this product to the soil and incorporate into the top 2 inches of soil within 14 days before planting, using a finishing disk, harrow, rolling cultivator, or similar implement capable of providing uniform 2-inch incorporation. Use a preplant incorporated application if furrow irrigation is used or when a period of dry weather after application is expected. If Field corn and Pop corn will be planted on beds, apply and incorporate this product after bed formation, unless specified otherwise.

C) Preemergence: Apply this product during planting (behind the planter) or after planting, but before weeds or Field corn and Pop corn emerge.

4) SPECIAL APPLICATION PROCEDURES

CA Only (Field corn and Pop corn)—Preplant Incorporated: Broadcast this product alone or with tank mix partners listed on this label to the soil and thoroughly incorporate with a disk or similar implement set to till 4 to 6 inches deep. For more thorough incorporation, till the soil in 2 different directions (cross-till). Field corn and Pop corn may be planted on flat surface or on beds. Caution should be used when forming the beds that

only soil from this product's treated zone is used (i.e., untreated soil should not be brought to soil surface). If the application is made to preformed beds, incorporate this product with a tillage implement set to till 2 to 4 inches deep. Care should be taken during tilling to keep the tilled (this product treated) soil on the beds.

Preemergence: Apply this product after planting. Water with sprinkler or flood irrigation within 7 to 10 days.

Fall Application (Only in IA, MN, ND, SD, WI, and portions of NE and IL - see specific instructions for timing of application and other information): Do not apply to frozen ground. Use on medium and fine soils with greater than 2.5% organic matter that will be planted to Field corn or Pop corn the next Spring. Ground may be tilled before or after application. Do not exceed a 2 to 3-inch incorporation depth if tilled after treatment. **Note:** If a Spring application is made, the total rate of the Fall plus Spring applications must not exceed the maximum total rate for the specific crop, or illegal residues may result.

Ground Application: Apply this product alone or in tank mixtures by ground equipment in a minimum of 10 gals. of spray mixture per acre, unless otherwise specified.

Use sprayers that provide accurate and uniform application. For tank mixtures of this product with wettable powder or dry flowable formulations, screens and strainers should be no finer than 50-mesh. Rinse sprayer thoroughly with clean water immediately after use.

Calculate the amount of herbicide needed for band treatment by the formula:

$$\frac{\text{band width in inches}}{\text{row width in inches}} \times \text{broadcast rate per acre} = \text{amount needed per acre of field}$$

Note: For information on applying in lower volumes of carrier, see Low Carrier Application in Appendix B.

For application by air or through center pivot systems, see Appendices C and D. Appendix C includes Aerial Drift Management and Aerial Drift Reduction Advisory Information sections.

For information on impregnating dry fertilizer, see Appendix E.

B. THIS PRODUCT APPLIED ALONE

1) WEEDS CONTROLLED

Barnyardgrass (watergrass)	Foxtail millet	Signalgrass
Bristly foxtail	Galinsoga	(<i>Brachiaria</i>)
Carpetweed	Giant foxtail	Southwestern cupgrass
Common waterhemp	Goosegrass	Tail waterhemp
Crabgrass	Green foxtail	Wild proso millet*
Crowfootgrass	Pigweed	Witchgrass
Eastern black nightshade	Prairie cupgrass	Woolly cupgrass*
Fall panicum	Red rice	Yellow foxtail
Florida pusley	Robust foxtails (purple, white)	Yellow nutsedge

*For control of these weeds in Field corn and Popcorn only refer to the Woolly Cupgrass and Wild Proso Millet Control Program section of this label.

WEEDS PARTIALLY CONTROLLED**:

Common purslane	Sandbur	Volunteer sorghum
Eclipta	Seedling Johnsongrass	Wild proso millet
Florida beggarweed**	Shattercane	Woolly cupgrass
Hairy nightshade	Texas panicum***	

** See General Information section. Control of these weeds can be erratic, due partially to variable weather conditions. Control may be improved by following these suggested procedures:

- 1) In Field corn and Popcorn use 2 to 2.67 pts./A or the preplant surface-applied rates for this product alone or in tank mixture, if allowed, when making preplant incorporated or preemergence applications.
- 2) Thoroughly till moist soil to destroy germinating and emerged weeds. If this product is to be applied preplant incorporated, this tillage may be used to incorporate this product if uniform 2-inch incorporation is achieved as recommended under Application Procedures.
- 3) Plant Field corn and Pop corn into moist soil immediately after tillage. If this product is to be used preemergence, apply at planting or immediately after planting.
- 4) If available, sprinkler irrigate within 2 days after application. Apply one-half to 1 inch of water. Use lower water volume (one-half inch) on coarse-textured soils and higher volume (1 inch) on fine-textured soils. Also, refer to the section on Center Pivot Irrigation Application for this method of applying this product.
- 5) If irrigation is not possible and rain does not occur within 2 days after planting and application, weed control may be decreased. Under these conditions, a uniform, shallow cultivation is recommended as soon as weeds emerge.

** For partial control of this weed, use a minimum of 2 pts./A and apply preemergence.

*** For partial control of this weed, use a minimum of 2 pts./A and apply through a center pivot irrigation system.

2) ROTATIONAL CROPS

This Product Alone: (1) If Field corn or Popcorn crop treated with this product alone is lost, either Field corn or Popcorn may be replanted immediately. Do not make a second broadcast application of this product. If the original application was banded and the second crop is planted in the untreated row middles, a second banded treatment may be applied. (2) Barley, Oats, Rye, or Wheat may be planted 4.5 months following treatment; Alfalfa may be planted 4 months following application. Tomatoes may be planted 6 months following application. (3) Field corn and Pop corn, in addition to Cotton, Peanuts, Pod crops, Potatoes, Safflower, Grain or Forage sorghum, Soybeans, Root crops, Tobacco, Barley, Buckwheat, Milo, Oats, Rice, Rye, Wheat, Cabbage, or Peppers, may be planted in the Spring following treatment. Clover may be seeded 9 months following application. Do not graze or feed forage or fodder from Cotton to livestock. (4) Following a lay-by treatment or multiple treatments applied the previous season, Field corn and Pop corn in addition to Tobacco, Cabbage, or Peppers may be planted in the Spring. All other rotational crops may be planted 12 months after a lay-by application.

This Product In Tank Mixtures:

For Rotational Crop restrictions for this product used in tank mixtures, refer to the statements/restrictions above for this product and to the respective product labels of any mixing partner(s) for additional statements/restrictions.

Important Notes: To avoid injury to rotational Alfalfa or Clover: (1) Do not apply more than 2 lbs. a.i. of metolachlor per acre (2 pts. of this product) preemergence (including preplant surface, preplant incorporated, postplant incorporated, etc.), and (2) Do not make lay-by or other postemergent applications of this product.

C. FIELD CORN AND POP CORN — THIS PRODUCT ALONE

Apply this product either preplant surface, preplant incorporated or preemergence using the appropriate rate specified below.

1) PREPLANT SURFACE-APPLIED

Refer to instructions for use of this product alone under Application Procedures.

- a. Fall Application (Apply after September 30 in MN, ND, SD, WI and north of Route 30 in IA; Apply after October 15 north of Route 91 in NE and south of Route 30 in IA; Apply after October 31 north of Route 136 in IL):

In all locations, apply to crop stubble after harvest when the sustained soil temperature at a 4-inch depth is less than 55°F and falling. In minimum-till or no-tillage systems on soils having greater than 2.5% organic matter, use 1.67 to 2 pts./A on medium-textured and 2 pts./A on fine-textured soils. Do not apply to frozen ground. A tillage operation may precede the application. A Fall and/or a Spring tillage may follow application, but do not exceed an incorporation depth greater than 2 to 3 inches. Minimize furrow and ridge formation in the tillage operations. **Note:** If a Spring application is made, the total rate of the Fall plus Spring applications must not exceed the maximum total rate for Field Corn and Pop corn, or illegal residues may result.

- b. Use on medium- and fine-textured soils with minimum-tillage or no-tillage systems in CO, IA, IL, IN, KS, KY, MN, MO, MT, ND, NE, SD, TN, WI, and WY. Apply two-thirds of the recommended rate of this product (1.67 pts./A on medium soils and 2 pts./A on fine soils) as a split treatment 30 to 45 days before planting and the remainder at planting. Applications made less than 30 days prior to planting may be as either a split or single treatment. Apply 1.33 pts./A on coarse soils not more than 2 weeks prior to planting.
- c. On medium- and fine-textured soils with minimum- or no-tillage systems in CT, DE, MA, MD, ME, MI, NH, NY, OH, PA, RI, VA, VT, and WV, preplant surface applications may be applied following the directions for use above. If the amount of rainfall results in unsatisfactory length of weed control following the earlier treatment, a postemergence application of an appropriately labeled broadleaf and/or grass weed herbicide may be used, i.e., Atrazine, Beacon®, Bicep®, Exceed®, Accent®, Banvel®, Basagran®, bromoxynil (Brominal® or Buctril®), or 2,4-D. If the postemergence treatment includes the herbicide used preplant surface-applied, do not exceed the total labeled rate for Field corn and Pop corn on a given soil texture. Observe all directions for use, use precautions, and limitations on the label of the postemergent herbicide.

2) PREPLANT INCORPORATED OR PREEMERGENCE

Follow instructions for use of this product alone under Application Procedures. On coarse soils, apply 1 to 1.33 pts./A of this product if organic matter content is less than 3%, or 1.33 pts./A if organic matter content is 3% or greater. On medium soils, apply 1.33 to 1.67 pts./A of this product. On fine soils, apply 1.33 to 1.67 pts./A of this product if organic matter content is less than 3%, or 1.67 to 2 pts./A if organic matter content is 3% or greater.

6/11

Lay-by: To extend the duration of weed control in Field corn and Pop corn, a maximum rate of 2 pts./A of this product may be applied after Corn emergence until the Corn plants reach 40 inches in height, following any preplant surface-applied, preplant incorporated, or preemergence herbicide application, including this product. For best results, applications should be made to soil free of emerged weeds and directed towards the base of Corn plants in excess of 5 inches tall. The total rate of this product applied on Field corn and Pop corn during any one crop year should not exceed 4 pts./A, depending on soil texture.

Note for all applications to Field corn and Pop corn: To avoid possible illegal residues, do not graze or feed forage from treated areas for 30 days following application.

Shattercane, Wild Proso Millet, Woolly Cupgrass, and Eclipta — Partial Control: For more consistent partial control of Shattercane, Wild proso millet, Woolly cupgrass, or Eclipta, apply 2 to 2.55 pts./A as a single application; or apply 1 to 1.33 pts./A of this product preplant incorporated followed by 1 to 1.33 pts./A of this product preemergence; however, do not apply more than a total of 2.55 pts./A. Make the preemergence application during or after planting, but before weeds and Corn plants emerge. Apply the 1.33 pts./A rate of this product when a heavy infestation of Shattercane, Wild proso millet, Woolly cupgrass, or Eclipta is expected. A shallow cultivation may be needed to control any late emerging weeds.

Woolly Cupgrass and Wild Proso Millet Control Program: For control of these species, use the following 3-step program:

- 1) Apply this product early preplant, preplant incorporated, or preemergence at 1.67 pts./A on *medium soils* and 2 pts./A on *fine-textured soils* up to the maximum label rate. Lightly incorporate with a rotary hoe if rainfall does not occur within 5 to 7 days.
- 2) Apply a postemergence tank mix of Beacon at 0.38 oz./A or Exceed at 1 packet per 4 acres plus Accent SP at 0.33 oz./A plus 1 qt. of crop oil concentrate plus 1 gal./A of 28% nitrogen, or the equivalent amount of ammonium sulfate, when grasses are 2 to 3 inches tall and the Field corn and Pop corn is at least 4 inches tall.
- 3) Cultivate 14 to 21 days after the postemergence application.

Notes:

- 1) Do not apply more than the labeled application rate for a given soil texture per year, either as a single or split treatment, or illegal residues may result.
- 2) In Field corn and Pop corn, this product may be used up to 2.75 pts./A as either a preplant surface, preplant incorporated, or preemergence treatment on soils having an organic matter content between 6% and 20% or up to 2 pts./A on any soil for extended residual control and where severe stands of problem weeds are expected.
- 3) In the event of escape of annual weeds following a preplant surface, preplant incorporated, or preemergence treatment of this product, follow with a postemergence application of an appropriately labeled broadleaf and/or grass weed herbicide, i.e., Atrazine, Beacon, Bicep, Exceed, Accent, Banvel, Basagran, bromoxynil (Brominal or Butril), or 2,4-D. If the postemergence treatment includes the herbicide used in earlier treatment, i.e., Atrazine, do not exceed the total labeled rate for Field corn and Pop corn on a given soil texture.
- 4) Brominal or Butril may be applied postemergence alone or in tank-mix combination with Atrazine. Do not exceed 1.2 lbs. a.i./A of Atrazine in tank-mix combination with Brominal or Butril postemergence. Refer to the Atrazine, Brominal, and Butril labels for specific rates and use precautions.
- 5) Do not use this product on peat or muck soils.

D. FIELD CORN AND POP CORN - Me-Too-Lachlor III Herbicide Combinations

This product in any tank mixture for Field corn and Pop corn (except this product + Atrazine postemergence and this product + Banvel postemergence) may be applied in water or fluid fertilizer. Use only water in this product + Atrazine or this product + Banvel postemergence tank mixes.

Note: For all applications to Field corn and Pop corn, do not graze or feed forage from treated areas for 30 days following application, or possible illegal residues may result.

IMPORTANT: FOR TANK MIXTURES WITH ATRAZINE (OR OTHER BRANDS OF ATRAZINE) — If applying this product in tank mixture with Atrazine, all the restrictions and rate limitations on the Atrazine label must be followed if more restrictive/protective than those on this label. In addition, if Atrazine is/must be applied at rates lower than those recommended on this label, broadleaf weed control may be affected. Refer to the Atrazine label for weeds controlled at the reduced rates.

COMMENTS — Chart 1: This Product In Tank Mixtures

- 1) **Special Mixing Instructions** for this product + Atrazine or Sim-Trol and Prowl.
 - (1) Fill the spray tank one-quarter full with water or fluid fertilizer and start agitation. (2) To aid compatibility, add a compatibility agent, such as Unite or X-77[®], at 4 pts./100 gals. of spray mixture. (3) Then add the Atrazine or Simazine and allow it to become dispersed. (4) Then add this product and Prowl 4E. (5) Add the rest of the water.
- 2) Although a single formulation for Atrazine or Sim-Trol is listed in the rate tables, other formulations may be substituted using the following formula:
 - 1 lb. of Atrazine 90DF or Sim-Trol 90DF = 1.8 pts. of Atrazine 4L or Sim-Trol 4L

Chart 1: This Product In Tank Mixtures for Field corn and Pop corn — Additional Weeds Controlled and Special Instructions

	This Product + Atrazine and/or Sim-Trol (Preplant Surface, PPI, PRE)	This Product + Atrazine (Post)	This Product + Banvel (Field Corn)	This Product + Atrazine + Lorox	This Product + Atrazine or Sim-Trol + Prowl	This Product + Marksman
Section	D1	D2	D3	D4	D5	D8
Special Mixing Instructions					1	
Comments	2,3, 4,5, 7,8	2,3, 4,5		2,3, 4,5, 6	2,3, 4,5	7
Browntop panicum	a			a	a	
Cocklebur	a	b	b	a	a	a
Common purslane	a			a	a	a
Hairy nightshade	a			a	a	a
Jimsonweed		a	b			a
Kochia		a				a
Lambsquarters	a	a	a	a	a	a
Morningglory	a	b	b	a	a	a
Mustard		a				a
Pigweed				a	a	a
Prickly sida		a				a
Ragweed	a	a	a	a	a	a
Smartweed	a	a	a	a	a	a
Velvetleaf	a	a	b	a	a	a

a = control; b = partial control

COMMENTS — Chart 1: This Product Tank Mixtures for Corn (Cont.)

- 3) Although directions specify Atrazine formulations, in tank mixture with this product, other brands of atrazine may be used. Follow the rates, recommendations, and limitations on the Atrazine label.
- 4) See additional mixing instructions on the Atrazine label.
- 5) **Use Precautions:** Do not exceed a total of 2.5 lbs. a.i. of atrazine per acre per year. However, certain states may have established rate limitations for atrazine within specific geographical areas. Consult your state lead pesticide control agency for additional information. It is a violation of this label to deviate from state use regulations.
- 6) Other formulations of Lorox can be used:
 - 1 lb. of Lorox DF = 1 pt. of Lorox 4L
- 7) In Minimum-Tillage and No-Tillage systems, mix with Gramoxone Extra for control of most emerged annual weeds and suppression of perennial weeds; or with Landmaster BW for suppression of emerged Field bindweed and control or suppression of annual weeds; or with Roundup for control of most emerged annual and perennial weeds.
- 8) Refer to label section D7 on specific directions for 2,4-D or Banvel burndown combinations in Minimum-Tillage and No-Tillage systems.

This product in any tank mixture for Field corn and Pop corn may be applied in water or fluid fertilizer, except as noted.

Notes:

- 1) For all applications to Field corn and Pop corn, do not graze or feed forage from treated areas for 30 days following application, or possible illegal residues may result.
- 2) When applying this product in tank mixture with Atrazine, do not exceed a total of 2.5 lbs. a.i. of atrazine per acre per year.
- 3) Refer to **Section C. Field corn and Pop corn — This Product Alone, Note 3** for recommended sequential postemergence treatments if escape weeds develop.
- 4) In Field corn and Pop corn, this product may be used up to 2 pts./A in combinations on any soil for extended residual control and where severe stands of problem weeds are expected.

1. TANK MIXTURE WITH ATRAZINE OR SIM-TROL, OR ATRAZINE + SIM-TROL — PREPLANT SURFACE, PREPLANT INCORPORATED, OR PREEMERGENCE

In addition to the weeds controlled by this product alone, this product + Atrazine or Sim-Trol, or this product + Atrazine + Sim-Trol, applied preplant surface, preplant incorporated, or preemergence, also controls the following weeds: Browntop panicum, Cocklebur, Common purslane, Hairy nightshade, Lambsquarters, Morningglory, Ragweed, Smartweed, and Velvetleaf.

Apply this product + Atrazine or Sim-Trol, or this product + Atrazine + Sim-Trol either preplant surface, preplant incorporated, or preemergence.

Preplant Surface-Applied: Follow instructions for use of this product alone under **Application Procedures** and under application instructions for this product alone on Field corn and Pop corn. Apply this product + Atrazine or Sim-Trol, or this product + Atrazine + Sim-Trol on *medium soils* (1.67 pts./A of this product + 3.2 to 4 pts./A of Atrazine 4L or Sim-Trol 4L, or Atrazine 4L + Sim-Trol 4L combined) and on *fine soils* (1.67 to 2 pts./A of this product + 4 to 5 pts./A of Atrazine 4L, or Sim-Trol 4L, or Atrazine 4L + Sim-Trol 4L combined) in minimum-tillage and no-tillage systems in CO, IA, IL, IN, KS, KY, MN, MO, MT, ND, NE, SD, TN, WI, and WY. Apply the tank mixtures as a split or single treatment in those states and as indicated in the **This Product — Alone Preplant Surface-Applied** section of the label. On *coarse soils*, apply 1.33 pts./A of this product and 3.2 pts./A of Atrazine 4L or Sim-Trol 4L, or Atrazine 4L + Sim-Trol 4L combined.

Preplant Incorporated or Preemergence: Follow instructions for use of this product alone under **Application Procedures**. Apply this product + Atrazine or Sim-Trol, or this product + Atrazine + Sim-Trol, using the appropriate rates from Table 1.

Note: Do not apply more than the labeled rate for a given soil texture per year, either as a split or single treatment, or illegal residues may result.

Shattercane, Wild Proso Millet, Woolly Cupgrass, and Eclipta — Partial Control

For more consistent partial control of Shattercane, Wild proso millet, Woolly cupgrass, or Eclipta, where this product is applied in tank mixture or sequentially with other registered Field corn and Pop corn herbicides apply 2 to 2.33 pts. as a single application or, the following applications may be made:

- 1) Apply 1 to 1.33 pts./A of this product plus 2 lbs. per acre of Atrazine or Sim-Trol preplant incorporated, followed by 1 to 1.33 pts./A of this product preemergence. Make the preemergence application during or after planting, but before weeds and Corn emerge.
- 2) Apply this product at 1.33 pts./A alone or in tank mix combination with up to 2 lbs. per acre of Atrazine or Sim-Trol preplant incorporated. Do not exceed the total rate of triazine herbicide recommended for Field corn and Pop corn grown on a given soil texture. Follow with a post-directed application of Evik® 80W at 2.5 lbs./A. Refer to the Evik 80W label for specific directions for the post-directed application.
- 3) Apply Eradicane® or Sutan® (or equivalent EPTC or butylate formulations) at labeled rates preplant incorporated, followed by a preemergence application of this product at 1 to 1.33 pts./A. Do not use EPTC or Butylate on soils where rapid degradation has been shown to occur. Make the preemergence application during or after planting, but before weeds and Corn emerge.

Use Precaution: When following the application regimes in numbers 1 to 3 above, a shallow cultivation may be needed after the preemergence or postemergence application to help control any late emerging Shattercane or Wild proso millet plants.

2. TANK MIXTURE WITH ATRAZINE — POSTEMERGENCE

Weeds Controlled			Weeds Partially Controlled
Barnyardgrass (watergrass)	Green foxtail	Pigweed	Cocklebur
Crabgrass	Yellow foxtail	Prickly sida	Morningglory
Crowfootgrass	Jimsonweed	Purslane	Yellow nutsedge
Fall panicum	Kochia	Ragweed	
Giant foxtail	Lambsquarters	Smartweed	
	Mustard	Velvetleaf	

Apply 1 pt./A of this product + 1.3 lbs./A of Atrazine 90DF* on *coarse soils*, 1.33 pts./A of this product + 1.8 lbs./A of Atrazine 90DF on *medium soils*, or 1.33 to 1.67 pts./A of this product + 1.8 to 2.2 lbs./A** of Atrazine 90DF on *fine soils*. Apply this tank mixture before grass and broadleaf weeds pass the 2-leaf stage and before Field corn and Pop corn exceeds 5 inches in height. Application to weeds larger than the 2-leaf stage will generally result in unsatisfactory control.

Lay-by: Apply to Field corn and Pop corn plants not more than 12 inches tall. Applications to Corn in excess of 5 inches should be directed to the base of the Corn plants; whereas, applications to Corn plants less than 5 inches tall may be made over the top. Occasionally, some Corn leaf burn may result, but this should not affect later growth or yield. Do not apply this postemergence tank mixture in fluid fertilizer, or severe crop injury may occur.

*When using Atrazine 4L, use equivalent rates. One pound of Atrazine 90DF equals 1.8 pts. of Atrazine 4L.

**For better control of Cocklebur, Morningglory, Velvetleaf, and Yellow nutsedge on *fine-textured soils* above 3% organic matter, apply 2.2 lbs./A of Atrazine 90DF, or equivalent rate of Atrazine 4L, with 1.33 to 1.67 pts./A of this product.

Table 1: This Product + Atrazine or Sim-Trol, or This Product + Atrazine + Sim-Trol, Preplant Incorporated, or Preemergence — Corn (Field, Pop)

Soil Texture	Broadcast Rates Per Acre			
	Less Than 3% Organic Matter		3% Organic Matter or Greater	
	This Product + Atrazine 90DF* OR Sim-Trol 90DF*	- OR - This Product + Atrazine 90DF** + Sim-Trol 90DF**	This Product + Atrazine 90DF* OR Sim-Trol 90DF*	- OR - This Product + Atrazine 90DF** + Sim-Trol 90DF**
COARSE	0.85-1.0 pt. + 1.1-2.2 lbs.	0.85-1.0 pt. + 0.6-1.1 lbs. + 0.6-1.1 lbs.	1.0 pt. + 1.3-2.2 lbs.	1.0 pt. + 0.7-1.1 lbs. + 0.7-1.1 lbs.
MEDIUM	1.0-1.33 pts. + 1.3-2.2 lbs.	1.0-1.33 pts. + 0.7-1.1 lbs. + 0.7-1.1 lbs.	1.33 pts. + 1.8-2.2 lbs.	1.33 pts. + 0.9-1.1 lbs. + 0.9-1.1 lbs.
FINE	1.33 pts. + 1.8-2.2 lbs.	1.33 pts. + 0.9-1.1 lbs. + 0.9-1.1 lbs.	1.33-1.67 pts. + 1.8-2.2 lbs.***	1.33-1.67 pts. + 0.9-1.1 lbs.*** + 0.9-1.1 lbs.
Muck or Peat (soils with more than 20% organic matter)	DO NOT USE			
<p>*Use Sim-Trol in preference to Atrazine when heavy infestations of Crabgrass or Fall panicum are expected. On soils having between 6% and 20% organic matter, this product may be used up to 2.33 pts./A in tank mix combination with 2.2 lbs./A of Atrazine 90DF, or equivalent rates of Atrazine 4L. Refer to the Atrazine label for weeds controlled at this reduced rate.</p> <p>**When using the tank mixture of this product + Atrazine 90DF + Sim-Trol 90DF, use equal rates of each as shown when heavy broadleaf weed infestations are expected. When heavy infestations of Crabgrass or Fall panicum are expected, use a 1:2 ratio of Atrazine + Sim-Trol, instead of the 1:1 ratio given in Table 1. (Example: Total Atrazine 90DF + Sim-Trol 90DF = 1.2 lbs./A, use 0.4 lb. of Atrazine 90DF + 0.8 lb. of Sim-Trol 90DF respectively.) Refer to Comment No. 2 following Chart 1 for Sipcam Atrazine 4L and Sim-Trol 4L conversions.</p> <p>***For Cocklebur, Yellow nutsedge, and Velvetleaf control on <i>fine-textured soils</i> above 3% organic matter, apply 2.25 lbs./A of Atrazine 90DF, or equivalent rates of Atrazine 4L, or the same total amount of Atrazine + Sim-Trol 1.33 to 1.67 pts./A of this product.</p>				

Tank mixtures of this product + Atrazine may be applied following use of any registered preplant surface-applied, preplant incorporated, or preemergence Field corn and Pop corn herbicide, including this product + Atrazine.

Note: The total rate of this product should not exceed 4 pts., nor the Atrazine rate more than 2.5 lbs. a.i./A during any one crop year, or illegal residues may result. Refer to the Atrazine label for geographic, soil-texture, and rotational restrictions.

3. TANK MIXTURE WITH BANVEL

Preemergence: Use this tank mixture only on Field corn which is flat-planted (no furrows) in CO, IA, IL, IN, KS, MN, NE, OH, SD, and WI.

In addition to the weeds controlled by this product alone, this product + Banvel, applied preemergence, also controls Lambsquarters, Ragweed, Smartweed, Cocklebur*, Jimsonweed*, Morningglory*, and Velvetleaf*.

*Partially controlled.

Apply this product + Banvel preemergence. Broadcast 1 pt./A of Banvel with 1.33 pts./A of this product on *medium soils*, or with 1.33 to 1.67 pts./A of this product on *fine soils*. Do not apply on *coarse soils* or on soils with less than 2.5% organic matter. Apply this tank mixture to the soil surface at planting or after planting, but before Field corn emerges. Plant seed at least 1.5 inches deep and apply behind planting equipment, avoiding incorporation by the planter wheel or other seed-covering device. Do not incorporate before Field corn plants emergence. If it is necessary to rotary hoe to break the soil crust, do not disturb the soil more than one-half inch deep.

Use Precautions: (1) Avoid drift to sensitive non-target plants, such as Soybeans, during application, or injury may occur. (2) Do not apply with aircraft.

Postemergence for Control of Pigweed (Mid-Atlantic states including DE, MD, PA, VA, and WV): Apply 1 to 1.5 pts. of this product + one-half to 1 pt./A of Banvel by ground equipment when Pigweed plants are less than 3 inches tall and before Field corn exceeds 5 inches in height in a minimum of 20 gals. of spray per acre. Use the lower rate on coarse-textured and low organic matter soils. Use the higher rate on fine-textured and high organic matter soils.

Banvel SGF* and Clarity* may be used at equivalent pounds of active ingredient per acre.

Use Precautions: (1) Avoid drift to sensitive non-target plants, such as Soybeans, during application, or injury may occur. (2) Do not apply with aircraft.

4. TANK MIXTURE WITH ATRAZINE AND LOROX FOR CONTROL OF LAMBSQUARTERS AND PIGWEED

For prolonged control of Lambsquarters and Pigweed in DE, MD, NJ, NY, PA, VA, and WV, this product may be applied preemergence in tank-mix combination with Atrazine + Linuron. Apply this product and Atrazine according to the rates in Table 1 and Linuron according to the following rates.

Soil Texture	Broadcast Rate per Acre
Sandy loam (1-3% organic matter)	0.67 lb. Lorox
Sandy loam (3-6% organic matter)	1.0 lb. Lorox
Medium- and fine-textured soils (1-6% organic matter)	1.0 lb. Lorox

Observe all directions for use, use precautions, and limitations on this product, Atrazine and Lorox labels when applying these products in tank-mix combinations.

5. TANK MIXTURE WITH ATRAZINE OR SIM-TROL + PROWL FOR PROLONGED CONTROL OF LAMBSQUARTERS AND PIGWEED IN FIELD CORN ONLY (NORTHEAST U.S., INCLUDING MI, IN, KY, AND STATES EAST OF THESE)

For prolonged control of Lambsquarters and Pigweed in addition to a broad spectrum of annual broadleaf and grass weeds, this product in tank mix combination with Atrazine* or Sim-Trol + Prowl may be applied after planting, but before Field corn or weeds emerge. Apply by ground equipment in a minimum of 10 gals. of water or 20 gals. of liquid fertilizer. Apply by air in a minimum of 5 gals. of water. Refer to Table 1 of this label for rates of this product, Atrazine, or Sim-Trol to be applied. Apply Prowl 4E according to the following rates below in Table 2.

*Do not apply this product in tank mix combination with Atrazine 80W + Prowl, as this combination is not compatible. Other Atrazine formulations may be used.

Mixing Instructions: See Tank Mixture with Atrazine-Postemergence.

Table 2: Prowl 4E — Broadcast Rates Per Acre

Soil Texture	Percent Organic Matter in Soil		
	Less Than 1.5%	1.5-3%	Over 3%
COARSE	1.5-2.0 pts.	2.0 pts.	3.0 pts.
MEDIUM	2.0 pts.	3.0 pts.	3.0 pts.
FINE	2.0 pts.	3.0 pts.	3.0 pts.

Observe all directions for use, use precautions and limitations on the respective product labels when applying these products in tank mix combination. Refer to the Prowl 4E label for replanting instructions in the event of crop loss.

6. TANK MIXTURE WITH ATRAZINE OR SIM-TROL, OR ATRAZINE + SIM-TROL, WITH GRAMOXONE EXTRA, LANDMASTER BW, OR ROUNDUP FOR MINIMUM-TILLAGE OR NO-TILLAGE SYSTEMS

*See Comment No. 1 following Chart 1 for special mixing instructions.

In minimum-tillage or no-tillage systems where Field corn and Pop corn is planted directly into a cover crop, stale seedbed, established sod, or previous crop residues, the contact herbicides Gramoxone Extra, Landmaster BW, or Roundup may be added to a tank mix of this product, Atrazine, or Sim-Trol, or this product + Atrazine + Sim-Trol. See Comment No. 7 following Chart 1. This product + Atrazine or Sim-Trol, or this product + Atrazine + Sim-Trol portion of the tank mixture provides preemergence control of the weeds listed on this label in the tank mixture section for This Product + Atrazine or Sim-Trol, or This Product + Atrazine + Sim-Trol-Preplant Surface, Preplant Incorporated, or Preemergence.

Application: Apply before, during, or after planting, but before the Field corn and Pop corn emerges, at the rates specified below. Add Gramoxone Extra, Landmaster BW, or Roundup at the following broadcast rates:

Gramoxone Extra: 1.5 to 2, 2 to 2.5, or 2.5 to 3 pts./A to 1 to 3, 3 to 6, or 6-inch tall weeds, respectively. Apply surfactant at 1 or 2 pts./100 gals. of spray mixture with 75% or greater or 50 to 74% nonionic active ingredient, respectively. This treatment will not control weeds taller than 6 inches.

Note: Do not apply combinations containing Gramoxone Extra in suspension-type liquid fertilizers because the activity of paraquat will be reduced.

Landmaster BW: 27 to 54 ozs./A depending on weed species and size. See the Landmaster BW label for weeds controlled, recommended rates for specific weeds, and other information concerning use.

Roundup: See the Roundup or Roundup RT label for weeds controlled, recommended rates, and other use precaution directions.

Apply in 20 to 60 gals. of water or fluid fertilizer per acre with ground equipment.

On *coarse soils*, apply 1 pt./A of this product with 1.3 lbs. of Atrazine 90DF* or Sim-Trol 90DF*, or with 0.7 lb. of Atrazine 90DF** + 0.7 lb. of Sim-Trol 90DF**. On *medium soils*, apply 1.33 pts./A of this product with 1.8 lbs. of Atrazine 90DF or Sim-Trol 90DF, or with 0.9 lb. of Atrazine 90DF + 0.9 lb. of Sim-Trol 90DF. On *fine soils****, apply 1.33 to 1.67 pts./A of this product with 1.8 to 2.2 lbs. of Atrazine 90DF or Sim-Trol 90DF, or with 0.9 to 1.1 lbs. of Atrazine 90DF + 0.9 to 1.1 lbs. of Sim-Trol 90DF.

*Use Sim-Trol in preference to Atrazine when heavy infestations of Crabgrass or Fall panicum are expected.

**When using the tank mixture of this product + Atrazine 90DF + Sim-Trol 90DF, use equal rates of Atrazine and Sim-Trol as shown when heavy broadleaf weed infestations are expected. When heavy infestations of Crabgrass or Fall panicum are expected, use a 1:2 ratio of Atrazine + Sim-Trol, instead of the 1:1 ratio given. (Example: Total Atrazine 90DF + Sim-Trol 90DF = 1.8 lbs./A, use 0.6 lb. of Atrazine 90DF + 1.2 lbs. of Sim-Trol 90DF, respectively.) Refer to Comment No. 2 following Chart 1 for Atrazine 4L and Sim-Trol 4L conversions.

***For Cocklebur, Yellow nutsedge, and Velvetleaf control on *fine-textured soils* above 3% organic matter, apply 2.25 lbs./A of Atrazine 90DF, or equivalent rate of Atrazine 4L, or the same total amount of Atrazine + Sim-Trol, with 1.33 to 1.67 pts./A of this product.

7. TANK MIXTURE WITH ATRAZINE; OR ATRAZINE + 2,4-D; OR ATRAZINE + 2,4-D + DICAMBA (BANVEL) FOR MINIMUM-TILLAGE OR NO-TILLAGE SYSTEMS

In minimum-tillage or no-tillage systems where Field corn and Pop corn is planted directly into a cover crop, stale seedbed, established sod, or previous crop residues, this product applied in combination with Atrazine will kill most emerged small annual weeds. Apply this product + Atrazine before, during, or after planting, but before Field corn and Pop corn emerges, according to the rates in Table 1. Where heavy crop residues exist, add 0.8 to 1.6 pts./A of an appropriately labeled 3.8 lbs. a.i./gal. of 2,4-D amine (such as Weedar® 64, Weedar 64A, DMA-4 Herbicide, or Formula 40®) to the spray tank last and apply in a minimum of 25 gals. of carrier per acre.

As carriers, nitrogen solutions and complete liquid fertilizers, applied before Corn emergence, enhance burndown of existing weeds, and therefore, are recommended instead of water. Add X-77 surfactant at 1 to 2 qts./100 gals. of diluted spray, or another appropriate surfactant at its recommended rate, or add crop oil concentrate plus 28% liquid nitrogen (or equivalent). Apply before weeds exceed 3 inches in height. If Alfalfa is present, add Dicamba (Banvel) to the spray mixture at 0.33 to 0.5 pt./A and apply before Alfalfa exceeds 6 inches in height.

For fields with existing sod grasses (e.g., Bromegrass, Orchardgrass, Rye, or Timothy), when existing weeds exceed 3 inches in height or when very dry conditions exist, add Gramoxone Extra at the rate of 2.5 pts./A in place of or in addition to 2,4-D, as indicated above. Do not apply Gramoxone Extra in suspension-type liquid fertilizer. Observe all directions for use, use precautions, and limitations on the respective product labels when applying these products in tank mix combination.

8. TANK MIXTURE WITH MARKSMAN IN CONSERVATION TILLAGE — FIELD CORN

In conservation tillage systems where Field corn is planted directly into a cover crop or previous crop residue, this product + Marksman will kill most emerged small annual weeds. Apply this product + Marksman before, during, or after planting, but before Field corn emergence on *medium* and *fine soils* with greater than 2.5% organic matter. For fields with existing vegetation exceeding 3 inches in height or when very dry conditions exist, add Gramoxone Extra at its standard rate. This product + Marksman may be applied postemergence to Field corn less than 3 inches tall and before weedy grasses exceed the 2-leaf stage.

As carriers, nitrogen solutions and complete liquid fertilizers, applied before Corn emergence, enhance burndown of existing weeds. Do not apply Gramoxone Extra in suspension-type liquid fertilizer or use on emerged crop.

Refer to the Marksman label and follow all directions, limitations, use precautions, and information regarding application and use in Field Corn.

ACCEPTED
with COMMENTS
by EPA Letter Dated
OCT 7 2003

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

19713-526

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal. Open dumping is prohibited.

PESTICIDE STORAGE: This product may be stored at temperatures down to -30°F. For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product are toxic. Improper disposal of unused pesticide, spray mixture, or rinsate is a violation of Federal law. Pesticide, spray mixture, or rinsate that cannot be used according to label instructions must be disposed of according to Federal, state, or local procedures. For guidance in proper disposal methods, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office.

CONTAINER DISPOSAL: Do not reuse empty container. Triple rinse (or equivalent), puncture and dispose of in a sanitary landfill, or by incineration, or by open burning, if allowed by state and local authorities. Keep out of smoke from burning containers.

For Bulk and Mini Bulk Containers:

CONTAINER DISPOSAL: Reseal container and offer for reconditioning or triple rinse (or equivalent) and offer for recycling or reconditioning, or clean in accordance with manufacturer's instructions. Caintainer Precautions: Before refilling, inspect thoroughly for damage such as cracks, punctures, bulges, dents, abrasions and damaged or worn threads on closure devices.

REFILL ONLY WITH THIS PRODUCT. The contents of this container cannot be removed completely by cleaning. Refilling with material other than this product will result in contamination and may weaken the container.

THIS CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.

CONDITIONS OF SALE AND WARRANTY

9/11
Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

OUR RECOMMENDATIONS FOR USE of this product are based upon tests believed reliable. Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the Seller. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith.

In no case shall the Manufacturer or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product when such use and/or handling is not in strict accordance with directions given herewith. The foregoing is a condition of sale by the Seller and is accepted as such by the Buyer.

APPENDICES

APPENDIX A: COMPATIBILITY TEST

Because liquid fertilizers can vary, even within the same analysis, always check compatibility with herbicide(s) each time before use. Be especially careful when using complete suspension or fluid fertilizers, as serious compatibility problems are more likely to occur. Commercial application equipment may improve compatibility in some instances. The following test assumes a spray volume of 25 gals./A. For other spray volumes, make appropriate changes in the ingredients. Check compatibility using this procedure:

1. Add 1 pt. of fertilizer to each of 2 one-quart jars with tight lids.
2. To one of the jars, add one-quarter teaspoon or 1.2 milliliters of a compatibility agent approved for this use, such as Compex[®] or Unite[®] (one-quarter tsp. is equivalent to 2 pts./100 gals. spray). Shake or stir gently to mix.
3. To both jars, add the appropriate amount of herbicide(s). If more than one herbicide is used, add them separately with dry herbicides first, flowables next, and emulsifiable concentrates last. After each addition, shake or stir gently to thoroughly mix. The appropriate amount of herbicides for this test follows:
Dry herbicides: For each pound to be applied per acre, add 1.5 level teaspoons to each jar.
Liquid herbicides: For each pint to be applied per acre, add one-half teaspoon or 2.5 milliliters to each jar.
Note: For this product in tank mixtures with Atrazine + Sim-Trol, use one-third to one-half the amount of Atrazine specified above and the remainder as Sim-Trol, depending on whether the 1:2 or 1:1 ratio of Atrazine to Sim-Trol is to be applied.
4. After adding all ingredients, put lids on and tighten, and invert each jar 10 times to mix. Let the mixtures stand 15 minutes, and then look for separation, large flakes, precipitates, gels, heavy oily film on the jar, or other signs of incompatibility. Determine if the compatibility agent is needed in the spray mixture by comparing the 2 jars. If either mixture separates, but can be remixed readily, the mixture can be sprayed as long as good agitation is used. If the mixtures are incompatible, test the following methods of improving compatibility: (A) slurry the dry herbicide(s) in water before addition, or (B) add one-half of the compatibility agent to the fertilizer and the other one-half to the emulsifiable concentrate or flowable herbicide before addition to the mixture. If incompatibility is still observed, do not use the mixture.

APPENDIX B: LOW CARRIER APPLICATION

For Broadcast Ground Application Only

Use sprayers, such as Ag-Chem RoGator[®], Hagie, John Deere Hi-Cycle[™], Melroe Spra-Coupe, Tyler Patriot[™], or Willmar Air Ride[®], that provide accurate and uniform application. Only water may be used as a carrier. Screens in suction and in-line strainers should be 50-mesh. Manufacturers may require that tip screens as fine as 100-mesh be used with some nozzles. Use a pump with capacity to: (1) maintain up to 35 to 40 psi at the nozzles, and (2) provide sufficient agitation in tank to keep mixture in suspension. Use a minimum of 5 gals. of spray mixture per acre. Maximum recommended sprayer speed is 15 mph. Rinse sprayer thoroughly with clean water immediately after each use.

Note: Low pressure nozzles are recommended to reduce drift and increase application accuracy. Care should be taken when using automatic rate controlling devices to spray the material within the rated working pressure and flow ranges of the nozzles selected. Nozzle screens should be used when recommended by the manufacturer. All nozzles should be placed on 20-inch centers, except flooding types which should be placed on 40-inch centers. When Flat Fan-type nozzles are used, angles of 80° or 110° are recommended. Always read and follow the manufacturer's directions for optimum setup and performance of their nozzles or tips.

APPENDIX C: AERIAL APPLICATION

Apply this product in water alone or in tank mixtures with Atrazine, or Linuron in a minimum total volume of 2 gals./A by aircraft. This product may also be applied by air in combination with Prowl. Avoid application under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur. In order to assure that spray will be controllable within the target area when used according to label directions, make applications at a maximum height of 10 feet, using low-drift nozzles at a maximum pressure of 40 psi, and restrict application to periods when wind speed does not exceed 10 mph. To assure that spray will not adversely affect adjacent sensitive nontarget plants, apply this product alone or this product + Atrazine by aircraft at a minimum upwind distance of 400 feet from sensitive plants, or apply this product + Lorox at a minimum upwind distance of 300 feet from sensitive plants.

Avoid application to humans or animals. Flagmen and loaders should avoid inhalation of spray mist and prolonged contact with skin.

Aerial Drift Management

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment- and weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

10/11
The following drift management requirements must be followed to avoid off-target drift movements from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

1. The distance of the outermost nozzles on the boom must not exceed three-fourths the length of the wingspan or rotor.
2. Nozzles must always point backwards parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory Information section below.

Aerial Drift Reduction Advisory Management: Information on Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversions).

Controlling Droplet Size

- Volume - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- Pressure - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- Number of nozzles - Use the minimum number of nozzles that provide uniform coverage.
- Nozzle Orientation - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- Nozzle Type - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

Boom Length

For some use patterns, reducing the effective boom length to less than three-fourths the wingspan or rotor length may further reduce drift without reducing swath width.

Application Height

Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

Swath Adjustment

When applications are made with a crosswind, the swath will be displaced downward. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.)

Wind

Drift potential is lowest between speeds of 2 to 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential.

Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

Temperature and Humidity

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights when there is limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas

This pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive area).

APPENDIX D: CENTER PIVOT IRRIGATION APPLICATION

This product alone in tank mixture with other herbicides on this label, which are registered for center pivot application, may be applied in irrigation water preemergence (after planting, but before weeds or crop emerge) at rates recommended on this label. This product also may be applied postemergence to the Field corn crop or Pop corn crop and preemergence to weeds in Field corn or Pop corn where postemergence applications are allowed on this label. Follow all restrictions (height, timing, rate, etc.) to avoid illegal residues. Apply this product only through a center pivot irrigation system. Do not apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Extension specialists, equipment manufacturers, or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system, unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut down the system and make the necessary adjustments should the need arise.

Operating Instructions

1. The system must contain a functional check-valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check-valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump or piston pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.
8. Prepare a mixture with a minimum of 1 part of water to 1 part herbicide(s) and inject this mixture into the center pivot system. Injecting a larger volume of more dilute mixture per hour will usually provide more accurate calibration of metering equipment. Maintain sufficient agitation to keep the herbicide in suspension.
9. Meter into irrigation water during entire period of water application.
10. Apply in one-half to 1 inch of water. Use the lower water volume (one-half inch) on *coarse-textured soils* and the higher volume (1 inch) on *fine-textured soils*. More than 1 inch of water at application may reduce weed control by moving the herbicide below the effective zone in the soil.

Use Precautions: for center pivot applications: Where sprinkler distribution patterns do not overlap sufficiently, unacceptable weed control may result. Where sprinkler distribution patterns overlap excessively, crop injury may result.

APPENDIX E: DRY BULK GRANULAR FERTILIZERS

Many dry bulk granular fertilizers may be impregnated or coated with this product alone or selected tank mixtures of this product which are registered for preplant incorporated or preplant surface application which are used to control weeds in crops on this product label and are not prohibited from use on dry bulk granular fertilizers.

When applying this product or this product mixtures with dry bulk granular fertilizers, follow all directions for use and use precautions on the respective product labels regarding Field corn and Pop corn, rates per acre, soil texture, application methods (including timing of application), and rotational crops.

All individual state regulations relating to dry bulk granular fertilizer blending, registration, labeling, and application are the responsibility of the individual and/or company selling the herbicide/fertilizer mixture.

Prepare the herbicide/fertilizer mixtures by using any closed drum, belt, ribbon, or other commonly used dry bulk fertilizer blender. Nozzles used to spray this product and this product mixtures onto the fertilizer must be placed to provide uniform spray coverage. Care should be taken to aim the spray directly onto the fertilizer only and to avoid spraying the walls of the blender.

If the herbicide/fertilizer mixture is too wet, add a highly absorptive material, such as Agsorb® or Celatom MP-79®, or similar granular clay or diatomaceous earth materials, to obtain a dry, free-flowing mixture. Absorptive materials should be added only after the herbicide has been thoroughly blended into the fertilizer mixture. Best application results will be obtained by using a granule of six-thirtieth particle size or of a size similar to that of the fertilizer material being used. Generally, less than 2% by weight of absorptive material will be needed. Avoid using more than 5% absorptive material by weight. Calculate amounts of this product, Atrazine, Atrazine + Sim-Trol or Sim-Trol by the following formula:

$$\frac{2,000}{\text{lbs. of fertilizer per acre}} \times \frac{\text{pts./A of liquid or flowable product}}{\text{or flowable product}} = \text{pints of liquid or flowable product per ton of fertilizer}$$

$$\frac{2,000}{\text{lbs. of fertilizer per acre}} \times \frac{\text{lbs./A of dry product}}{\text{product}} = \text{pounds of dry product per ton of fertilizer}$$

Pneumatic (Compressed Air) Application (This Product Alone):

High humidity, high urea concentrations, low fertilizer use rates, and dusty fertilizer may cause fertilizer mixture to build up or plug the distributor head, air tubes, or nozzle deflector plates. To minimize buildup, premix this product with Exxon Aromatic 200 at a rate of 1 to 4 pts./gal. of this product. Aromatic 200 is a noncombustible/non-flammable petroleum product. Aromatic 200 may be used in either a fertilizer blender or through direct injection systems. Drying agents should not be used when using Aromatic 200.

Notes: (1) Mixtures of this product and Aromatic 200 must be used on dry fertilizer only. Poor results or crop injury may result if these mixtures are used in water or liquid fertilizer solutions for spraying applications. (2) When impregnating this product in a blender before application, a drier mixture can be attained by substituting a drying agent for Aromatic 200. The use of Agsorb FG or drying agents of six-thirtieth particle size are recommended. (3) Drying agents are not recommended for use with the On-The-Go impregnation equipment.

Use Precautions: To avoid potential for explosion, (1) Do not impregnate this product or this product mixtures on ammonium nitrate, potassium nitrate, or sodium nitrate, either alone or in blends with other fertilizers. (2) Do not use this product or this product mixtures on straight limestone, since absorption will not be achieved. Fertilizer blends containing limestone can be impregnated.

Application

Apply 200 to 700 lbs. of the herbicide/fertilizer mixture per acre. For best results, apply the mixture uniformly to the soil with properly calibrated equipment immediately after blending. Uniform application of the herbicide/fertilizer mixture is essential to prevent possible crop injury. Non-uniform application may also result in unsatisfactory weed control. In areas where conventional tillage is practiced, a shallow incorporation of the mixture into the soil may improve weed control. On *fine- or medium-textured soils* in areas where soil incorporation is not planned, i.e., reduced tillage situations or in some conventional till situations, make applications approximately 30 days before planting to allow moisture to move the herbicide/fertilizer mixture into the soil. On *coarse-textured soils*, make applications approximately 14 days prior to planting.

Use Precaution: To avoid crop injury, do not use the herbicide/fertilizer mixture on Field corn and Pop corn where bedding occurs.

Sim-Trol is a registered trademark of Sipcam Agro USA, Inc.

Beacon, Bicep, Concep, Evik, Exceed, Eradicane, Gramoxone, and Sutan, are registered trademarks of Syngenta Crop Protection, Inc.

Accent, and Lorox are registered trademarks of E.I. duPont de Nemours and Company, Inc.

Ag-Chem RoGator is a registered trademark of Ag-Chem Equipment Company

Agsorb is a registered trademark of Oil-Dri Corporation

Banvel, Banvel SGF, Basagran, Clarity, Marksman, and Prowl, are registered trademarks of Nufarm, Inc.

Celatom MP-79 is a trademark of Eagle-Picher Industries, Inc.

Complex is a registered trademark of KALO Agricultural Chemicals, Inc.

Hi-Cycle is a trademark of John Deere Company

Landmaster, and Roundup, are trademarks of Monsanto Company

Buctril, and Formula 40 are registered trademarks of Bayer AG

Tyler Patriot is a trademark of Tyler Ltd. Partnership

Unite is a registered trademark of HACO, Inc.

Willmar Air Ride is a registered trademark of Willmar Manufacturing

X-77 is a registered trademark of Loveland Industries, Inc.