



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

November 10, 2021

Luz G Chan
Registration Manager
DREXEL CHEMICAL COMPANY
P.O. Box 13327
Memphis, TN 38113-0327

Subject: Registration Review Label Amendments for ATRAZINE Incorporating Mitigation Measures from the Interim Decision and the Technical Registrants' Commitments for the Endangered Species Act (ESA) Biological Evaluation
Product Name: DREXEL AUGUZINE
EPA Registration Number: 19713-567
Application Date: 10/22/2020
Decision Number: 567141

Dear Luz G Chan:

The Agency, in accordance with the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the ATRAZINE Interim Decision and with the technical registrants' commitments for the ESA Biological Evaluation. The Agency has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved

Page 2 of 2
EPA Reg. No. 19713-567
Decision No. 567141

labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If you have any questions about this letter, please contact Ben Tweed at tweed.benjamin@epa.gov.

Sincerely,



Kelly Sherman
Chief, Risk Management and Implementation
Branch III (RMIB III)
Pesticide Re-Evaluation Division
Office of Pesticide Programs
U.S. Environmental Protection Agency

Enclosure

ACCEPTED
11/10/2021
 Under the Federal Insecticide, Fungicide
 and Rodenticide Act as amended, for the
 pesticide registered under
 EPA Reg. No. 19713-567



ATRAZINE GROUP 5 HERBICIDE

AuguzineTM
 St. Augustine Weed Killer

For the control of both emerged weeds and weeds from seeds in St. Augustinegrass and Centipedegrass, use 5 teaspoonfuls per 100 square feet.

ACTIVE INGREDIENTS:

Atrazine 4.00%
 Related compounds 0.08%

OTHER INGREDIENTS: 95.92%

TOTAL: 100.00%

**KEEP OUT OF REACH OF CHILDREN
 CAUTION**

[See FIRST AID Below]
 [See Side (Back) Panel for FIRST AID]; [See Page ___ for FIRST AID]
 [See Container Labeling for (FIRST AID and) Complete Directions for Use]
 [See (Attached) Booklet (Container Labeling) for Complete Directions for Use]

[SHAKE WELL BEFORE USING]
 [Recirculate Contents Before Use]

EPA Reg. No. 19713-567

EPA Est. No. 19713-XX-XXX

Net Content: ___(Gal.) (Fl. Oz.) (___L)

FIRST AID
<p>IF SWALLOWED:</p> <ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious or convulsing person.
<p>IF ON SKIN OR CLOTHING:</p> <ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also call CHEMTREC 800-424-9300 for emergency medical treatment information.</p>
<p>NOTE TO PHYSICIAN: There is no specific antidote for Atrazine. If this product is ingested, induce emesis or lavage stomach. The use of an aqueous slurry of activated charcoal may be considered.</p>

567SP-0621*P

PRECAUTIONARY STATEMENTS

Hazards To Humans And Domestic Animals

CAUTION: Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes, or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Wear long-sleeved shirt, long pants, chemical-resistant gloves made of any waterproof material including barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, natural rubber \geq 14 mils, polyethylene, polyvinyl chloride or viton \geq 14 mils, and shoes plus socks.

USER SAFETY RECOMMENDATIONS

Users should: 1) Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. 2) Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. 3) Remove clothing/PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Atrazine can travel (seep or leach) through soil and can enter groundwater which may be used as drinking water. Atrazine has been found in groundwater. Users are advised not to apply atrazine to Sand and Loamy sand soils where the water table (groundwater) is close to the surface and where these soils are very permeable, i.e., well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

This pesticide is toxic to aquatic invertebrates. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters or rinsate.

NON-TARGET ORGANISMS ADVISORY STATEMENT

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

ENDANGERED SPECIES

It is a Federal offense to use any pesticide in a manner that results in an unauthorized "take" (e.g., kill or otherwise harm) of an endangered species under the Endangered Species Act section 9. When using this product, you must follow the measures contained in the Endangered Species Protection Bulletin for the area in which you are applying for the product. You must obtain a Bulletin no earlier than six months before using this product. To obtain Bulletins consult <http://www.epa.gov/espp/>, call 1-877-447-3813, or email ESPP@epa.gov. You must use the Bulletin valid for the month in which you will apply the product.

NOT FOR USE ON TURF BEING GROWN FOR SALE OR OTHER COMMERCIAL USE AS SOD, OR FOR COMMERCIAL SEED PRODUCTION, OR FOR RESEARCH PURPOSES.

PRECAUTIONS AND RESTRICTIONS: Do not apply this product in a way that will contact any persons or pet, either directly or through drift. Keep people and pets out of the area during application.

Do not allow people or pets to enter the treated area until sprays have dried.

IMPORTANT: Read the entire Directions for Use, all precautions and Warranty – Conditions of Sale before using this product.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE, RESTRICTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY, ILLEGAL RESIDUES OR PERSONAL INJURY.

WEED RESISTANCE MANAGEMENT

ATRAZINE	GROUP	5	HERBICIDE
----------	-------	---	-----------

For resistance management, this product is a Group 5 mode of action herbicide. Any weed population may contain or develop plants naturally resistant to this product and other Group 5 mode of action herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance management strategies should be followed.

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of this product or other Group 5 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank-mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Drexel Chemical Company representatives at (901) 774-4370.

MANDATORY SPRAY DRIFT MANAGEMENT

Aerial Applications:

*Do not release spray at a height greater than 10 ft above the ground or vegetative canopy, unless a greater application height is necessary for pilot safety.

*Applicators are required to use a coarse or coarser droplet size (ASABE S572).

*If the windspeed is 10 miles per hour or less, applicators must use ½ swath displacement upwind at the downwind edge of the field. When the windspeed is between 11 – 15 miles per hour, applicators must use ¾ swath displacement upwind at the downwind edge of the field.

*Do not apply when windspeeds exceed 15 mph at the application site. If the windspeed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% of less of the rotor diameter for helicopters.

*Do not apply during temperature inversions.

*User must maintain a 150 foot (46 meter) in-field downwind buffer (in the direction in which the wind is blowing) from the edge of streams and rivers, as well as high-tide line for all estuarine/marine environments.

Ground Boom Applications:

*User must only apply with the release height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy.

*Applicators are required to use a coarse or coarser droplet size (ASABE S572).

*Do not apply when wind speeds exceed 10 miles per hour at the application site.

*User must maintain a 15 foot (4.6 meter) in-field downwind buffer (in the direction in which the wind is blowing) from the edge of streams and rivers, as well as high-tide line for all estuarine/marine environments.

*Do not apply during temperature inversions.

Boomless Ground Applications:

*Applicators are required to use a coarse or coarser droplet size (ASABE S572).

*Do not apply when wind speeds exceed 10 miles per hour at the application site.

*User must maintain a 15 foot (4.6 meter) in-field downwind buffer (in the direction in which the wind is blowing) from the edge of streams and rivers, as well as high-tide line for all estuarine/marine environments.
*Do not apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.

BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size - Ground Boom

- Volume- Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure- Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle- Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

BOOM HEIGHT – GROUND BOOM

For ground equipment, the boom should remain level with the crop and have minimal bounce.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

BOOMLESS GROUND APPLICATIONS

Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

HANDHELD TECHNOLOGY APPLICATIONS

Take precautions to minimize spray drift.

USE SITE RESTRICTION

Not for use in the states of Alaska or Hawaii or in the U.S. Territories (American Samoa, Guam, North Mariana Islands, Puerto Rico, U.S. Virgin Islands).

APPLICATION PROCEDURES

- Do not use near adjacent desirable plants or in greenhouse, or injury may occur.
- To avoid spray drift, apply with a coarse low pressure spray.
- Do not apply under windy conditions.
- Avoid spray overlap.

- This product is to be mixed with water only. Combinations with other chemicals in the application mixture are not recommended.
- Do not apply when temperatures exceed 90°F.
- Use only on healthy Turfgrass which is not under heat or drought stress and reasonably free of infestations of insects, nematodes, and diseases.
- Applications during the summer should be limited to spot treatment only.
- Do not use on muck or alkaline soils.
- Do not use over rooting area of trees, ornamentals, vegetables or other desirable plants other than the indicated Turfgrasses.
- Do not apply to areas where the natural drainage would carry the atrazine to adjacent desirable plants.
- Do not replant treated areas (except with listed Turfgrass) within 12 months after treatment or injury may result.
- Do not overseed turf areas within 6 months after treatment.
- Sprigging of listed Turfgrass is allowed in treated areas, however, some yellowing or stunting may occur.

APPLICATION RESTRICTIONS

- Maximum rate per application on Turfgrass (including lawns) is 8.6 fluid ounces per 1,000 square feet (23.5 pt./A) (1 lb. a.i./A/application).
- Maximum of two applications per year is permitted.
- Do not use backpack spray application to landscape Turf except for spot treatments only.

USE DIRECTIONS

Weeds Controlled or Suppressed:

Annual Bluegrass (*Poa annua*), Chickweed (Common and Mouseear), Crabgrass (suppression only), Cransbill, Cudweed, Dichondra, Florida Betony, Henbit, Knotweed, Lespedeza, Moneywort, Mustards, Narrowleaf Vetch, Parsley-Piert, Pennywort (Dollarweed), Sandspur, Smutgrass, Spurge, Spurweed, Swinecrest, Woodsorrel and various annual Clovers.

Established St. Augustinegrass and Centipedegrass:

This product may be applied during both dormant and growing seasons. Best results are usually obtained when applied in early Spring or dormant periods when weeds are small or have not emerged. Do not apply more than two treatments per year.

This product controls Spurweed, Florida Betony, Annual Bluegrass and many other problem weeds in St. Augustinegrass and Centipedegrass turfs.

This product will control both emerged weeds and weeds from seeds. Rain or water within 2 or 3 days of application may decrease the effectiveness on emerged weeds. However, for the control of weeds from seeds, rainfall or watering is necessary within 7 to 10 days after treatment.

Spurweed:

Control of Spurweed can best be obtained by applying when Spurweed has newly emerged (December and January.)

Florida Betony:

This weed emerges in the Fall. Apply this product in mid to late October followed by a second application in mid to late February.

Dichondra, Moneywort:

Apply this product in early April followed by a second application in July. **Note:** Applications for Spurweed or Florida Betony generally will give control or suppression of the other listed weeds. However, as a general rule, this product will give best control when applied to young tender weeds or just prior to weed emergence.

RATE OF APPLICATION* FOR ST. AUGUSTINEGRASS AND CENTIPEDEGRASS		
Areas To Be Treated	Amount of This Product	Minimum Amount of Water
100 sq. ft.	5 tsps.	1 qt.
500 sq. ft.	4.3 fl. oz.	1 gal.
1,000 sq. ft.	8.6 fl. oz.	2 gal.
2,000 sq. ft.	17.6 fl. oz. (1.1 pt.)	4 gal.
43,560 sq. ft. (1 acre)	376.5 fl. oz. (23.5 pt.)	87 gal.
*Determine total area to be sprayed and base rate of application on the above rates. Do not exceed the maximum application rate of 376.5 fl. oz. (23.5 pt.) of this product (1 lb. a.i.) per acre per application on Residential Turf. Maximum number of application per year is two.		

METHODS OF APPLICATION

Conventional Sprayers:

Application should be made with conventional compressed air garden sprayer equipped with a flat fan spray tip. Correct calibration of the sprayer is essential. Overdosing will cause severe plant injury and or death of the desired Turf. Insufficient application may not give anticipated weed control.

Measure the amount of this product into the spray tank and dilute with water according to the above Table. Close tank sprayer and shake well before applying.

Use of backpack sprayers to landscape Turf is restricted to spot treatments only.

Hose-End Sprayers With Ready-Spray Nozzle:

One quart (32 fl. oz.) of this product will treat 3,720 square feet. of listed grasses.

DO NOT apply one quart of this product to an area less than 3,720 square feet. DO NOT apply this product to the same area more than twice per year.

Using the Ready-Spray Nozzle, apply this product following the below procedures:

1. Shake container well before using.
2. Connect a garden hose to the Ready-Spray nozzle. Make sure that the dial nozzle is in the "OFF" position with the [green] safety tab in the valve notch.
3. Turn on water at faucet. Extend hose to the farthest area to be treated and work back toward the faucet so you don't come in contact with the treated area.
4. To begin spraying, point nozzle toward treatment side and:
 - a) Bend back the [green] safety tab [located at the right of the yellow dial] with your thumb.
 - b) Hold while turning the yellow dial clockwise quickly until it stops. Water will automatically mix with this product.
5. Spray until wet to control the weeds listed on this label. Walk at a steady pace while spraying using an even sweeping motion.
6. To STOP spraying, quickly turn the dial counter clockwise until it stops and the safety tab engages into the notch on the valve. Turn water OFF at faucet. To relieve pressure before removing from nozzle from the hose, bend the safety tab back and turn dial "ON" until water stops spraying.
7. To store unused product, make sure the dial is in the "OFF" position with the [green] safety tab in the valve notch. Place in cool area, away from heat, sunlight or open flame.

Read the 'APPLICATION PROCEDURES' section of this label for additional application instructions.

For Connect and Spray Applicator:

1. Make sure knob control lever is in the OFF position. Shake well and attach to hose.

([Optional] Make sure the handle is in the upright position and locked OFF position. Shake container well and attach spray nozzle to hose.)

2. Turn water on at faucet, extend hose to the farthest area to be treated and work back toward the faucet so you don't come in contact with the treated area. Aim nozzle toward lawn and turn control lever to ON to begin spraying. The product will automatically siphon and mix with water. Walk at a steady pace while spraying, using an even sweeping motion. Spray weeds until wet.

([Optional] Turn water on at faucet, aim nozzle towards lawn, bend tab back, and rotate handle clockwise towards the hose connection in the ON position to begin spraying. The product will automatically siphon and mix with water.) [Optional Icon]

3. When finished, turn control lever to OFF position. Turn faucet off. Discharge residual water pressure by turning control lever ON and OFF again.

([Optional] When finished, turn the handle counterclockwise until the handle is in the upright position and locked position. Turn faucet off. Discharge residual water pressure by turning the handle ON and OFF again. Unscrew spray nozzle from hose.)

[ALTERNATE TEXT]

[Hose-End Ready to Spray System]

One quart (32 fl. oz.) of this product will treat 3,720 square feet of listed grasses.

DO NOT apply one quart of this product to an area less than 3,720 square feet. DO NOT apply this product to the same area more than twice per year.

How to Use the Hose-End Ready-To-Spray System

Connect

1. Shake well before using.
2. Connect sprayer to hose.
3. Turn on water.

Spray

1. To begin spraying, point nozzle in the direction you want to spray.
2. [Bend small plastic tab back and] Turn knob [clockwise] to ON position.
3. Spray evenly to the area to be treated. [Refer to the [sight gauge] clear view strip (graduated scale) on the side of the container to determine the amount of product sprayed] [Optional Marketing Statement]

Finish

1. To stop spraying, [turn knob counter clockwise] to OFF position.
2. Turn off water.
3. Relieve water pressure slowly by [bending plastic tab back and] turning knob to ON position until water slows to a drip. Then turn knob back to OFF position.
4. Disconnect sprayer from hose.]

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store pesticide in the original container in a locked storage area.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL:

Nonrefillable Container (rigid material; ≤ 5 gallons): Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container one-fourth full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Nonrefillable Container (rigid material; > 5 gallons up to < 250 gallons): Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after

emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container one-fourth full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Refillable Container (≥ 250 Gals. & Bulk): Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in sanitary landfill, or by incineration or by other procedures allowed by state and local authorities.

For Residential/Household Use:

PESTICIDE DISPOSAL: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. **If empty**, place in trash or offer for recycling if available. **If partially filled**, call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

WARRANTY—CONDITIONS OF SALE

OUR DIRECTIONS FOR USE of this product are based upon tests believed reliable. Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically directed and other influencing factors in the use of this product are beyond the control of the Seller. To the extent consistent with applicable law, Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith. To the extent consistent with applicable law, in no case shall the Manufacturer or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product when such use and/or handling is not in strict accordance with directions given herewith. The foregoing is a condition of sale by the Seller and is accepted as such by the Buyer.

Manufactured By:



Drexel Chemical Company

P.O. BOX 13327, MEMPHIS, TN 38113-0327

SINCE 1972

AUGUZINE and the Drexel logo are either trademarks or registered trademarks of Drexel Chemical Company.

Optional Marketing Statements:

- * For St. Augustine Grass
- * Auguzine will control (followed by any or all of this list) Annual Bluegrass (*Poa annua*), Chickweed (Common and Mouseear), Cudweed, Dichondra, Florida Betony, Henbit, Knotweed, Lespedeza, Moneywort, Pennywort (Dollarweed), Mustards, Narrowleaf Vetch, Parsley-Piert, Smutgrass, Spurge, Spurweed, Swinecrest, Woodsorrel and various annual clovers.
- * Auguzine supresses Crabgrass
- * Controls Listed Tough Weeds
- * Controls both listed emerged weeds and weeds from seeds
- * One pint covers 1,852 sq. ft.
- * One Quart covers 3,720 sq. ft.
- * 0.5 Gallon covers 7,408.2 sq. ft.
- * One Gallon covers 14,816.3 sq. ft.
- * Kills Listed Tough Weeds
- * Kills Other Listed Weeds
- * Kills & Prevents Listed Tough Weeds
- * Kills Weeds...Not the Lawn
- * With Atrazine
- * For outdoor use around the home only
- * [One quart] Treats 3,720 sq. ft.
- * Covers 3,720 sq. ft.
- * Kills listed tough weeds [including Dollarweed and Sandspur] [and other listed weeds]
- * Contains atrazine
- * Attach to hose and spray
- * Ready to spray [apply]
- * For St. Augustinegrass and Centipedegrass [Lawns]
- * Kills Dollarweed, Sandspur and other listed weeds
- * [Front panel starburst: Connect and Spray Applicator]
- * [Front panel: Connect & Spray Applicator]
- * Product Fact Box

PRODUCT FACT BOX	
What it Does:	
Kills established listed weeds including Dollarweed, Sandspur and Florida Betony and prevents weeds from emerging in the Lawn	
Where to Use:	
(icon outdoors) St. Augustinegrass and Centipedegrass Lawns: Kills the weeds, not the Lawn	
When to Use:	
Anytime temperatures are below 90°F	
Questions & Comments:	
(phone icon) (computer icon) Call (901) 774-4370 or visit our website at www.drexchem.com	

* Product Usage Box

Product Usage	
Kills & Prevents	Annual Bluegrass, Dollarweed, Florida Betony, Sandspur, Spurge, Spurweed, Woodsorrel, various annual clovers, and other listed weeds
Use On	Established St. Augustinegrass and Centipedegrass
Amount to Use	<ul style="list-style-type: none"> • 4.3 fl. ozs. mixed with 1 gallon of water treats 500 sq. ft. (<i>[Optional] One quart (32 fl. ozs.) of this product will treat 3,720 sq. ft.</i>) • 32 fl. ozs. mixed with 7.4 gallons of water treats 3,720 sq. ft. <i>Wait 72 hours before watering treated, emerged weeds.</i>
When to Use	When air is calm and temperature below 90°F – up to 2 applications per year.