



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

July 31, 2020

Luz G. Chan
Registration Manager
Drexel Chemical Company
P.O. Box 13327
Memphis, TN 38113-0327

Subject: Label Amendment – Incorporation of Supplemental Labels
Product Name: DREXEL DE-AMINE 4
EPA Registration Number: 19713-650
Application Date: January 28, 2019
Decision Number: 548022

Dear Ms. Chan:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance

Page 2 of 2
EPA Reg. No. 19713-650
Decision No. 548022

with FIFRA section 6. If you have any questions, please contact Kable Bo Davis by phone at 703-306-0415, or via email at davis.kable@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read 'M. Ondish', followed by the word 'for' written in a similar cursive style.

Mindy Ondish
Product Manager 23
Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

Enclosure

ACCEPTED
 07/31/2020
 Under the Federal Insecticide, Fungicide
 and Rodenticide Act as amended, for the
 pesticide registered under
 EPA Reg. No. 19713-650

2,4-D	GROUP	4	HERBICIDE
-------	-------	----------	-----------



De-Amine® 4

Contains Dimethylamine Salt of 2,4-D.
 For selective control of many broadleaf weeds in various sites listed on the label.

ACTIVE INGREDIENT:		
2,4-Dichlorophenoxyacetic acid, dimethylamine salt*	47.5%
OTHER INGREDIENTS:	52.5%
TOTAL:	100.0%

*2,4-Dichlorophenoxyacetic acid equivalent (a.e.) – 39.5%; 3.8 lbs. per gallon. Isomer specific by AOAC Method No. 978.05.

KEEP OUT OF REACH OF CHILDREN DANGER / PELIGRO

Si usted no entiende a la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

[See First Aid Below]

[See Side (Back) Panel for FIRST AID]; [See Page ___ for FIRST AID]
 [See Container Labeling for (FIRST AID and) Complete Directions for Use]
 [See (Attached) Booklet (Container Labeling) for Complete Directions for Use]

EPA Reg. No. 19713-650
 EPA Est. No. 19713-XX-XXX

Net Content: _____ Gals. (L)

FIRST AID
<p>IF IN EYES:</p> <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
<p>IF SWALLOWED:</p> <ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
<p>IF ON SKIN OR CLOTHING:</p> <ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
<p>IF INHALED:</p> <ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also call CHEMTREC at 800-424-9300 for emergency medical treatment information.</p>
<p>Note to Physician: This product is a phenoxy type of herbicide. There is no specific antidote. Base all treatments on observed signs and symptoms of distress in the patient. Probable mucosal damage may contraindicate the use of gastric lavage.</p>

650SP-0720*P

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

DANGER: Corrosive. Causes irreversible eye damage. Harmful if swallowed. Do not get in eyes or on clothing. Avoid breathing vapor or spray mist.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

All mixers, loaders, flaggers, other applicators and handlers must wear: Long-sleeved shirt and long pants, shoes plus socks, chemical-resistant gloves such as barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils or viton \geq 14 mils, protective eyewear (goggles, safety glasses or face shield), and chemical-resistant apron when mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.

See Engineering Controls for additional requirements.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. After each day of use, clothing or PPE must not be reused until it has been cleaned.

ENGINEERING CONTROLS

When handlers use enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Pilots must use an enclosed cockpit that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.240(d)(6)].

USER SAFETY RECOMMENDATIONS

Users should: 1) Wash hands thoroughly before eating, drinking, chewing gum, using tobacco, or using the toilet. 2) Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water. 3) Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. For terrestrial uses, do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas and nontarget plants. Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

Aquatic Weed Control: Fish breathe dissolved oxygen in the water and decaying weeds also use oxygen. When treating continuous, dense weed masses, it may be appropriate to treat only part of the infestation at a time. For example, apply the product in lanes separated by untreated strips that can be treated after vegetation in treated lanes has disintegrated. During the growing season, weeds decompose in a 2 to 3 week period following treatment. Begin treatment along the shore and proceed outwards in bands to allow fish to move into untreated areas. Waters having limited and less dense weed infestations may not require partial treatments. Do not contaminate water used for irrigation or domestic purposes (except as directed on this label) especially in areas where Cotton, Grapes, Tomatoes or other susceptible plants are grown.

Do not treat irrigation ditches in areas where water will be used to overhead (sprinkler) irrigate susceptible crops especially Cotton, Grapes, Tobacco and Tomatoes.

Do not apply this product directly to, or permit to drift onto Cotton, Flowers, Fruit trees, Grapes, Okra, Tomatoes, Vegetables or other desirable plants which are susceptible to 2,4-D. Do not apply near susceptible plants since very small quantities of 2,4-D will cause severe injury during the growing or dormant periods. Crops contacted by sprays or spray drift of this product may be killed or suffer significant stand loss with extensive quality and yield reduction.

Mixing and Loading: Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Exercise precautions when handling 2,4-D pesticides at such sites to prevent contamination of ground water supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

WEED RESISTANCE MANAGEMENT

2,4-D	GROUP	4	HERBICIDE
-------	-------	---	-----------

For resistance management, this product is a Group 4 mode of action herbicide. Any weed population may contain or develop plants naturally resistant to this product and other Group 4 mode of action herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance management strategies should be followed.

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of this product or other Group 4 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank-mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Drexel Chemical Company representatives at (901) 774-4370.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all directions for use carefully before applying.

Do not apply this product in a way that will contact workers or other person, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard (WPS), 40 CFR Part 170. This standard contains the requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers or agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered in the WPS.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours. PPE required for early entry to treated areas that is permitted under the WPS and that involves contact with anything that has been treated such as plants, soil, or water is: Coveralls, shoes plus socks, chemical-resistant gloves made of any waterproof material, and protective eyewear.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Entry Restrictions for Non-WPS Uses: When this product is applied in the rangeland and established pastures not harvested for hay or seed; non-cropland areas, when tree injection method only in forest sites, and when applied in aquatic areas, do not enter or allow people (other than applicator) or pets on treatment area during application. Do not enter treated areas until sprays have dried.

USE INFORMATION

DE-AMINE 4 is a herbicide that contains 2,4-D Dimethylamine salt active ingredient. It is intended for selective control of many broadleaf weeds in the various sites listed on this label.

Apply this product as a water or oil-water spray during warm weather when target weeds or woody plants are actively growing. Application under drought conditions will often give poor results. Use low spray pressure to minimize drift. Generally, the lower dosages specified on this label will be satisfactory for young, succulent growth of susceptible weed species. For less susceptible species and under conditions where control is more difficult, use the higher specified rates. Deep-rooted perennials such as Canada thistle and Field bindweed and many woody plants usually require repeated applications for satisfactory control. Consult your State Agricultural Experiment stations or Extension Service Weed Specialist for recommendations from this label that best fit local conditions.

USE PRECAUTIONS

- Excessive amounts of 2,4-D in the soil may inhibit seed germination and plant growth temporarily.

USE RESTRICTIONS

- DO NOT apply this product through any type of irrigation system.
- Use of this product in certain portions of California, Oregon, and Washington is subject to the January 22, 2004 Order for injunctive relief in *Washington Toxics Coalition et. al. v. EPA*, C01-0132C, (W.D. W.A.). For further information, please refer to EPA website: <http://www.epa.gov/espp/litstatus/wtc/index.htm>.

SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of application (e.g., ground, aerial, airblast) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Droplet Size

When applying sprays that contain 2,4-D as the sole active ingredient, or when applying sprays that contain 2,4-D mixed with active ingredients that require a coarse or coarser spray, apply only as a coarse or coarser spray (ASAE standard 572) or a volume mean diameter of 385 microns or greater for spinning atomizer nozzles.

When applying sprays that contain 2,4-D mixed with other active ingredients that require a medium or more fine spray, apply only as a Medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Wind Speed

Do not apply at wind speeds greater than 15 mph. Only apply this product if the wind direction favors on target deposition and there are not sensitive areas (including, but not limited to, residential areas, bodies of water, known habitat for non-target species, non-target crops) within 250 feet downwind. If applying a medium spray, leave one swath unsprayed at the downwind edge of the treated field.

Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if: a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to Cotton, Okra, flowers, fruit trees, Grapes (in growing stage), fruit trees (foliage), Soybeans (vegetative stage), Ornamentals, Sunflowers, Tomatoes, Beans, and other vegetables or Tobacco. Small amounts of spray drift that may not be visible may injure susceptible broadleaf plants.

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

Equipment

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

Aerial Application

The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter. Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety. This requirement does not apply to forestry or Rights-of-way applications. When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this by adjusting the path of the aircraft upwind.

Ground Boom Application

Groundboom: Do not apply with a nozzle height greater than 4 feet above the crop canopy.

Handguns and Boomless Nozzles: Applications for Rights-of-way vegetation management are best applied with specialized nozzles delivering a coarse or very coarse spray volume.

MIXING

Mix this product only with water, unless otherwise directed on this label. Add about half the water to the mixing tank, then, add this product with agitation, and finally the rest of the water with continuing agitation.

Note: Adding oil, wetting agent, or other surfactant to the spray mixture may increase effectiveness on weeds, but also may reduce selectivity to crops resulting in crop damage.

Tank-mixing

When tank-mixing, read and follow the label of each tank-mix product used for precautionary statements, directions for use, weeds controlled, and geographic and other restrictions. Use in accordance with the most restrictive of label limitations and precautions. No label dosages should be exceeded. Do not tank-mix this product with any product containing a label prohibition against tank-mixing with 2,4-D.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank-mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank-mixture.

Tank-mix Compatibility Testing

If compatibility with the tank-mix product is not known, perform a jar test prior to tank-mixing to ensure compatibility of this product and other pesticides. Use a clear glass quart jar with lid and mix the tank-mix ingredients in their relative proportions. Invert the jar containing the mixture several times and observe the mixture for approximately one-half hour. If the mixture balls-up, jells, forms flakes, sludge, oily films or layers, or other precipitates, it is not compatible and the tank-mix combination should not be used.

Mixing With Liquid Fertilizer

This product may be combined with liquid nitrogen fertilizer suitable for foliar application to accomplish broadleaf weed control and fertilization of Corn, small grains or Pastures in a single operation. Use this product in accordance with directions for these crops provided in this label. Use liquid fertilizer at rates specified by the supplier or Extension Service Specialist.

Test for mixing compatibility as describe above before mixing in spray tank. A compatibility aid such as Mix™, Unite or Compex may be needed in some situations. Compatibility is best with liquid fertilizer solutions containing only nitrogen. Mixing with N-P-K solutions may not be satisfactory, even with the addition of a

compatibility aid. Premixing 1 part this product with up to 4 parts water may help in situations when mixing difficulty occurs. Fill the tank about half full with the liquid fertilizer, then add the required amount of this product with agitation. Maintain agitation and complete filling the tank with liquid fertilizer. Apply immediately and continue agitation in spray tank during application. Do not store the spray mixture. Application during very cold weather (near freezing) is not advisable.

SPRAYER CLEAN-OUT

To avoid injury to desirable plants, equipment used to apply this product should be thoroughly cleaned before re-use or applying other chemicals.

1. Rinse and flush application equipment thoroughly after use at least three times with water. Dispose of all rinse water by application to treatment area or apply to non-cropland area away from water supplies.
2. During the second rinse, add 1 quart of household ammonia for every 25 gallons of water. Circulate the solution through the entire system so that all internal surfaces are contacted (15 to 20 minutes). Let the solution stand for several hours, preferably overnight.
3. Flush the solution out of the spray tank through the boom.
4. Rinse the system twice with clean water, recirculating and draining each time.
5. Remove nozzles and screens and clean separately.
6. If equipment is to be used to apply another pesticide or agricultural chemical to a 2,4-D susceptible crop, additional steps may be required to remove all traces of 2,4-D, including cleaning of disassembled parts and replacement of hoses or other fittings that may contain absorbed 2,4-D.

APPLICATION

Apply with calibrated air or ground equipment using sufficient spray volume to provide adequate coverage of target weeds or as otherwise directed in specific use directions. For broadcast application, use a spray volume of 1 or more gallons per acre by air and 5 or more gallons per acre for ground equipment unless otherwise specified. Where states have regulations which specify minimum spray volumes, they should be observed. In general, spray volume should be increased as crop canopy, height and weed density increase in order to obtain adequate spray coverage.

RANGE OF RATES AND APPLICATION TIMING

Generally, the lower dosages given will be satisfactory for young, succulent growth of sensitive weed species. For less sensitive species and under conditions where control is more difficult, the higher dosages will be needed. Apply this product during warm weather when weeds are young and actively growing.

SPOT TREATMENT

To prevent misapplication, spot treatments should be applied with a calibrated boom or with hand sprayers using a fixed spray volume per 1,000 square feet as indicated below.

Handheld Sprayers: Handheld sprayers may be used for spot applications of this product. Care should be taken to apply the spray uniformly and at a rate equivalent to a broadcast application. Application rates in the table are based on the application rate for an area of 1,000 square feet. Mix the amount of this product corresponding to the desired broadcast rate in 1 to 3 gallons of spray. To calculate the amount of this product required for larger areas, multiply the table value (fl. oz. or ml) by the thousands of square feet to be treated. An area of 1000 square feet is approximately 10.5 X 10.5 yards (strides) in size.

Rate Conversion Table for Spot Treatment

Broadcast Rate (Pint/Acre)							
0.5	0.66	0.75	1	2	3	4	8
Equivalent Amount of This Product per 1,000 sq. ft. (fl. oz.)							
1/5 (5.5 mL)	1/4 (7.3 mL)	1/3 (8.3 mL)	3/8 (11 mL)	3/4 (22 mL)	1 (33 mL)	1.5 (44 mL)	3 (88 mL)

Conversion factor: 1 pt. = 16 fl. ozs.; 1 fl. oz. = 29.6 (30 mL)

BAND APPLICATION

This product may be applied as a band treatment. Use the formula below to determine the appropriate rate and volume per treated acre.

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast rate per acre} = \text{Band rate per treated acre}$$

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast volume per acre} = \text{Band volume per treated acre}$$

WEEDS CONTROLLED

ANNUALS OR BIENNIALS		
Beggarticks*	Kochia	Ragweed, common
Bittercress, small-flowered	Lambsquarters, common	Ragweed, giant
Bitterweed	Lettuce, prickly*	Rape, wild
Broomweed, common*	Lettuce, wild	Rocket, yellow
Burdock, common	Lupines	Salsify, common*
Buttercup, small-flowered*	Mallow, little*	Salsify, western*
Carpetweed	Mallow, Venice*	Shepherdspurse
Cinquefoil, common	Marshelder	Sicklepod
Cinquefoil, rough	Morningglory, annual	Smartweed (annual species)*
Cocklebur, common	Morningglory, common	Sneezeweed, bitter
Coffeeweed	Morningglory, ivy	Sowthistle, annual
Copperleaf, Virginia	Morningglory, woolly	Sowthistle, spiny
Croton, Texas	Mousetail	Spanishneedles
Croton, woolly	Mustards (except blue mustard)	Sunflower
Fleabane, rough	Parsnip, wild	Sweetclover
Flixweed	Pennycress, field	Tansymustard
Galinsoga	Pepperweed*	Thistle, bull
Geranium, Carolina	Pigweed (<i>Amaranthus</i> spp.)*	Thistle, musk*
Hemp, wild	Poorjoe	Thistle, Russian (tumbleweed)*
Horseweed, (marestail)	Primrose, common	Velvetleaf
Jewelweed	Purslane, common	Vetch
Jimsonweed	Pusley, Florida	
Knotweed*	Radish, wild	

*These weeds are only partially controlled and may require repeated applications and/or use of higher specified rates of this product even under ideal conditions of application.

PERENNIALS		
Alfalfa*	Coffeeweed	Nettles (including stinging)*
Artichoke, Jerusalem*	Cress, hoary*	Onion, wild*
Aster, many-flower*	Dandelion	Pennywort
Austrian fieldcress*	Docks*	Plantains
Bindweed (hedge, field, European)*	Dogbanes*	Ragwort, tansy*
Blue lettuce	Evening primrose, cutleaf	Sowthistle, perennial
Blueweed, Texas	Garlic, wild*	Speedwell
Broomweed	Goldenrod	Spotted catsear
Bullnettle *	Hawkweed, orange*	Thistle, Canada*
Carrot, wild*	Healal	Vervains*
Catnip	Ironweed, western	Waterplantain
Chicory	Ivy, ground*	Wormwood
Clover, red*	Loco, bigbend	

*These weeds are only partially controlled and may require repeated applications and/or use of higher specified rates of this product even under ideal conditions of application.

OTHERS			
Alder	Devil's claw (<i>Proboscidea louisianica</i>)	Orange hawkweed*	Sunflower
Alligator weed	Duckweed	Parrot feather	Tanweed
American lotus	Elderberry	Poison hemlock	Tarweed
Biden	Frenchweed	Poison ivy	Toadflax
Bittersweet	Goatsbeard	Pokeweed	Tumbleweed
Bitter wintercress	Goosefoot	Povertyweed	Virginia creeper
Blackeyed Susan	Gumweed	Puncture vine	Water hyacinth
Blessed thistle	Henbit	Purslane	Water lily
Boxelder	Hoary cress*	Rush	Water primrose
Buckhorn	Horsetail	Sagebrush	Water shield
Bulrush	Honeysuckle	Saltcedar*	Wild strawberry
Bur ragweed	Indigo	Sheperdspurse	Wild sweet potato
Chickweed	Indian mallow	Southern wild rose	Willow
Cockle	Locoweed	Spatterdock	Witchweed
Common waterplantain	Marijuana	St. Johnswort	Wormseed
Creeping Jenny	Mexican weed	Stinkweed	Yellow rocket
Curly indigo	Nut sedge	Sumac	

*May require repeated application and/or use of the higher specified label rate of this product even under ideal conditions. In California, DO NOT use this product to control Saltcedar.

USE SITES

ASPARAGUS

Time of Application	This Product (Pt./Ac.)	Specific Use Instructions
Spring	3 to 4	Apply this product in the Spring on actively growing weeds in 50 to 60 gallons of water per acre by ground or in 12 gallons of water per acre by air. Refer to the "WEEDS CONTROLLED" section of this label for list of weeds. If spears are present, apply this product immediately after cutting. Spears contacted by the spray may be malformed and off-flavored. If spears are malformed by spray, cut immediately and discard. Only apply as post-harvest spray using drop nozzles to avoid spraying the fern.

Use Restrictions:

- Do not apply more than 4.2 pints of this product (2 lbs. a.e.) per acre per application.
- Do not make more than 2 applications per crop cycle.
- Allow at least 30 days between applications.
- Do not harvest within 30 days of application.
- When tank-mixing with products that contain 2,4-D, do not exceed a combined total of 4 pounds a.e. per acre per year.

BLUEBERRIES*

Time of Application	This Product (Pt./Ac.)	Specific Use Instructions
First Application: Spring If necessary, second application: After Harvest	2.9	High Bush Berries (Vegetative Strips between Rows) Broadleaf weeds (Broadleaf dock, Canada thistle, Dandelion, Spotted catsear): Apply 2.9 pts. of this product in 50 gallons of water per acre using ground equipment only. Make the first application in the Spring as a direct, shielded spray to the vegetative strip between Blueberry rows, avoiding contact with the Blueberry plant foliage. If necessary, a second application at 3 pts. of this product in 50 gallons of water per acre may be made after harvest to control regrowth of broadleaf weeds.
*High Bushberries (Vegetative strips between rows); For use in the states of Massachusetts, New Jersey, Oregon, Washington and Wisconsin only.		

Use Precautions:

- INJURY TO CROPS FROM THIS HERBICIDE MAY OCCUR. IF YOU ARE NOT PREPARED TO ACCEPT SOME DEGREE OF CROP INJURY DO NOT USE THIS PRODUCT ON BLUEBERRIES.
- For optimum herbicide performance, mowing between rows for at least 7 days before or after the application is not advised.
- To avoid injury to Blueberry plants, apply when temperature is less than 65°.
- Avoid drip or contact of this product to Blueberry plants in the growing or dormant period as it may kill or cause significant plant injury that could result in grade or yield loss.
- Soil residue of this product may temporarily inhibit seed germination and plant growth.

Use Restrictions:

- The pre-harvest interval (PHI) is 30 days.
- Limited to 2 application per year.
- Do not exceed 2.9 pints (1.4 lbs. a.e.) per acre per application.
- Do not exceed 5.8 pints (2.8 lbs. a.e.) per acre per year.
- When tank-mixing with products that contain 2,4-D, do not exceed a combined total of 2.8 pounds of a.e. per acre per year.
- Do not apply through any type of irrigation system.
- Do not apply by air.
- Do not apply in or near greenhouses.

CEREAL GRAINS (Barley Millet, Rye, Triticale, Teff*, Wheat)

Target Weeds	This Product (Pt./Ac.)	Specific Use Instructions
Not Underseeded With Legumes Postemergence: Annual and Biennial broadleaf weeds	0.5 to 1.3**	Apply after grain is well-tillered (usually about 4 to 8 inches high). Do not spray grain in the boot to dough stage.
Not Underseeded With Legumes Postemergence: Perennial broadleaf weeds	1 to 2.6**	
Underseeded With Legumes	0.25 to 0.5**	Apply after grain is 8 inches tall. Do not spray grain in boot to dough stage. Do not spray Alfalfa or sweet clover unless the infestation is severe and injury to these legumes can be tolerated.
Emergency Weed Control in Triticale, Wheat: Perennial broadleaf weeds	2.6	Apply when weeds are approaching bud stage after the grain dough stage. Do not spray during the boot to dough stage. The 2.6 pints per acre per application can produce injury to Wheat. Balance the severity of the weed problem against the possibility of crop damage. Where perennial weeds are scattered, make spot treatment to minimize the extent of crop injury.

Pre-harvest Application	1.0	Apply using air or ground equipment to control weeds that could interfere with harvest or to suppress perennial weeds. Apply when grain is in dough stage. Do not apply from early boot through the milk stage of growth.
<p>*Not for use on Teff in California. **Use the lower rate if small annual and biennial weeds are the major problem. Use the higher rate if perennial weeds or annual and biennial weeds are present which are in the hard-to-kill categories as determined by local experience. The higher rates increase the risk of grain injury and should be used only where the weed control problem justifies the grain damage risk. Do not apply this product to grain in the seedling stage.</p>		

Tank-mixtures:

This product can be used in combination with Metsulfuron-methyl, Metsulfuron-methyl plus Chlorsulfuron, Thifensulfuron-methyl plus Tribenuron-methyl or Tribenuron-methyl in Spring and Winter Wheat and Barley to control resistant weeds such as Kochia and Russian thistle.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank-mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank-mixture.

Use Precautions:

- For aerial application on grain, apply this product in 3 to 10 gallons of water per acre.
- For ground application, apply in a minimum of 10 to 15 gallons of water per acre for proper spray coverage.

Use Restrictions:

- Do not harvest within 14 days of application.
- Post-emergence: Maximum of 2.6 pints (1.25 lbs. a.e.) per acre per application. Limit to 1 post-emergence application per crop cycle.
- Pre-harvest: Maximum of 1 pint (0.5 lb. a.e.) per acre per application. Limit to 1 pre-harvest application per crop cycle.
- Limit to 3.6 pints product (1.75 lbs. a.e.) per acre per crop cycle.
- Do not feed treated straw to livestock if an emergency treatment as described above is applied.

CEREAL GRAINS (Oats*)

Crop	This Product (Pt./Ac.)	Specific Use Instructions
Not underseeded with Legumes: Spring Planted Oats	0.5 to 2**	Apply in sufficient water to give good coverage. Apply after fully tillered except during the boot to dough stage.
Not underseeded with Legumes: Fall Planted Oats	1 to 1.5**	Apply after full tillering but before early boot stage. Some difficult weeds may require higher application rates per acre for maximum control, but injury may result. Do not spray during or immediately following cold weather.
Underseeded with Legumes: Oats	0.25 to 0.5**	Apply after grain is 8 inches tall. Do not apply before tillering or from early boot through the milk stage of growth. Do not spray Alfalfa or Sweet clover unless the infestation is severe and injury to these legumes can be tolerated.
Pre-harvest	1	Apply when grains are in the hard dough stage to control large weeds that may interfere with harvest. Best results will be obtained when soil moisture is sufficient to cause succulent weed growth.

*Oats are less tolerant to 2,4-D than Wheat or Barley and more likely to be injured.
**Use the lower rate if small annual and biennial weeds are the major problem. Use the higher rate if perennial weeds or annual and biennial weeds are present which are in the hard-to-kill categories as determined by local experience. The higher rates increase the risk of grain injury and should be used only where the weed control problem justifies the grain damage risk. Do not apply this product to grain in the seedling stage.

Use Restrictions:

- Do not harvest within 14 days of application.
- Post-emergence: Maximum of 2.6 pints (1.25 lbs. a.e.) per acre per application. Limit to 1 post-emergence application per crop cycle.
- Pre-harvest: Maximum of 1 pint (0.5 lb. a.e.) per acre per application. Limit to 1 pre-harvest application per crop cycle.
- Limit to 3.6 pints product (1.75 lbs. a.e.) per acre per crop cycle.

CORN (Field, Pop, Sweet)

Time of Application / Growth Stage	This Product (Pt./Ac.)	Specific Use Instructions
Pre-plant (Burndown)	1 to 2	Use high rate in rate range for less susceptible weed or cover crops, weeds in advanced stages of development, or under less favorable growth conditions.
Pre-emergence and Reduced tillage	2	Pre-plant: Apply 7 to 14 days before planting Corn to control emerged broadleaf weed seedling or existing cover crops. Pre-emergence: Apply any time after planting but before Corn emerges to control broadleaf weed seedlings or existing cover crops. The seed furrow must be completely closed at application or severe crop injury may result.
Post-emergence Annual broadleaf weeds: Crop up to 8 inches tall	0.5 to 1	Apply when weeds are small and Corn is less than 8 inches tall (to top of crop canopy). If Corn is more than 8 inches tall, use drop nozzles and directed sprays to keep spray off foliage.
Post-emergence Annual broadleaf weeds: Crop 8 inches tall to tasseling (directed spray only)	1	Treat perennial weeds when they are in bud to bloom stage. Do not apply from tasseling to hard dough stage.
Post-emergence: Perennial broadleaf weeds	1	
Pre-harvest (Field Corn & Popcorn only)	Up to 3	Apply after Corn is in hard dough (or denting) stage. Do not apply pre-harvest to Sweet Corn.

Use Precautions:

- Do not apply pre-plant or pre-emergence to light sandy soils.
- Corn hybrids vary in tolerance to 2,4-D. Some are easily injured. Apply only to varieties known to be tolerant to 2,4-D. Consult the seed company or your Agricultural Experiment Station or Extension Service Weed Specialist for this information.
- Corn treated with 2,4-D may exhibit stem brittleness for 8 to 10 days following application. During this period, the crop is more susceptible to stem breakage from cultivation or wind.

Use Restrictions (Field Corn and Popcorn):

- Do not harvest within 7 days of application.
- Do not use treated crop as fodder for 7 days following application.
- Do not apply more than 6.3 pints of this product (3 lbs. a.e.) per acre per crop cycle.
- Pre-plant or Pre-emergence: Do not apply more than 2.1 pints of this product (1 lb. a.e.) per acre per application. Limit to 1 pre-plant or 1 pre-emergence application.
- Post-emergence: Do not apply more than 1 pint of this product (0.5 lb. a.e.) per acre per application. Limit to 1 post-emergence application per crop cycle.
- Pre-harvest: Do not apply more than 3.1 pints of this product (1.5 lbs. a.e.) per acre per application. Limit to 1 pre-harvest application per crop cycle.

Use Restrictions (Sweet Corn):

- Do not apply within 45 days of harvest of Corn.
- Do not use treated crop as fodder for 7 days following application.
- Minimum of 21 days between applications.
- Do not apply more than 3.1 pints of this product (1.5 lbs. a.e.) per acre per crop cycle.
- Pre-plant or Pre-emergence: Do not apply more than 2.1 pints of this product (1 lb. a.e.) per acre per application. Limit to 1 pre-plant or pre-emergence application per crop cycle.
- Post-emergence: Do not apply more than 1 pint of this product (0.5 lb. a.e.) per acre per application. Limit to 1 post-emergence application per use season.

CRANBERRIES*

Time of Application	This Product (Pt./Ac.)	Specific Use Instructions
Late June and July (for best results)	2.4	To control tall weeds in Cranberry bogs, apply with a wooden frame or similar device shaped like a hockey stick with its lower member wrapped with several thicknesses of Turkish toweling (or other suitable material). Apply by soaking the toweling in one part of this product to two parts water. Then, with swabbed portion of the stick horizontal, wave left and right above the cranberry vines, wiping small quantities of the herbicide onto tall weeds above the crop level.
*For use in the states of Massachusetts, New Jersey, Oregon, Washington and Wisconsin only.		

Use Precautions:

- INJURY TO CROPS FROM THIS HERBICIDE MAY OCCUR. IF YOU ARE NOT PREPARED TO ACCEPT SOME DEGREE OF CROP INJURY, DO NOT USE THIS PRODUCT ON CRANBERRIES.
- Avoid drip or contact of this product to Cranberry vines as it may kill or cause significant injury to Cranberry.

Use Restrictions:

- The pre-harvest interval (PHI) is 30 days.
- Limited to 2 applications per year.
- Do not exceed 2.5 pints (1.2 lbs. a.e.) per acre per application.
- When tank-mixing with products that contain 2,4-D, do not exceed a combined total of 1.2 pounds of a.e. per acre per year.
- Do not apply through any type of irrigation system.
- Do not apply by air.

FALLOW LAND* AND CROP STUBBLE

Target Weeds	This Product (Pt./Ac.)	Specific Use Instructions
Annual broadleaf weeds	1 to 2	Use a lower rate in the rate range when weeds are small (2 to 3 inches tall) and actively growing. Use a higher in the rate range when weeds are larger and under less favorable growth conditions.
Biennial broadleaf weeds	2 to 4	Apply when musk thistles or other biennial species are in the seedling to rosette stage and before development of flower stalks. The lower rate can be used in the Spring during the rosette stage. Use the highest rate in the Fall or after flower stalks have developed.
Perennial broadleaf weeds	2 to 4	Apply when perennial weeds are in bud to early bloom stage or while in good vegetative growth.
Wild garlic and Onion in crop stubble	4	Apply to new regrowth of wild garlic or onion which occurs in the Fall after harvest of small grains, Corn or grain Sorghum.
*Fallow land is defined as idle land, post-harvest to crops or between crops.		

Use Precautions:

- For best weed control results, do not cultivate for at least 2 weeks after application or until top growth is dead.

Use Restrictions:

- Plant only labeled crops within 29 days following application.
- Minimum of 30 days between applications.
- Do not apply more than 4.2 pints of this product (2 lbs. a.e.) per acre per application.
- Limit to 2 applications per year.

Planting in Treated Areas

Labeled Crops: Within 29 days after an application of this product, plant only those crops listed on this or other registered 2,4-D labels. Follow more stringent limitations, if any, provided in directions for specific crops. Labeled crops may be at risk of crop injury or loss if planted soon after application, especially during the first 14 days. When weighing this risk, consider the degradation factors described below.

Other Crops: All other crops may be planted 30 or more days after application without concern for illegal residues in the planted crop. However, under certain conditions, there may be a risk of injury to susceptible crops. Degradation factors described below should be considered in weighing this risk. Under normal conditions, any crop may be planted without risk of injury if at least 90 days of soil temperatures above freezing have elapsed since application.

Degradation Factors: When planting into treated areas, the risk of crop injury is less if lower rates of product were applied and conditions following application have included warm, moist soil conditions that favor rapid breakdown of 2,4-D. Risk is greater if higher rates of product were applied and soil temperatures have been cold and/or soils have been excessively wet or dry in the days following application. Consult your local agricultural extension service or information about susceptible crops and typical conditions in your area.

FILBERTS (HAZELNUT)*

Target	This Product (Pt./Ac.)	Specific Use Instructions
Annual broadleaf weeds; Suckers	1.5 to 2	Apply in 100 gallons of water per acre. Spray to the point of runoff when suckers are 6 to 9 inches tall. Use nozzles with large orifice nozzles and low tank pressure. Apply when needed from April through August.
*Not for use in California.		

Use Restrictions:

- Do not apply within 45 days of harvest.
- Allow at least 30 days between applications.
- Do not make more than 4 applications per year.
- Do not apply more than 2.1 pints of this product (1 lb. a.e.) per 100 gallons of spray solution per application.

GRAPE VINEYARDS*

Time of Application	This Product (Pt./Ac.)	Specific Use Instructions
After shatter following bloom and before Grape shoots reach the ground or during dormant season	1.8 to 2.8	Use this product on Grape vineyards established for at least 3 years to control Canada thistle, Field bindweed (Morningglory) and other 2,4-D susceptible broadleaf weeds when weeds are in the bud to early bloom stage and are growing vigorously. Dilute in 10 to 100 gallons of water to treat 1 acre of ground to be sprayed. For band or spot treatment, calculate rates according to the actual portion of acre treated. Use a hooded boom and low pressure flooding nozzles to deliver coarse droplets.
*For use in California only.		

Use Precautions:

- Grapes are extremely sensitive to 2,4-D. Use directed application so no 2,4-D contacts Grape leaves and young shoots or stems.

Use Restrictions:

- The pre-harvest interval (PHI) is 100 days.
- Limited to 1 application per crop cycle.
- Maximum of 2.8 pints (1.36 lbs. a.e.) per acre per application.
- When tank-mixing with products that contain 2,4-D, do not exceed a combined total of 1.36 pounds of a.e. per acre per year.

HOPS

Time of Application	This Product (Pt./Ac.)	Specific Use Instructions
Post-emergence	1	Direct application to the row middles.

Use Restrictions:

- Do not apply within 28 days of harvest.
- Allow at least 30 days between applications.
- Do not make more than 3 applications per crop cycle.
- Do not apply more than 1 pint of this product (0.5 lb. a.e.) per acre per application.
- Do not apply more than 3.1 pints of this product (1.5 lbs. a.e.) per acre per crop cycle.

ORCHARD FLOORS (Pome Fruits*, Stone Fruits, Tree Nuts*** and Pistachios)**

Time of Application / Target Weeds	This Product (Pt./Ac.)	Specific Use Instructions
Post-emergence: Annual and Biennial weeds	1 to 2	For application to orchard floors, use coarse, low pressure sprays and sufficient water for thorough coverage of weeds.
Post-emergence: Perennial weeds	Up to 4	Apply to annual weeds when small and actively growing. Apply to perennial weeds from bud to bloom stage.
<p>*Pome fruits including Apples, Crabapples, Loquat, Mayhaw, Oriental pear, Pears, Quince **Stone fruits including Apricot, Chickasaw plum, Damson plum, Fresh prunes, Japanese plums, Nectarines, Peaches, Plums, Plumcot, Sweet cherry, Tart Cherry ***Tree nuts including Almonds, Beech nut, Black walnut, Brazil nut, Butternut, Cashew, chestnut, Chinquapin, English walnut, Hickory nut, Macadamia nut (bush nut), Pecan. (Excludes Filberts. For use on Filberts, see "FILBERTS" section.)</p>		

Use Precautions:

- To avoid tree injury, do not allow spray drift to contact foliage, fruit, stems, trunks or trees or exposed roots.
- Avoid application immediately before irrigation and withhold irrigation for 2 days before and 3 days after application.
- Newly established trees or young orchards are more susceptible to 2,4-D injury. Apply only to orchards that have been established for at least one year and are in vigorous growth condition.

Use Restrictions:

- Do not apply when orchards are blooming.
- Do not make orchard floor applications in areas with light sandy soils.
- Pome Fruits: Allow at least 75 days between applications. Do not apply within 14 days of harvest.
- Stone Fruits: Allow at least 75 days between applications. Do not apply within 40 days of harvest.
- Tree Nuts and Pistachios: Allow at least 30 days between applications. Do not apply within 60 days of harvest.
- Do not apply more than 4.2 pints of this product (2 lbs. a.e.) per acre per application.
- Do not cut orchard floor forage or hay within 7 days after last application;
- Do not make more than 2 applications per year.
- Do not apply more than a total of 8.4 pints of this product (4 lbs. a.e.) per acre per use season.

POTATOES (RED)*

Time of Application	This Product (Pt./Ac.)	Specific Use Instructions
Post-emergence	0.15 (2.35 fl. ozs.)	<p>Properly timed applications of this product generally enhance red color, aid in storage retention of red color, improve skin appearance, increase tuber set, and improve tuber size uniformity (fewer jumbos). Crop response may vary depending on variety, stress factors, and local conditions. Varieties with naturally dark red color generally benefit less from treatment.</p> <p>Apply 2.35 fl. ozs. of this product per acre in 5 to 25 gallons of water using ground or aerial equipment when Potatoes are in the pre-bud stage (about 7 to 10 inches high) followed by a second application about 10 to 14 days later. The specific spray volume selected should be sufficient for good coverage of plants.</p> <p>Consult with Agricultural Extension Service and other qualified crop advisors for local recommendations.</p>

*For use on Red Potatoes intended for fresh market only.

Use Restrictions:

- The pre-harvest interval (PHI) is 45 days.
- Minimum of 10 days between applications.
- Maximum of 2.35 fluid ounces (0.07 lb. a.e.) per acre per application.
- Limited to 2 post-emergence application per crop cycle.

RICE*

Time of Application	This Product (Pt./Ac.)	Specific Use Instructions
Pre-plant	1 to 2	Apply 2 to 4 weeks before planting rice to control emerged broadleaf weeds.
Post-emergence	1 to 2**	Apply when rice is in late tillering stage and at the time of first joint development (first to second green ring).

*Not for use in California.
**Up to 3 pints of this product (1.5 lbs. a.e.) per acre may be applied post-emergence for difficult weed control situations. However, there is greater risk of crop injury at rates greater than 2 pints per acre. Use such rates only when the need for weed control justifies additional risk to the crop.

Use Precautions:

- Do not apply at early seedling stage or after rice internodes exceed one-half inch or panicle initiation.
- Some rice varieties under certain conditions or stages of growth may be injured by 2,4-D. Before applying, consult local university or agricultural extension service specialists regarding for local treatment recommendations for various rice varieties.

Use Restrictions:

- Do not apply within 60 days of harvest.
- Pre-plant: Do not apply more than 2.1 pints of this product (1 lb. a.e.) per acre per application. Do not apply more than 1 pre-plant application per crop cycle.
- Post-emergence: Do not apply more than 3.1 pints of this product (1.5 lb. a.e.) per acre per application. Do not apply more than 1 post-emergence application per crop cycle.
- Do not apply more than 3.1 pints of this product (1.5 lbs. a.e.) per acre per crop cycle.

RICE (WILD)*

Target Weed	This Product (pt./Ac.)	Specific Use Instructions
Common water plantain	0.5	Broadcast in 4 to 10 gallons of total spray when wild rice is in 1 to 2 aerial leaf to early tillering state and after Water plantain has emerged from the water and before wild rice has reached the boot stage. Do not spray after wild rice has reached the boot stage.
*For use in Minnesota only on Wild rice grown in commercial paddies.		

Use Precautions:

- Do not apply to Wild rice growing in lakes or streams.
- Do not use water drained out of Wild rice paddies to irrigate other crops.
- In order to protect federally listed endangered species, Minnesota Dept. of Agriculture has a program to pre-notify landowners where pesticide applications may affect federally listed endangered or threatened species.

Use Restrictions:

- Do not apply within 60 days of harvest.
- Do not make more than 1 application per crop cycle.
- Do not apply more than 0.5 pint of this product (0.25 lb. a.e.) per acre per application.

SORGHUM [Grain Sorghum (Milo) and Forage Sorghum]

Time of Application / Growth Stage	This Product (Pt./Ac.)	Specific Use Instructions
Post-emergence – Directed Spray Crop 6 to 8" tall	0.5 to 1.5	Apply when Sorghum is 6 to 15 inches tall. If Sorghum is more than 8 inches tall (to top of crop canopy), use drop nozzles and apply as a directed spray to keep spray off foliage.
Post-emergence - Directed Spray Only: Crop 8 to 15" tall	0.75 to 1.5	

Use Precautions:

- Temporary crop injury can be expected under conditions of high soil moisture and high air temperatures. If it is necessary to apply this product under these conditions, use no more than 0.66 pint of this product per acre.
- Sorghum hybrids vary in tolerance to 2,4-D. Some are easily injured. Apply only to varieties known to be tolerant to 2,4-D. Consult the seed company or your agricultural experiment station or extension service weed specialist for this information.

Use Restrictions:

- Do not apply during boot or later stages of growth.
- Do not apply within 30 days of harvest of Sorghum.
- Do not permit meat or dairy animals to consume treated crop as fodder or forage within 30 days following application.
- Do not apply more than 1 post-emergence application per crop cycle.
- Do not apply more than a total of 2.1 pints of this product (1 lb. a.e.) per crop cycle.

SOYBEANS

Time of Application	This Product (Pt./Ac.)	Specific Use Instructions
Pre-plant burndown	0.75 to 1	Apply no less than 15 days before planting Soybeans when weeds are small and actively growing. Use the higher rate on larger weeds and when perennial weeds are present. See "Use Precautions" and "Use Restrictions" below.
	1 to 2	Apply not less than 30 days before planting Soybeans when weeds are small and actively growing. Use the higher rate on larger weeds and when perennial weeds are present. See "Use Precautions" and "Use Restrictions" below.

Tank-mixture:

This product may be applied pre-plant to Soybeans in tank-mixtures with other herbicides such as Glyphosate, Imazaquin, Paraquat, Pendimethalin, Pendimethalin plus Imazethapyr, Sethoxydim and others that are registered for pre-plant use in Soybeans.

Crop oil concentrates, agricultural surfactants and fluid fertilizers approved for use on growing crops may be added to spray mixtures of this product to increase the herbicidal effectiveness on certain weeds.

Read and follow all directions and precautions on this label and on the label of each product added to the spray mixture. Follow the most restrictive labeling. Refer to the "MIXING" section of this label for tank-mixing instructions and compatibility testing.

Use Precautions:

- Unacceptable injury to Soybeans planted in treated fields may occur. Whether or not Soybean injury occurs and the extent of such injury will depend on weather (temperature and rainfall) from herbicide application until Soybean emergence and agronomic factors such as the amount of weed vegetation and previous crop residue present at the time of application. Injury is more likely under cool rainy conditions and where there is less weed vegetation and crop residue present.
- Do not disturb treated soil through tillage between application and planting of Soybeans.
- In treated fields, plant Soybean seed as deep as practical, but not less than 1 inch deep. Adjust the planter, if necessary, to ensure that planted seed is adequately covered.
- **Unless you are prepared to accept the results of soybean injury, including possible stand loss and/or yield reduction, do not pre-plant apply this product in Soybeans**

Use Restrictions:

- Do not use on sandy soils with less than 1% organic matter.
- Do not apply more than 2.1 pints of this product (1 lb. a.e.) per acre per crop cycle.
- One (1) or 2 pre-plant applications are allowed per crop cycle. If a single pre-plant application is made, do not apply more than 2.1 pints of this product (1 lb. a.e.) per acre per application. Apply no less than 30 days prior to planting Soybeans. If 2 pre-plant applications are made, do not apply more than 1 pint of this product (0.5 lb. a.e) per acre per application. Apply no less than 15 days prior to planting Soybeans.
- During the growing season following application, do not replant treated fields with crops other than those labeled for use with this product.

STRAWBERRIES (Established Plantings Only)*

Time of Application	This Product (Pt./Ac.)	Specific Use Instructions
Early spring	2 to 3	Apply in 25 to 50 gallons of water per acre in early Spring on established Strawberry plantings when Strawberries are dormant or immediately after the last picking. DO NOT apply unless possible injury to Strawberry crop is acceptable. Follow recommendations of State Extension Horticultural Specialist in your area.
*Not for use in California and Florida.		

Use Restrictions:

- Do not apply more than 3.1 pints of this product (1.5 lbs. a.e.) per acre per application.
- Do not apply more than 1 application per crop cycle.

SUGARCANE

Time of Application	This Product (Pt./Ac.)	Specific Use Instructions
Pre-emergence, Post-emergence	2 to 4	Consult your agricultural experiment station or extension service weed specialist for local recommendations. Pre-emergence: Apply before cane emerges to actively growing weeds. Post-emergence: Apply after cane emerges through canopy closure. Use higher rate for perennial weeds and difficult-to-control species.

Use Restrictions:

- Do not harvest cane prior to maturity.
- Pre-emergence: Do not apply more than 4.2 pints of this product (2 lbs. a.e.) per acre per application. Limit to 1 pre-emergence application per crop cycle.
- Post-emergence: Do not apply more than 4.2 pints of this product (2 lbs. a.e.) per acre per application. Limit to 1 post-emergence application per crop cycle.
- Do not apply more than a total of 8.4 pints of this product (4 lbs. a.e.) per acre per use season.

SUGARCANE*

Time of Application	This Product (Pt./Ac.)	Specific Use Instructions
Pre-emergence, Post-emergence	1 to 3	If used in the islands of Maui and Kauai, the general wind restriction is raised to 20 mph. When applying in winds in excess of 15 mph, the following requirements are in effect: Aerial Application: <ul style="list-style-type: none"> • No application shall be made within a distance of 1000 feet of sensitive areas such as Nature Preserves, Wildlife Refuges, Parks, Lakes, Reservoirs, Rivers, Streams, Non-irrigation Canals, Natural Ponds, Estuaries, Wetlands, Intertidal Areas, Ecologically Significant Grasslands, homes, public or private buildings, or fields with crops other than sugarcane whenever these sensitive areas are downwind from the spray areas and subject to possible spray drift. In instances where these sensitive areas are upwind from the spray area, the minimum restricted distance shall be 300 feet. • Apply only as a coarse or coarser spray (ASAE standard 572 or a volume mean diameter of 385 microns). • Use a spray drift retardant and/or other measures known to control drift. Ground Broadcast Applications: <ul style="list-style-type: none"> • Apply by ground boom with nozzle height no more than 2 feet above ground (pre-emergence) or crop canopy (post emergent broadcast) applications or, for directed sprays, no more than 1 foot above the ground, or 1.25 feet (15 inches) for better spray patterns without boom levelers on uneven terrain. • Apply only as a coarse or coarser spray (ASAE standard 572) or a volume mean diameter of 385 microns. • Use spray drift retardants and/or other measures known to control drift. Applications techniques to reduce off-site drift include, but are not limited to, the use of hooded or shielded sprayers or other means to reduce drift.
*For use in Hawaii only.		

Use Restrictions:

- Do not harvest cane prior to crop maturity.
- Do not apply more than 8.4 pints of this product (4 lbs. a.e.) per acre per year.
- Limit to 1 pre-emergence application per crop cycle. Do not apply more than 4.2 pints of this product (2 lbs. a.e.) per acre per application.
- Post-emergence: Do not apply more than 4.2 pints of this product (2 lbs. a.e.) per acre per application. Lay-by application can be made but crop damage may occur in some Sugarcane varieties,

FORESTRY USES

Forest Site Preparation, Forest Roadsides, Brush Control, Established Conifer Release (Including Christmas Trees and Reforestation Areas)

When this product is applied by tree injection, follow re-entry requirements given in the "NON-AGRICULTURAL USE REQUIREMENTS" section of this label.

Site / Application Method	This Product	Specific Use Instructions
Annual weeds	2 to 4 pts. per acre	Apply when weeds are small and growing actively before the bud stage. Apply when biennial and perennial species are in the seedling to rosette stage and before flower stalks appear. For difficult to control perennial broadleaf weeds and woody species, use up to 1 gallon of this product and appropriate rates of Triclopyr per acre. For conifer release, make application in early Spring before budbreak of Conifers when weeds are small and actively growing.
Biennial broadleaf weeds, Perennial broadleaf weeds, Susceptible woody plants	4 to 8 pts. per acre	
Spot Treatment: Broadleaf weeds	See <i>"Specific Use Instructions"</i>	To control broadleaf weeds in small areas with a handheld sprayer, use an application rate equivalent to the specified broadcast rate. Spray to thoroughly wet all foliage. Mix 1.28 fluid ounces per gallon of spray solution and apply through pump-up or backpack sprayer. A non-ionic surfactant may be added to improve coverage. Refer to <i>"Rate Conversion Table for Spot Treatment"</i> and instructions for <i>"Spot Treatment"</i> in the <i>"APPLICATION"</i> section of this label.
Poplar/Cottonwood Trees Grown for Pulp: Broadleaf weeds	0.5 to 3 pts. per acre	Apply using wick applicators or conventional ground sprayers (except for irrigation systems). Do not allow this product to contact leaves or green bark of the tree. Apply in sufficient water for uniform coverage to or after planting. Application during warm weather is preferred. Apply when weeds are actively growing, preferably before bud stage. Repeat application may be necessary for less susceptible weeds. Reapply as needed. This product may be tank-mixed with Glyphosate to broaden weed control. A spreader such as Induce® may be added to improve efficacy. Read and follow all directions and precautions on this label and on the label of each product added to the spray mixture. Follow the most restrictive labeling. Tree Injection: This product may also be used as tree injection to control unwanted trees in Poplar/Cottonwood trees grown for pulp. Refer to the <i>"TREE INJECTION"</i> section below for use directions.

(Continued)

(Continuation)

Site / Application Method	This Product	Specific Use Instructions
Conifer Release: Species such as Balsam fir, Pines (Jack, Ponderosa, Red, White) Spruce (Black, White)	3 to 6 pts. per acre	To control competing hardwood species such as Alder, Aspen, Birch, Hazel, and Willow, apply from mid- to late summer when growth of conifer trees has hardened off and woody plants are still actively growing. Apply with ground or air equipment, using sufficient spray volume to ensure complete coverage. Because this treatment may cause occasional conifer injury, do not apply if such injury cannot be tolerated.
Directed Spray: Conifer plantations including Pines	8 pts. per 100 gals.	Apply when brush or weeds are actively growing by directing the spray so as to avoid contact with conifer foliage and injurious amounts of spray. Apply in oil, oil-water, or water carrier in a spray volume of 10 to 100 gallons per acre.
Basal Spray (May also be used in Rangeland, Pastures, and Noncropland)	16 pts. per acre	Thoroughly wet the base and root collar of all stems until the spray begins to accumulate around the root collar at the ground line. Wetting stems with the mixture may also aid in control.
Surface of Cut Stumps	or	Apply as soon as possible after cutting trees. Thoroughly soak the entire stump with the 2,4-D mixture including cut surface, bark and exposed roots.
Frill and Girdle	2.6 fl. ozs. per gal. of water	Cut frills (overlapping V-shaped notches cut downward through the bark in a continuous ring around the base of the tree) using an axe or other suitable tool. Treat freshly cut frills with as much of the 2,4-D mixture as they will hold.
Tree Injection	1 to 2 mL per injection site	To control unwanted hardwood trees as Alder, Aspen, Birch, Blackgum, Cherry, Elm, Hickory, Oak, Sweetgum, and Tulip poplar in forests and other noncrop areas, apply by injecting 1 ml of this product, undiluted, per inch of trunk diameter at breast height (DBH) as measured approximately 4.5 feet above the ground. Make injections as close to the root collar as possible and the injection bit must penetrate the inner bark. Applications may be made throughout the year, but for best results, apply between May 15 and October 15. For hard to control species such as Ash, Maple, and Dogwood, use 2 mL of this product, undiluted, per injection site or double the number of 1 mL injections. Do not treat Maples during the Spring sap flow. Note: No Worker Protection Standard worker entry restrictions or worker notification requirements apply when this product is directly injected into agricultural plants.

Use Precautions:

- Do not allow sprays to contact conifer shoot growth (current year's new growth) or injury may occur.
- Do not apply to nursery seed beds.
- For conifer release, do not use on plantations where Pine or Larch are among the desired species.

Use Restrictions:

- For broadcast applications, do not apply more than a total of 8.4 pints of this product (4 lbs. a.e.) per acre per year. Limit to 1 broadcast application per year.
- For basal spray, cut surface stumps, and frill applications, do not apply more than 16.8 pints of this product (8 lbs. a.e.) per 100 gallons of spray solution. Limit to 1 basal spray or cut surface application.
- For tree injection, do not apply more than 2 mL of this product per injection site. Limit to 1 injection application per year.

WEED CONTROL IN HYBRID POPLAR TREES, COTTONWOOD TREES AND WILLOW TREES GROWN AS BIOENERGY CROPS*

Follow re-entry requirements given in the "NON-AGRICULTURAL USE REQUIREMENTS" section of this label.

Target Weeds	This Product (Pt./Ac.)	Specific Use Instructions
Broadleaf weeds	0.5 to 4	Apply when weeds are actively growing preferably before bud stage. Application during warm weather is preferred. Repeat treatments may be necessary for less susceptible weeds. Reapply as needed. For hybrid poplar, cottonwood and willow, apply prior to or after planting. By ground, use 0.5 to 3 pints of this product per acre in minimum 10 gallons of water per acre for broadcast application. For wick type applicators, use 1 to 4 pints per acre. Crop injury may result if wick, wick solution or spray solution contact leaves or green bark of crop trees. Do not spray immediately before irrigation. Withhold above-ground irrigation for 3 days after application. Tank-mixture: This product may be tank-mixed with Glyphosate to provide broader spectrum of weed control. Read and follow all directions and precautions on this label and on the label of the product added to the spray mixture. Follow the most restrictive labeling.
*Not for use in California.		

Use Precautions:

- Exercise extreme care to avoid contact of spray solution, spray drift, or mist with tree foliage, green bark of trunks, stems or exposed roots of poplar, cottonwood and willow trees. Contact of this product can result in serious damage. Even when using extreme care in application of this product, injury to these trees from this herbicide may occur. DO NOT use this product if you are not prepared to accept some degree of crop injury.
- Treated plantings are not for human or animal consumption.

Use Restrictions:

- Limit to 1 broadcast application per year.
- Do not apply more than 4.2 pints of this product (2 lbs. a.e.) per acre per application.
- Minimum of 30 days between applications.
- Do not use in or near greenhouses.
- Do not apply by air or through any type of irrigation system.
- Do not use treated vegetation for forage or hay or allow livestock to graze treated fields.

WEED CONTROL IN GIANT MISCANTHUS, GIANT REEDGRASS, SWITCHGRASS, AND OTHER NONFOOD GRASS BIOENERGY CROPS*

Follow re-entry requirements given in the "NON-AGRICULTURAL USE REQUIREMENTS" section of this label.

Target Weeds	This Product (Pt./Ac.)	Specific Use Instructions
Broadleaf weeds	0.5 to 4	Apply 0.5 to 2 pints of this product per acre to seedling grasses in minimum 10 gallons of water by ground or in minimum 2 gallons of water by air. Use the rate of 1 to 4 pints per acre when grasses are well established. Do not spray immediately before irrigation. Withhold above-ground irrigation for 3 days after application.
*Not for use in California.		

Use Precautions:

- Do not apply through any type of irrigation system.
- Treated plantings are not for human or animal consumption.

Use Restrictions:

- Do not use treated grass for forage or hay or allow livestock to graze treated fields.
- Limit to 2 broadcast applications per year.
- Do not apply within 30 days of last application.
- Do not apply more than 4.2 pints of this product (2 lbs. a.e.) per acre per application.

RANGELAND, ESTABLISHED GRASS PASTURES (INCLUDING PERENNIAL GRASSLANDS NOT IN AGRICULTURAL PRODUCTION SUCH AS CONSERVATION RESERVE PROGRAMS [CRP] ACRES)

When this product is applied to rangeland and established Pastures not harvested for hay or seed, and when applied by tree injection, follow re-entry requirements given in the "NON-AGRICULTURAL USE REQUIREMENTS" section of this label.

Target / Application Method	This Product	Specific Use Instructions
Annual broadleaf weeds	2* pts. per acre	For best results, apply when weeds are small and when growing actively before the bud stage. Apply when musk thistles or other biennial species are in the seedling to rosette stage and before flower stalks appear. Refer to the "WEEDS CONTROLLED" section of this label for a listing of susceptible weeds and weed species that may only be partially controlled and require repeat applications and/or use of higher specified rates, even under ideal conditions of application.
Biennial broadleaf weeds, Perennial broadleaf weeds	2 to 4* pts. per acre	* In CRP areas, the rates may be reduced to 0.5 to 1 pint per acre to control annual broadleaf weeds in young grasses and 0.5 to 2 pints per acre to control annual broadleaf weeds in established grasses. Do not apply to young grasses with fewer than 6 leaves or prior to tillering as excessive injury may occur.
Spot treatment to control broadleaf weeds	See "Specific Use Instructions"	To control broadleaf weeds in small areas with a handheld sprayer, use an application rate equivalent to the specified broadcast rate. Spray to thoroughly wet all foliage. Mix 1.28 fl. ozs. per gallon of spray solution and apply through pump up or backpack sprayer. A non-ionic surfactant may be added to improve coverage. Refer to "Rate Conversion Table for Spot Treatment" and instructions for "Spot Treatment" in the "APPLICATION" section of this label.
Tree Injection Application	1 to 2 mL per injection site	See "Tree Injection" under the "FORESTRY USES" section for instructions.
Wild garlic and Wild onion	4 pts. per acre	Make 3 applications (Fall-Spring-Fall or Spring-Fall-Spring) starting in late Fall or early Spring.
Broadleaf weed control in newly sprigged coastal Bermudagrass	2 to 4 pts. per acre	Applications may be made either pre-emergence or post-emergence. Follow the above specific use directions for annual, biennial and perennial broadleaf weed control.
Sand shinnery oak, Sand sagebrush	2 pts. per acre	Sand shinnery oak: Apply by aircraft between May 15 and June 15. Sand sagebrush: Apply by ground or aircraft when foliage is fully expanded and plants are actively growing. Use a 1:4 oil-water emulsion as carrier and a spray volume of 3 to 5 gallons per acre.
Big sagebrush, Rabbitbrush	4 pts. per acre	Apply by ground or aircraft when foliage is fully expanded and plants are actively growing. Use a 1:4 oil-water emulsion as carrier and a spray volume of 3 to 5 gallons per acre. Retreatment may be needed.
Buckbrush, Chamise, Chaparral species, Coastal sage, Coyotebrush, Manzanita	4 pts. per acre	Apply by ground or aircraft when foliage is fully expanded and plants are actively growing. Use water or 1:4 oil-water emulsion as carrier and a spray volume of 5 to 10 gallons per acre. Retreatment may be needed.

(Continued)

(Continued)

Target / Application Method	This Product	Specific Use Instructions
Southern Wild Rose: Broadcast application	Up to 4 pts. per acre	Broadcast: Apply in a spray volume of 5 or more gallons per acre by aircraft or 10 or more gallons per acre by ground equipment. Spot treatment: Apply when foliage is well developed. Thorough coverage is required. Mix 1.28 fl. ozs. per gallon of spray solution. Apply through pump up or backpack sprayer. Add a non-ionic surfactant to improve coverage. Two or more treatments may be required.
Southern Wild Rose: Spot Treatment	1.28 fl. ozs. per gal. of spray solution	
Basal spray	16 pts. per 100 gals.	Thoroughly wet the base and root collar of all stems until the spray begins to accumulate around the root collar at the ground line. Wetting stems with the mixture may also aid in control.
Surface of Cut Stumps	or 2.5 fl. ozs. per gal. of water	Apply as soon as possible after cutting trees. Thoroughly soak the entire stump with the 2,4-D mixture including cut surface, bark and exposed roots.
Frill and Girdle		Cut frills (overlapping V-shaped notches cut downward through the bark in a continuous ring around the base of the tree) using an axe or other suitable tool. Treat freshly cut frills with as much of the 2,4-D mixture as they will hold.

Use Precautions:

- For program lands such as Conservation Reserve Program (CRP), consult program rules to determine whether grass or hay may be used. The more restrictive requirements of the program rules or this label must be followed.
- For annual, biennial and perennial broadleaf weeds control in CRP areas, use at least 2 gallons of water per acre by air and 5 gallons of water per acre by ground.

Use Restrictions:

- If grass is to be cut for hay, the Agricultural Use Requirements for the Worker Protection Standard are applicable.
- Do not use on Bentgrass, Alfalfa, Clover, or other legumes.
- Do not use on newly seeded areas until grass is well established.
- Do not use from early boot to milk stage where grass seed production is desired.
- Do not cut forage for hay within 7 days of application.
- Do not graze dairy cattle in treated areas for 7 days after application.
- Minimum of 30 days between applications.
- For susceptible annual and biennial broadleaf weeds, do not apply more than 2 pints of this product (1 lb. a.e.) per acre per application.
- For moderately susceptible biennial, perennial broadleaf weeds, use 2.1 to 4.2 pints of this product (1 to 2 lbs. a.e.) per acre per application.
- For difficult to control weeds and woody plants, use 4.2 pints of this product (2 lbs. a.e.) per acre per application.
- For spot treatment, use 4.2 pints of this product (2 lbs. a.e.) per acre per application.
- Do not make more than 2 applications per year.

NONCROPLAND**(Fencerows, Hedgerows, Roadsides, Drainage Ditches, Rights-of Way, Utility Power Lines, Railroads, Airports, and Industrial Sites)**

When this product is used in noncropland, follow re-entry requirements given in the "NON-AGRICULTURAL USE REQUIREMENTS" section of this label.

Site / Application Method	This Product	Specific Use Instructions
Annual broadleaf weeds	2 to 4 pts. per acre	Apply when annual weeds are small and actively growing before the bud stage. Biennial and perennial weeds should be at the rosette to bud stage but not flowering at the time of application. For difficult to control perennial broadleaf weeds and woody species, tank-mix up to 1 gallon of this product with appropriate rates of Triclopyr per acre.
Biennial broadleaf weeds, Perennial broadleaf weeds, Susceptible woody plants	4 to 8 pts. per acre	For ground application: (High volume) apply a total of 100 to 400 gallons per acre; (low volume) apply a total of 10 to 100 gallons per acre. For helicopter: Apply a total of 5 to 30 gals. per acre spray volume.
Spot treatment to control broadleaf weeds	See "Specific Use Instructions"	To control broadleaf weeds in small areas with a handheld sprayer, use an application rate equivalent to the specified broadcast rate. Spray to thoroughly wet all foliage. Mix 1.28 fluid ounces per gallon of spray solution and apply through pump up or backpack sprayer. A non-ionic surfactant may be added to improve coverage. Refer to "Rate Conversion Table for Spot Treatment" and instructions for "Spot Treatment" in the "APPLICATION" section of this label.
Tree Injection Application	1 to 2 mL per injection site	See "Tree Injection" under the "FORESTRY USES" section for instructions.
Southern Wild Rose: Broadcast application	Up to 4 pts. per acre	Broadcast: Apply in a spray volume of 5 or more gallons per acre by aircraft or 10 or more gallons per acre by ground equipment.
Southern Wild Rose: Spot treatment	1.28 fl. ozs. per gal. of spray solution	Spot treatment: Apply when foliage is well developed. Thorough coverage is required. Mix 1.28 fluid ounces per gallon of spray solution. Apply through pump up or backpack sprayer. Add a non-ionic surfactant to improve coverage. Two or more treatments may be required.

Use Precautions:

- Bentgrass, St Augustine, Clover, Legumes and Dichondra may be severely injured or killed by this treatment.

Use Restrictions:

- Do not apply to newly seeded areas until grass is well established.
- Applications to noncropland areas are not applicable to treatment of commercial timber or other plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.
- Annual and perennial broadleaf weeds: Do not apply more than 4.2 pints of this product (2 lbs. a.e.) per acre per application. Do not make more than 2 applications per season. Do not reapply to a treated area within 30 days of a previous application.
- Woody plants: Do not apply more than a total of 8.4 pints of this product (4 lbs. a.e.) per acre per year. Do not make more than 1 application per year.

GRASSES GROWN FOR SEED OR SOD FARMS

When this product is used in grasses grown for seed or sod, follow the PPE and re-entry requirements in the "AGRICULTURAL USE REQUIREMENTS" section of this label.

Site / Time of Application	This Product (Pt./Ac.)	Specific Use Instructions
Annual and perennial broadleaf weeds	2 to 4	Apply to established stands in Spring from tiller to early boot stage. Do not spray in boot stage. New Spring seedings may be treated with the lower rate after grass seedlings have at least 5 leaves. perennial weed regrowth may be treated in the Fall.
Grasses Grown for Seed (Post-emergence): Seedling grass (5-leaf stage or later)	0.75 to 1	Apply when weeds are small and actively growing. For best results, apply when soil moisture is adequate for active weed growth. Do not apply to newly seeded grasses until well established (5 leaf stage or later). Use a maximum of 1 pint of this product per acre. Cool season grasses are tolerant to higher rates. Do not apply to grass in the early boot through milk stage if seed production is desired. When grass is well established, higher rates of up to 4 pints per acre may be applied to control of hard-to-kill annual or perennial weeds.
Grasses Grown for Seed (Post-emergence): Well-established grasses	1 to 4	
Sod farms (Post-emergence)	0.5 to 4	Deep-rooted perennials such as Bindweed and Canada thistle may require repeat applications. Avoid mowing sod farms for 1 to 2 days before or after application. Delay irrigation until the day following application.

Use Precautions:

- Legumes may be damaged or killed.

Use Restrictions:

- Do not use on creeping grasses such as Bent except for spot treatment.
- Do not use on injury-sensitive southern grasses such as St. Augustinegrass.
- Do not use on Dichondra or other herbaceous ground covers.
- Reseeding: Delay reseeding at least 30 days following application. Preferably, with Spring application, reseed in the Fall and with Fall application, reseed in the Spring.
- Do not apply more than 4.2 pints of this product (2 lbs. a.e.) per acre per application.
- Minimum of 21 days between applications.
- Do not make more than 2 applications per year (excluding spot treatments).

ORNAMENTAL TURF (Includes Lawns, Golf Courses, Cemeteries, Parks, Airfields, Roadsides, Vacant Lots, Drainage Ditch Banks)

When this product is used in ornamental turf, follow the PPE and re-entry requirements in the "NON-AGRICULTURAL USE REQUIREMENTS" section of this label.

Site / Application Method	This Product (Pt./Ac.)	Specific Use Instructions
Post-emergence: Seedling grass (5-leaf stage or later)	0.75 to 1	Apply when weeds are small and actively growing. For best results, apply when soil moisture is adequate for active weed growth. Deep-rooted perennial weeds such as Bindweed and Canada thistle may require repeat applications. Do not apply to newly seeded grasses until well established (5-leaf stage or later). Use a maximum of 1 pint of this product per acre. Cool season grasses are tolerant of higher rates.
Post-emergence: Annual broadleaf weeds, Well-established grasses	2 to 3	
Post-emergence: Biennial broadleaf weeds, Perennial broadleaf weeds	3	

Use Precautions:

- Do not use on creeping grasses such as Bent except for spot treatment.
- Do not use on injury-sensitive southern grasses such as St. Augustinegrass.
- Do not use on Dichondra or other herbaceous ground covers. Legumes may be damaged or killed.
- Reseeding: Delay reseeding at least 30 days following application. Preferably, with Spring application, reseed in the Fall and with Fall application, reseed in the Spring.

Use Restrictions:

- Do not apply more than 3.1 pints of this product (1.5 lbs. a.e.) per acre per application.
- Do not apply more than 6.3 pints of this product (3 lbs. a.e.) per acre per year (excluding spot treatments).
- Do not make more than 2 broadcast applications per year (excludes spot treatments).

AQUATIC USES

CONTROL OF WEEDS AND BRUSH ON BANKS OF IRRIGATION CANALS AND DITCHES

Target Plants	This Product (Pt./Ac.)	Specific Use Instructions
Annual weeds	2 to 4	Use 2 gallons or more of spray solution per acre. Apply using low pressure spray (10 to 40 psi) in a spray volume of 20 to 100 gallons per acre using power operated spray equipment. Apply when wind speed is low, 5 mph or less. Apply working upstream to avoid accidental concentration of spray into water. Cross-stream spraying to opposite banks is not permitted and avoid boom spraying over water surface. When spraying shoreline weeds, allow no more than a 2 foot overspray onto water surface with an average of less than 1 foot of overspray to prevent significant water contamination. Apply when weeds are small and growing actively before the bud stage. Apply when biennial and perennial species are in the seedling to rosette stage and before flower stalks appear. For hard-to-control weeds, a repeat application after 30 days at the same rate may be needed. For woody species and patches of perennial weeds, mix 1 gallon of this product in 64 to 150 gallons of total spray. Wet foliage by applying about 3 to 4 gallons of spray per 1,000 square feet (10.5 X 10.5 steps).
Biennial and Perennial broadleaf weeds, Susceptible woody plants	4	

Use Restrictions:

- Limited to 2 applications per season.
- Do not apply within 30 days of first application.
- Do not apply more than 4.2 pints of this product (2 lbs. a.e.) per acre per application.
- Do not apply more than a total of 8.4 pints of this product (4 lbs. a.e.) per acre per use season.
- Spot treatment is permitted.
- Do not use on small canals with a flow rate less than 10 cubic feet per second (CFS) where water will be used for drinking purposes.

CFS may be estimated by using the formula below.

The approximate velocity needed for the calculation can be determined by observing the length of time that it takes a floating object to travel a defined distance. Divide the distance (ft.) by the time (sec.) to estimate velocity (ft. per sec.). Repeat 3 times and use the average to calculate CFS.

CFS = Average Width (ft.) x Average Depth (ft.) x Average Velocity (ft. per sec.)

For Ditchbank Weeds:

- Do not allow boom spray to be directed onto water surface.
- Do not spray cross-stream to opposite bank.

For Shoreline Weeds:

- Boom spraying onto water surface must be held to a minimum.
- Allow no more than 2 foot overspray onto water with an average of less than 1 foot overspray to prevent introduction of greater than negligible amounts of chemical into the water.

CONTROL OF AQUATIC WEEDS IN PONDS, LAKES, RESERVOIRS, MARSHES, BAYOUS, DRAINAGE DITCHES, CANALS, RIVERS AND STREAMS THAT ARE QUIESCENT OR SLOW MOVING INCLUDING PROGRAMS OF THE TENNESSEE VALLEY AUTHORITY

Notice to Applicators: Before application, coordination and approval of local and state authorities may be required, either by letter or agreement or issuance of special permits for such use.

Target Weeds	Rate Per Acre (Pints)	Specific Use Instructions
Floating and Emergent Aquatic Weeds	2.5 to 8	Apply when leaves are fully developed above waterline and are actively growing. Spray to wet foliage thoroughly. Contact your State Department or Game and Fish Commission for assistance in determining the best time and rate of application under your local conditions. Perennial and other hard-to-control weeds may require repeat applications for adequate control.
Water hyacinth (<i>E. crassipes</i>)	4 to 8	Apply on actively growing plants by surface and air applications. Spray the weed mass only. Use 8 pints when Water hyacinth plants are mature and when weed mass is dense. Repeat application as necessary to kill regrowth and plants missed in previous operation. Surface Application: Use power operated sprayers with boom or spray gun mounted on boat, tractor or truck. Thorough wetting of foliage is essential for maximum control. Use 100 to 400 gallons of spray mixture per acre. Take special precautions such as use of low pressure, large nozzles and spray thickening agents to avoid spray drift to susceptible crops. Follow label directions for use of any drift control agent. For Directaspra™ operation, use this product with 1 pint of drift control agent in 50 to 100 gallons of water. For other applications, follow the label of the drift control agent for mixing directions. Aerial Application: Use drift control spray equipment or thickening agent mixed in the spray mixture. Apply 8 pints of this product per acre using standard boom systems using a minimum spray volume of 5 gallons per acre. For Microfoil® drift control spray systems, apply this product in a total spray volume of 12 to 15 gallons per acre.

Use Restrictions:

- Do not use more than 8.4 pints of this product (4 lbs. a.e.) per surface acre per application.
- Do not make more than 2 applications per season.
- Minimum of 21 days between applications.
- Spot treatments are permitted.

Fish breathe dissolved oxygen in the water and decaying weeds also use oxygen. When treating continuous, dense weed masses, it may be appropriate to treat only part of the infestation at a time. For example, apply the product in lanes separated by untreated strips that can be treated after vegetation in treated lanes has disintegrated. During the growing season, weeds decompose in a 2 to 3 week period following treatment. Waters having limited and less dense weed infestations may not require partial treatments. Other local factors such as water exchange and sediment load can also influence the dissolved oxygen level. Coordination and approval of local and state authorities may be required, either by letter of agreement or issuance of special permits for aquatic applications.

Water Use:**1 . Water for Irrigation or Sprays:**

- A. If treated water is intended to be used only for crops or non-crop areas that are labeled for direct treatment with 2,4-D such as Pastures, Turfgrass or Cereal grains, the treated water may be used to irrigate and/or mix sprays for these sites at any time after the 2,4-D aquatic application.
- B. Due to potential phytotoxicity considerations, the following restrictions are applicable: If treated water is intended to be used to irrigate or mix sprays for plants grown in commercial nurseries and greenhouses; and other plants or crops that are not labeled for direct treatment with 2,4-D, the water must not be used unless one of the following restrictions has been observed:
 - i. A setback distance from functional water intake(s) of > 600 feet was used for the application, or,
 - ii. A waiting period of 7 days from the time of application has elapsed, or,
 - iii. An approved assay indicates that the 2,4-D concentration is 100 ppb (0.1 ppm) or less at the water intake. Wait at least 3 days after application before initial sampling at water intake.

2. Drinking Water (Potable Water):

- A. Consult with appropriate state or local water authorities before applying this product to public waters. State or local agencies may require permits. The potable water use restrictions on this label are to ensure that consumption of water by the public is allowed only when the concentration of 2,4-D in the water is less than the MCL (Maximum Contaminant Level) of 70 ppb. Applicators should consider the unique characteristics of the treated waters to assure that 2,4-D concentrations in potable water do not exceed 70 ppb at the time of consumption.
- B. For floating and emergent weed applications, the drinking water setback distance from functioning potable water intakes is > 600 feet.
- C. If no setback distance of > 600 feet is used for the application, applicators or the authorizing organization must provide a drinking water notification prior to a 2,4-D application to the party responsible for a public water supply or to individual private water users. Notification to the party responsible for a public water supply or to individual private water users must be done in a manner to assure that the party is aware of a water use restrictions when this product is applied to potable water.

The following is an example of notification via posting, but other methods of notification which convey the above restrictions may be used and may be required in some cases under state or local law or as a condition of a permit.

EXAMPLE:

Posting notification should be located every 250 feet including the shoreline of the treated area and up to 250 feet of shoreline past the application site to include immediate public access points.

Posting must include the day and time of application. Posting may be removed if analysis of a sample collected at the intake 3 days or more following application shows that the concentration in the water is less than 70 ppb (100 ppb for irrigation or sprays), or after 7 days following application, whichever occurs first.

Text of Notification: Wait 7 days before diverting functioning surface water intakes from the treated aquatic site to use as drinking water, irrigation, or sprays, unless water at functioning drinking water intakes is tested at least 3 days after application and is demonstrated by assay to contain no more than 70 ppb 2,4-D (100 ppb for irrigation or sprays).

Application Date: _____ Time: _____

- D. Following each application of this product, treated water must not be used for drinking water unless one of the following restrictions has been observed:
- i. A setback distance from functional water intake(s) of ≥ 600 feet was used for the application, or,
 - ii. A waiting period of at least 7 days from the time of application has elapsed, or,
 - iii. An approved assay indicates that the 2,4-D concentration is 70 ppb (0.07 ppm) or less at the water intake. Sampling for drinking water analysis should occur no sooner than 3 days after 2,4-D application. Analysis of samples must be completed by a laboratory that is certified under the Safe Drinking Water Act to perform drinking water analysis using a currently approved version of analytical Method Number 515, 555, other methods for 2,4-D as may be listed in Title 40 CFR Part 141.24, or Method Number 4015 (immunoassay of 2,4-D) from U.S. EPA Test Methods for Evaluating Solid Waste SW-846.
- E. **Note:** Existing potable water intakes that are no longer in use, such as those replaced by a connection to a municipal water system or a potable water well, are not considered to be functioning potable water intakes.
- F. Drinking water setback distances do not apply to terrestrial applications of 2,4-D adjacent to water bodies with potable water intakes.

Target Weeds / Treatment Site	Application Rate per Acre	Specific Use Instructions
Submerged Aquatic Weeds including Eurasian Water Milfoil (<i>Myriophyllum spicatum</i>) in Ponds, Lakes, Reservoirs, Marshes, Bayous, Drainage ditches, Canals, Rivers and Streams that are Quiescent or Slow moving including Programs of the Tennessee Valley Authority	2.8 gals. (10.8 lbs. a.e)	<ul style="list-style-type: none"> • Application Timing: For best results, apply in Spring or early Summer when aquatic weeds appear. Check for weed growth in areas heavily infested the previous year. A second application may be needed when weeds show signs of recovery, but no later than mid-August in most areas. • Subsurface Application: Apply this product, undiluted, directly to the water through a boat mounted distribution system. Shoreline areas should be treated by subsurface injection application by boat to avoid aerial drift. • Surface Application: Use power operated boat mounted boom sprayer. If rate is less than 5 gallons per acre, dilute to a minimum spray volume of 5 gallons per surface acre. • Aerial Application: Use drift control spray equipment or thickening agents mixed with sprays to reduce drift. Apply through standard boom systems in a minimum spray volume of 5 gallons per surface acre. For Microfoil drift control spray systems, apply this product in a total spray volume of 12 to 15 gallons per acre. • Apply to attain a concentration of 2 to 4 ppm (see "Table 1" below).

Table 1. Amount of 2,4-D to Apply for a Target Subsurface Concentration

Surface Area	Average Depth (ft.)	For Typical Conditions - 2 ppm 2,4-D lbs. a.e./Acre-foot	For Difficult Conditions* - 4 ppm 2,4-D lbs. a.e./Acre-foot
1 acre	1	5.4	10.8
	2	10.8	21.6
	3	16.2	32.4
	4	21.6	43.2
	5	27.0	54
*Examples include spot treatment of pioneer colonies of Eurasian Water milfoil and certain difficult to control aquatic species.			

Use Restrictions:

- Maximum application rate is 2.8 gallons of this product (10.8 lbs. a.e.) per acre-foot per application.
- Do not make more than 2 applications per season.
- Do not apply within 21 days of previous application.
- When treating moving bodies of water, applications must be made while traveling upstream to prevent concentration of 2,4-D downstream from the application.

Fish breathe oxygen in the water and a water-oxygen ratio must be maintained. Decaying weeds use up oxygen, but during the period when applications should be made, the weed mass is fairly sparse and the weed decomposition rate is slow enough that the water-oxygen ratio is not disturbed by treating the entire area at one time. If treatments must be applied later in the season when the weed mass is dense and repeat treatments are needed, apply product in lanes, leaving buffer strips which can then be treated when vegetation in treated lanes has disintegrated. During the growing season, weeds decompose in a 2 to 3 week period following treatment.

Water Use:

1. Water for Irrigation or Sprays

- A. If treated water is intended to be used only for crops or non-crop areas that are labeled for direct treatment with 2,4-D such as Pastures, Turfgrass or Cereal grains, the treated water may be used to irrigate and/or mix sprays for these sites at anytime after the 2,4-D aquatic application.
- B. Due to potential phytotoxicity and/or residue considerations, the following restrictions are applicable:
If treated water is intended to be used to irrigate or mix sprays for unlabeled crops, non-crop areas or other plants not labeled for direct treatment with 2,4-D, the water must not be used unless one of the following restrictions has been observed:
 - i. A setback distance described in the Drinking Water Setback Table was used for the application, or,
 - ii. A waiting period of 21 days from the time of application has elapsed, or,
 - iii. An approved assay indicates that the 2,4-D concentration is 100 ppb (0.1 ppm) or less at the water intake. See "Table 3" for the waiting period after application but before taking the initial sampling at water intake.

2. Drinking Water (Potable Water):

- A. Consult with appropriate state or local water authorities before applying this product to public waters. State or local agencies may require permits. The potable water use restrictions on this label are to ensure that consumption of water by the public is allowed only when the concentration of 2,4-D in the water is less than the MCL (Maximum Contaminant Level) of 70 ppb. Applicators should consider the unique characteristics of the treated waters to assure that 2,4-D concentrations in potable water do not exceed 70 ppb at the time of consumption.
- B. For submersed weed applications, the drinking water setback distances from functioning potable water intakes are provided in "Table 2 Drinking Water Setback Distance" (below).
- C. If no setback distance from the Drinking Water Setback Table (Table 2) is to be used for the application, applicators or the authorizing organization must provide a drinking water notification and an advisory to shut off all potable water intakes prior to a 2,4-D application.
Notification to the party responsible for a public water supply or to individual private water users must be done in a manner to assure that the party is aware of the water use restrictions when this product is applied to potable water.
The following is an example of notification via posting, but other methods of notification which convey the above restrictions may be used and may be required in some cases under state or local law or as a condition of a permit.

EXAMPLE:

Posting notification should be located every 250 feet including the shoreline of the treated area and up to 250 feet of shoreline past the application site to include immediate public access points.

Posting should include the day and time of application. Posting may be removed if analysis of a sample collected at the intake no sooner than stated in Table 3 (below) shows that the concentration in the water is less than 70 ppb (100 ppb for irrigation or sprays), or after 21 days following application, whichever occurs first.

Text of Notification: Wait 21 days before diverting functioning surface water intakes from the treated aquatic site to use as drinking water, irrigation, or sprays, unless water at functioning drinking water intakes is tested no sooner than (insert days from Table 3) and is demonstrated by assay to contain no more than 70 ppb 2,4-D (100 ppb for irrigation or sprays).

Application Date: _____ Time: _____

- D. Following each application of this product, treated water must not be used for drinking water unless one of the following restrictions has been observed:
 - i. A setback distance described in the Drinking Water Setback Distance Table was used for the application, or,
 - ii. A waiting period of at least 21 days from the time of application has elapsed, or,
 - iii. An approved assay indicates that the 2,4-D concentration is 70 ppb (0.07 ppm) or less at the water intake. Sampling for drinking water analysis should occur no sooner than stated in Table 3. Analysis of samples must be completed by a laboratory that is certified under The Safe Drinking Water Act to perform drinking water analysis using a currently approved version of analytical Method Number 515, 555, other methods for 2,4-D as may be listed in Title 40 CFR, Part 141.24, or Method Number 4015 (immunoassay of 2,4-D) from U.S. EPA Test Methods for Evaluating Solid Waste SW-846.

- E. **Note:** Existing potable water intakes that are no longer in use, such as those replaced by a connection to a municipal water system or a potable water well, are not considered to be functioning potable water intakes.
- F. Drinking water setback distances do not apply to terrestrial applications of 2,4-D adjacent to water bodies with potable water intakes.

Table 2. Drinking Water Setback Distance for Submersed Weed Applications

Application Rate and Minimum Setback Distance (feet) From Functioning Potable Water

1 ppm*	2 ppm*	3 ppm*	4 ppm*
600	1200	1800	2400

*ppm acid equivalent target water concentration

Table 3. Sampling for Drinking Water Analysis After 2,4-D Application for Submersed Weed Applications

Minimum Days After Application Before Initial Water Sampling at the Functioning Potable Water Intake

1 ppm*	2 ppm*	3 ppm*	4 ppm*
5	10	10	14

*ppm acid equivalent target water concentration

STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed by storage or disposal.

PESTICIDE STORAGE: Keep container tightly closed when not in use. If exposed to subfreezing temperatures, the product should be warmed up to at least 40°F and mixed thoroughly before using.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

Nonrefillable Container (rigid material; ≤ 5 gallons): Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container one-fourth full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Nonrefillable Container (rigid material; > 5 gallons up to < 250 gallons): Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container one-fourth full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Refillable Container (≥ 250 gallons & Bulk): Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

WARRANTY – CONDITIONS OF SALE

Our DIRECTIONS FOR USE for use of this product are based upon tests believed reliable. Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically directed, and other influencing factors in the use of this product are beyond the control of the Seller. To the extent consistent with applicable law, Buyer assumes all risks of use, storage and handling of this material not in strict accordance with the directions given herewith. To the extent consistent with applicable law, in no case shall the Manufacturer or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product when such use and/or handling is not in strict accordance with the directions given herewith. The foregoing is a condition of sale by the Seller and is accepted as such by the Buyer.



DE-AMINE, MIX and the Drexel logo are either trademarks or registered trademarks of Drexel Chemical Company. All other brand names, product names, or trademarks belong to their respective holders.

[EDITORIAL NOTES]

The following unit of measures found in the Tables under the “*DIRECTIONS FOR USE*” section can be formatted using the following abbreviations:

- gallon(s) - gal(s).
- gallons per acre – gals./Ac.
- pint(s) – pt(s).
- pint(s) per acre – pt(s)/Ac.
- pints per 100 gallons – pts./100 gals.
- fluid ounces – fl. ozs.
- fluid ounces per gallon – fl. ozs./gal.
- per - /
- square feet – sq. ft.
- Rate Per Acre – Rate/Ac.
- Acre-foot – Ac.-Ft.

ACCEPTED

07/31/2020

Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered under
EPA Reg. No. 19713-650

SUPPLEMENTAL LABEL

2,4-D GROUP 4 HERBICIDE

Drexel

De-Amine® 4

(EPA Reg. No. 19713-650)

**FOR USE IN BLUEBERRIES (High Bush Berries – Vegetative Strips Between Rows)
(MA, NJ, OR, WA & WI Only)**

**THIS SUPPLEMENTAL LABEL EXPIRES ON JULY 31, 2022 AND MUST NOT BE USED OR
DISTRIBUTED AFTER THIS DATE.**

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the entire Drexel De-Amine 4 label before proceeding with the use directions contained in this supplemental labeling.
- Follow all applicable directions, restrictions, Worker Protection Standard (WPS) requirements, and precautions on the registered label for EPA Reg. No. 19713-650.

Time of Application	This Product (Pt./Ac.)	Specific Use Instructions
First Application: Spring If necessary, second application: After Harvest	2.9	High Bush Berries (Vegetative Strips between Rows) Broadleaf weeds (Broadleaf dock, Canada thistle, Dandelion, Spotted catsear): Apply 2.9 pts. of this product in 50 gallons of water per acre using ground equipment only. Make the first application in the Spring as a direct, shielded spray to the vegetative strip between Blueberry rows, avoiding contact with the Blueberry plant foliage. If necessary, a second application at 3 pts. of this product in 50 gallons of water per acre may be made after harvest to control regrowth of broadleaf weeds.

Use Precautions:

- INJURY TO CROPS FROM THIS HERBICIDE MAY OCCUR. IF YOU ARE NOT PREPARED TO ACCEPT SOME DEGREE OF CROP INJURY DO NOT USE THIS PRODUCT ON BLUEBERRIES.
- For optimum herbicide performance, mowing between rows for at least 7 days before or after the application is not advised.
- To avoid injury to Blueberry plants, apply when temperature is less than 65°.
- Avoid drip or contact of this product to Blueberry plants in the growing or dormant period as it may kill or cause significant plant injury that could result in grade or yield loss.
- Soil residue of this product may temporarily inhibit seed germination and plant growth.

Use Restrictions:

- The pre-harvest interval (PHI) is 30 days.
- Limited to 2 application per year.
- Do not exceed 2.9 pints (1.4 lbs. a.e.) per acre per application.
- Do not exceed 5.8 pints (2.8 lbs. a.e.) per acre per year.
- When tank mixing with products that contain 2,4-D, do not exceed a combined total of 2.8 pounds of a.e. per acre per year.
- Do not apply through any type of irrigation system.
- Do not apply by air.
- Do not apply in or near greenhouses.



DE-AMINE and the Drexel logo are registered trademarks of Drexel Chemical Company.

ACCEPTED

07/31/2020

Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered under
EPA Reg. No. 19713-650

SUPPLEMENTAL LABEL

2,4-D GROUP 4 HERBICIDE

Drexel

De-Amine® 4

(EPA Reg. No. 19713-650)

FOR USE IN CRANBERRIES
(MA, NJ, OR, WA & WI Only)

THIS SUPPLEMENTAL LABEL EXPIRES ON JULY 31, 2022 AND MUST NOT BE USED OR DISTRIBUTED AFTER THIS DATE.

KEEP OUT OF REACH OF CHILDREN
CAUTION

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the entire Drexel De-Amine 4 label before proceeding with the use directions contained in this supplemental labeling.
- Follow all applicable directions, restrictions, Worker Protection Standard (WPS) requirements, and precautions on the registered label for EPA Reg. No. 19713-650.

Time of Application	This Product (Pt./Ac.)	Specific Use Instructions
Late June and July (for best results)	2.4	To control tall weeds in Cranberry bogs, apply with a wooden frame or similar device shaped like a hockey stick with its lower member wrapped with several thicknesses of Turkish toweling (or other suitable material). Apply by soaking the toweling in one part of this product to two parts water. Then, with swabbed portion of the stick horizontal, wave left and right above the cranberry vines, wiping small quantities of the herbicide onto tall weeds above the crop level.

Use Precautions:

- INJURY TO CROPS FROM THIS HERBICIDE MAY OCCUR. IF YOU ARE NOT PREPARED TO ACCEPT SOME DEGREE OF CROP INJURY, DO NOT USE THIS PRODUCT ON CRANBERRIES.
- Avoid drip or contact of this product to Cranberry vines as it may kill or cause significant injury to Cranberry.

Use Restrictions:

- The pre-harvest interval (PHI) is 30 days.
- Limited to 2 applications per year.
- Do not exceed 2.5 pints (1.2 lbs. a.e.) per acre per application.
- When tank-mixing with products that contain 2,4-D, do not exceed a combined total of 1.2 pounds of a.e. per acre per year.
- Do not apply through any type of irrigation system.
- Do not apply by air.

Manufactured By:

Drexel Chemical Company
P.O. BOX 13327 MEMPHIS, TN 38113-0327
EST. 1972

DE-AMINE and the Drexel logo are registered trademarks of Drexel Chemical Company.

ACCEPTED

07/31/2020

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 19713-650

SUPPLEMENTAL LABEL

2,4-D GROUP 4 HERBICIDE

Drexel

De-Amine® 4

(EPA Reg. No. 19713-650)

**FOR USE IN GRAPE VINEYARDS
(CALIFORNIA Only)**

THIS SUPPLEMENTAL LABEL EXPIRES ON JULY 31, 2022 AND MUST NOT BE USED OR DISTRIBUTED AFTER THIS DATE.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the entire Drexel De-Amine 4 label before proceeding with the use directions contained in this supplemental labeling.
- Follow all applicable directions, restrictions, Worker Protection Standard (WPS) requirements, and precautions on the registered label for EPA Reg. No. 19713-650.

Time of Application	This Product (Pt./Ac.)	Specific Use Instructions
After shatter following bloom and before Grape shoots reach the ground or during dormant season	1.8 to 2.8	Use this product on Grape vineyards established for at least 3 years to control Canada thistle, Field bindweed (Morningglory) and other 2,4-D susceptible broadleaf weeds when weeds are in the bud to early bloom stage and are growing vigorously. Dilute in 10 to 100 gallons of water to treat 1 acre of ground to be sprayed. For band or spot treatment, calculate rates according to the actual portion of acre treated. Use a hooded boom and low pressure flooding nozzles to deliver coarse droplets.

Use Precautions:

- Grapes are extremely sensitive to 2,4-D. Use directed application so no 2,4-D contacts Grape leaves and young shoots or stems.

Use Restrictions:

- The pre-harvest interval (PHI) is 100 days.
- Limited to 1 application per crop cycle.
- Maximum of 2.8 pints (1.36 lbs. a.e.) per acre per application.
- When tank-mixing with products that contain 2,4-D, do not exceed a combined total of 1.36 pounds of a.e. per acre per year.

Manufactured By:

Drexel Chemical Company
P.O. BOX 13327, MEMPHIS, TN 38113-0327
EST. 1972

DE-AMINE and the Drexel logo are registered trademarks of Drexel Chemical Company.