



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

19713-735

Date of Issuance:

6/12/2023

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

Drexel Imazapyr 4SL

Name and Address of Registrant (include ZIP Code):

Drexel Chemical Company
P.O.BOX. 13327
Memphis, TN 38113-0327

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above-named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.
2. **The data requirements for storage stability and corrosion characteristics (Guidelines 830.6317 and 830.6320) are not satisfied. A one-year study is required to satisfy these data requirements. You have 18 months from the date of registration to provide these data.**

Signature of Approving Official:

Heather McFarley, Acting Product Manager 24
Fungicide and Herbicide Branch, Registration Division (7505P)
Office of Pesticide Programs

Date:

6/12/2023

3. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, “EPA Reg. No. 19713-735.”
4. Submit one copy of the revised final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 7/27/2022

If you have any questions, please contact Sayed Islam by phone at 202-566-2796, or via email at islam.sayed@epa.gov

Enclosure:

- Accepted label

ACCEPTED

06/12/2023

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 19713-735

IMAZAPYR GROUP 2 HERBICIDE

Drexel

IMAZAPYR 4SL

For control of undesirable vegetation growing within certain Aquatic sites, Forestry sites, Pasture/Rangeland, Non-agricultural lands, establishment and maintenance of Wildlife openings, release of unimproved Bermudagrass and Bahiagrass, bareground weed control, for use under certain paved areas, Industrial non-cropland areas including Highway, Pipeline, Railroad and Utility rights-of-way, Utility plant sites, Petroleum tank farms, Pumping installations, Fence rows, Storage areas, Non-irrigation ditchbanks including grazed or hayed areas within these sites, roads and transmission lines.

ACTIVE INGREDIENT:

Isopropylamine salt of Imazapyr: (2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-3-pyridinecarboxylic acid)*... 53.10%

OTHER INGREDIENTS: 46.90% TOTAL: 100.00%

* Equivalent to 43.3% 2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-3-pyridinecarboxylic acid or 4 pounds acid per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

[See FIRST AID Below]

[See Side (Back) Panel for FIRST AID];

[See Page ___ for FIRST AID]

[See Container Labeling for (FIRST AID and Complete Directions for Use]

[See (Attached) Booklet (Container Labeling) for Complete Directions for Use]

EPA Reg. No. 19713-XX

Net Content:

EPA Est. No. 19713-XX-X

Gals. (___) L

FIRST AID

IF INHALED:

- Move person to fresh air.
• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.
• Call a poison control center or doctor for further treatment advice.

IF IN EYES:

- Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.
• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
• Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
• Rinse skin immediately with plenty of water for 15 to 20 minutes.
• Call a poison control center or doctor for treatment advice.

IF SWALLOWED:

- Call a poison control center or doctor immediately for treatment advice.
• Have person sip a glass of water if able to swallow.
• DO NOT induce vomiting unless told to do so by a poison control center or doctor.
• DO NOT give anything by mouth to an unconscious person.

(Continued)

FIRST AID (Cont.)

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also call CHEMTREC at 800-424-9300 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Harmful if swallowed, absorbed through skin or inhaled. Avoid contact with eyes, skin or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants,
• Chemical-resistant gloves made of any waterproof material including polyethylene or polyvinyl chloride, and
• Shoes plus socks.

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. DO NOT reuse them.

ENGINEERING CONTROLS

Pilots must use an enclosed cockpit that meet the requirements listed in the Worker Protection Standard (WPS) for Agricultural Pesticides [40 CFR 170.240(d) (6)].

USER SAFETY RECOMMENDATIONS

Users should: 1) Wash hands thoroughly before eating, drinking, chewing gum, using tobacco or using the toilet. 2) Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. 3) Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to plants. Drift and runoff may be hazardous to plants in water adjacent to treated areas. DO NOT apply directly to water except as specified on the label. Treatment of aquatic weeds may result in oxygen depletion or loss due to decomposition of dead plants. DO NOT treat more than one half the surface area of the water in a single operation and wait at least 10 to 14 days between treatments. Begin treatment along the shore and proceed outward in bands to allow aquatic organisms to move into untreated areas. DO NOT contaminate water when disposing of equipment washwater or rinsate. This pesticide is toxic to vascular plants and must be used strictly in accordance with the drift precautions of the label.



Manufactured By: Drexel Chemical Company P.O. Box 13327, Memphis, TN 38113-0327

SINCE 1972

THE DREXEL logo is a registered trademark of Drexel Chemical Company.

IMAZAPYR 2SL DRAFTSP-0722 IMAZAPYR 4SL Page 1 of 15

PHYSICAL AND CHEMICAL HAZARDS

Spray solutions of this product must be mixed, stored and applied only in stainless steel, fiberglass, plastic and plastic-lined steel containers. **DO NOT** mix, store or apply this product or spray solutions of this product in unlined steel (except stainless steel) containers or spray tanks.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the Restricted Entry Interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil, or water, is:

- Coveralls
- Shoes plus socks
- Chemical-resistant gloves made of any waterproof material
- Protective eyewear

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard (WPS) for Agricultural Pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

DO NOT enter or allow others to enter treated areas until sprays have dried.

PRODUCT INFORMATION

This product is an aqueous solution to be mixed with water and a surfactant and applied as a spray solution to control undesirable vegetation growing within certain Aquatic sites, Forestry sites, Pasture/Rangeland, and Non-agricultural lands. Aquatic sites consist of standing and flowing water, Estuarine/Marine, Wetland, and Riparian areas. Non-agricultural lands include private, public and military land as follows: uncultivated non-agricultural areas (including Airports, Highway, Railroad and Utility rights-of-way and Sewage disposal areas), uncultivated agricultural areas – non-crop producing (including Farmyards, Fence rows, Fuel storage areas, Non-irrigation ditch banks and barrier strips), industrial sites – outdoor (including Lumber yards, Pipeline and Tank farms) and natural areas (including Wildlife management areas, Wildlife openings, Wildlife habitats, Recreation areas, Campgrounds, Trailheads and Trails). This product may also be used for the release of unimproved Bermudagrass and Bahiagrass, for bareground weed control, and for use under certain paved surfaces.

Herbicidal Activity: This product will control most annual and perennial grasses and broadleaf weeds in addition to many brush and vine species with some residual control of undesirable species that germinate above the waterline. This product is readily absorbed through emergent leaves and stems and is translocated rapidly throughout the plant, with accumulation in the meristematic regions. For maximum activity, weeds must be growing robustly at the time of application, and the spray solution must include a surfactant (see "ADJUVANTS" section for specific use directions).

Treated plants stop growing soon after spray application. Chlorosis appears first in the newest leaves, and necrosis spreads from this point. In perennials, the herbicide is translocated into, and kills, underground or submerged storage organs, which prevents regrowth.

Chlorosis and tissue necrosis may not be apparent in some plant species until two or more weeks after application. Complete kill of plants may not occur for several weeks. Applications of this product are rainfast one hour after treatment.

RESTRICTIONS

Maximum Rate – Annual

- **DO NOT** apply more than 48 fluid ounces of this product (1.5 lb. (ae) Imazapyr) per acre per year.

Maximum Rate – Single Application

- **DO NOT** apply more than 48 fluid ounces of this product (1.5 lb. (ae) Imazapyr) per single application.

Maximum Number of Applications Per Year

- **DO NOT** apply more than 1 application per year.

DO NOT use on food or feed crops.

DO NOT apply this product to water within 0.5 mile upstream of an active potable water intake in flowing water (i.e. river, stream, etc.) or within 0.5 mile of an active potable water intake in a standing body of water, including a lake, pond or reservoir.

DO NOT apply to water used for irrigation except as described in "USE PRECAUTIONS AND RESTRICTIONS" section of this label.

Keep from contact with fertilizers, insecticides, fungicides and seeds.

DO NOT drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the treated soil may be washed or moved into contact with their roots.

DO NOT use on lawns, walks, driveways, tennis courts.

DO NOT side trim desirable vegetation with this product unless severe injury and plant death can be tolerated. Prevent drift of spray to desirable plants.

If treated vegetation is to be removed from the application site, **DO NOT** use the vegetative matter as mulch or compost on or around desirable species.

Clean application equipment after using this product by thoroughly flushing with water.

Non-agricultural Lands and Forestry Sites

- **DO NOT** apply more than 48 fluid ounces of this product (1.5 lb. (ae) Imazapyr) per acre per year.
- **DO NOT** apply more than 48 fluid ounces of this product (1.5 lb. (ae) Imazapyr) per single application.
- **DO NOT** apply more than 1 application per year.

Pasture/Rangeland Sites

- **DO NOT** apply more than 24 fluid ounces of this product (0.75 lb. (ae) Imazapyr) per acre per year.
- **DO NOT** apply more than 24 fluid ounces of this product (0.75 lb. (ae) Imazapyr) per single application.
- **DO NOT** apply more than 1 application per year.
- **DO NOT** treat more than one-tenth of the available area to be grazed or cut for hay.
- For spot treatment only.

Aquatic Sites

- **DO NOT** apply more than 48 fluid ounces of this product (1.5 lb. (ae) Imazapyr) per acre per year.
- **DO NOT** apply more than 48 fluid ounces of this product (1.5 lb. (ae) Imazapyr) per single application.
- **DO NOT** apply more than 1 application per year.
- **DO NOT apply to aquatic uses in New York State.**

Aerial application - Aerial application to Aquatic sites is restricted to helicopter only.

Irrigation water - Application to water used for irrigation that results in residues greater than 1.0 part per billion (ppb) **MUST NOT** be used for irrigation purposes for 120 days after application or until residue levels of this product are determined by laboratory analysis or other appropriate means of analysis to be 1.0 ppb or less. When applications are made within 500 feet of an active irrigation intake, **DO NOT** irrigate for at least 24 hours following application to allow for dissipation.

Quiescent or Slow-moving Waters - In lakes and reservoirs, **DO NOT** apply this product within 1 mile of an active irrigation water intake during the irrigation season. Applications less than 1 mile from an active irrigation water intake may be made during the off-season, provided that the irrigation intake will remain inactive for a minimum of 120 days after application or until residue levels of this product are determined by laboratory analysis or other appropriate means of analysis to be 1.0 ppb or less.

Restrictions for potable water intakes - **DO NOT** apply this product directly to water within 0.5 mile upstream of an active potable water intake in flowing water (i.e. river, stream, etc.) or within 0.5 mile of an active potable water intake in a standing body of water including

a lake, pond or reservoir. To make Aquatic applications around and within 0.5 miles of active potable water intakes, the water intake must be turned off during application and for a minimum of 48 hours after the application. These Aquatic applications may be made only in the cases where there are alternative water sources or holding ponds that would permit the turning off of an active potable water intake for a minimum period of 48 hours after the applications.

NOTE: Existing potable water intakes that are no longer in use, including those replaced by connections to wells or a municipal water system, are not considered to be active potable water intakes. This restriction does not apply to intermittent, inadvertent overspray of water in terrestrial use sites.

Permitting - Consult local state fish and game agency and water control authorities before applying this product to public water. Permits may be required to treat such water.

Public waters - Application of this product to water can only be made by federal or state agencies, including Water Management District personnel, municipal officials, and the U.S. Army Corps of Engineers, or those applicators who are licensed or certified as aquatic pest control applicators and are authorized by the state or local government. Treatment to other than non-native invasive species is limited to only those plants that have been determined to be a nuisance by a federal or state government entity.

Private waters - Applications may be made to private waters that are still, including ponds, lakes and drainage ditches where there is minimal or no outflow to public waters.

Recreational use of water in treatment area - There are no restrictions on the use of water in the treatment area for recreational purposes, including swimming and fishing.

Livestock use of water in/from treatment area - There are no restrictions on livestock consumption of water from the treatment area.

Precautions for Avoiding Injury to Non-target Plants

Untreated desirable plants can be affected by root uptake of this product from treated soil. Injury or loss of desirable plants may result if this product is applied on or near desirable plants, on areas where their roots extend, or in locations where the treated soil may be washed or moved into contact with their roots. When making applications along shorelines where desirable plants may be present, caution must be exercised to avoid spray contact with their foliage or spray application to the soil in which they are rooted. Shoreline plants that have roots that extend into the water in an area where this product has been applied generally will not be adversely affected by uptake of the herbicide from the water.

MANDATORY SPRAY DRIFT

Aerial Applications:

- **DO NOT** release spray at a height greater than 10 ft. above the vegetative canopy, unless a greater application height is necessary for pilot safety.
- Applicators are required to use a coarse or coarser spray droplet size (ASABE S572.1) for all applications.
- The boom length must not exceed 65% of the wingspan for airplanes or 75% of the rotor blade diameter for helicopters.
- Applicators must use one-half swath displacement upwind at the downwind edge of the field.
- Nozzles must be oriented so the spray is directed toward the back of the aircraft.
- **DO NOT** apply when wind speeds exceed 10 mph at the application site.
- **DO NOT** apply during temperature inversions.

Ground Applications:

- Apply with the nozzle height specified by the manufacturer, but no more than 3 ft. above the ground or crop canopy.
- Applicators are required to use a coarse or coarser spray droplet size (ASABE S572.1) for all applications.
- **DO NOT** apply when wind speeds exceed 10 mph at the application site.
- **DO NOT** apply during temperature inversions.

Boomless Ground Applications:

- Applicators are required to use a coarse or coarser droplet size (ASABE S572.1) for all applications.
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

SPRAY DRIFT ADVISORIES

Boomless Ground Applications:

- Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

Handheld Technology Applications:

- Take precautions to minimize spray drift.
- THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- Volume – Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure – Use the lowest spray pressure specified for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle – Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size – Aircraft

- Adjust Nozzles – Follow nozzle manufacturers directions for setting up nozzles. Generally, to reduce fine droplets, nozzles must be oriented parallel with the airflow in flight.

Boom Height – Ground Boom

Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom must remain level with the crop and have minimal bounce.

Release Height – Aircraft

Higher release heights increase the potential for spray drift. When applying aerially to crops, **DO NOT** release spray at a height greater than 10 feet above the crop canopy, unless a greater application height is necessary for pilot safety.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

WIND EROSION

Avoid treating powdery dry or light sandy soils when conditions are favorable for wind erosion. Under these conditions, the soil surface must first be settled by rainfall or irrigation.

WEED RESISTANCE MANAGEMENT

IMAZAPYR GROUP 2 HERBICIDE

For resistance management, this product is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to this product and other Group 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same area. Appropriate resistance management strategies must be followed.

To delay herbicide resistance take one or more of the following steps:

- Rotate the use of this product or other Group 2 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds.

- Use tank-mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use [and crop rotation] and that considers mechanical control methods, cultural (e.g., timing to favor the Turf [or crop] [higher crop seeding rates; precision fertilizer application method] and not the weeds), biological (weed competitive [crops] [or] varieties) and other management practices.
- Scout before and after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method [including hoeing or tillage]. Prevent movement of resistant weed seeds to other areas by cleaning equipment [when moving between fields, and planting clean seed].
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management directions for specific crops and weed biotypes.
- For further information or to report non-performance or suspected resistance, contact Drexel Chemical Company representatives at (901) 774-4370.

It is advisable to keep accurate records of pesticides applied to individual fields to help obtain information on the spread and dispersal of resistant biotypes. Contact your local sales representative, crop advisor, or extension agent to find out if suspected resistant weeds to this MOA have been found in your region. **DO NOT** assume that each listed weed is being controlled by this mechanism of action. Co-formulated active ingredients are intended to broaden the spectrum of weeds that are controlled. Some weeds may be controlled by only one of the active ingredients in this product.

Suspected herbicide-resistant weeds may be identified by these indicators:

- Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds.
- A spreading patch of non-controlled plants of a particular weed species; and
- Surviving plants mixed with controlled individuals of the same species.

INTEGRATED PEST MANAGEMENT

Drexel advises the use of Integrated Pest Management (IPM) programs to control pests. This product may be used as part of an Integrated Pest Management (IPM) program which can include biological, cultural, and genetic practices including field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop or site systems in your area.

ADJUVANTS

Post-emergence applications of this product require the addition of a spray adjuvant for optimum herbicide performance. Only spray adjuvants that are approved or appropriate for aquatic use can be utilized.

Non-ionic Surfactants: Use a non-ionic surfactant at the rate 0.25% v/v or higher (see manufacturer's label) of the spray solution (0.25% v/v is equivalent to 1 qt. in 100 gal.). For best results, select a non-ionic surfactant with a HLB (Hydrophilic to Lipophilic Balance) ratio between 12 and 17 with at least 70% surfactant in the formulated product. Alcohols, fatty acids, oils, ethylene glycol or diethylene glycol must not be considered as surfactants to meet the above requirements.

Methylated Seed Oils or Vegetable Oil Concentrates: Instead of a surfactant, a methylated seed oil or vegetable-based seed oil concentrate may be used at the rate of 24 to 32 fluid ounces per acre. When using spray volumes greater than 30 gallons per acre, methylated seed oil or vegetable based seed oil concentrates must be mixed at a rate of 1 % of the total spray volume, or alternatively use a non-ionic surfactant as described above. Research indicates that these oils may aid in product deposition and uptake by plants under moisture or temperature stress.

Silicone Based Surfactants: See manufacturer's label for specific rate recommendations. Silicone-based surfactants may reduce the surface tension of the spray droplet, allowing greater spreading on the leaf surface as compared to conventional non-ionic surfactants. However, some silicone-based surfactants may dry too quickly, limiting herbicide uptake.

Invert Emulsions: This product can be applied as an invert emulsion. The spray solution results in an invert (water-in-oil) spray emulsion designed to minimize spray drift and spray runoff, resulting in more herbicide on the target foliage. The spray emulsion may be formed in a single tank (batch mixing) or injected (in-line mixing). Consult the invert chemical label for proper mixing directions. **DO NOT** apply more than 48 fluid ounces of this product per acre (1.5 lb. (ae) Imazapyr) in an invert emulsion.

Fertilizer/Surfactant Blends: Nitrogen based liquid fertilizers including 28%N, 32%N, 10-34-0 or ammonium sulfate, may be added at the rate of 32 to 48 fluid ounces per acre in combination with the specified rate of non-ionic surfactant, methylated seed oil or vegetable/seed oil concentrate. The use of fertilizers in a tank-mix without a non-ionic surfactant, methylated seed oil or vegetable/seed oil concentrate is not advised.

Other: An antifoaming agent, spray pattern indicator or drift reducing agent may be applied at the product labeled rate if necessary or desired.

TANK-MIXES

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product label involved in tank-mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank-mixture.

APPLICATION METHODS

This product may be selectively applied by using low volume directed application techniques or may be broadcast applied using ground equipment, watercraft or aircraft. Aerial applications to Aquatic sites must be made by helicopter. In addition, this product may also be applied using cut stump, cut stem, and frill or girdle treatment techniques within Non-agricultural lands, Pasture/Rangeland and Aquatic sites. See "AERIAL APPLICATION" and "GROUND APPLICATION" sections for additional details.

COMPATIBILITY

Before full-scale mixing of this product with other pesticides, emulsifiers, fertilizers, surfactants or oils, determine the compatibility of the proposed mixture. Use proportionate quantities of each ingredient and mix in a small container. Always mix one product thoroughly with the diluent before adding another product. If no incompatibility is evident after 30 minutes, the mixture is generally compatible for spraying. To evaluate potential short term effects of applying the mixture, test the tank-mix combination on a few plants or a small area before larger-scale treatments. Wait at least 2 to 3 days for problems to become apparent.

IMPORTANT: MIXING WITH OTHER SUBSTANCES MAY INCREASE THE RISK OF MIXING INCOMPATIBILITIES, REDUCED EFFECTIVENESS AND/OR CAUSE CROP INJURY OR LOSS. ANY LIABILITY FOR LOSS, INJURY OR DAMAGE RESULTING FROM A MIXTURE NOT SPECIFIED ON THIS LABEL OR IN MANUFACTURER'S SUPPLEMENTAL LABELING DISTRIBUTED FOR THIS PRODUCT IS SPECIFICALLY DISCLAIMED BY MANUFACTURER.

AERIAL APPLICATION

All precautions must be taken to minimize or eliminate spray drift. Both helicopter and fixed wing aircraft can be used to apply this product, but applications to Aquatic sites are restricted to helicopter only. **DO NOT** make applications by helicopter or fixed wing aircraft unless appropriate buffer zones can be maintained to prevent spray drift out of the target area, or when spray drift as a result of helicopter application can be tolerated. Aerial equipment designed to minimize spray drift including a helicopter equipped with a Microfoil™ boom Thru-Valve™ boom or raindrop nozzles must be used and calibrated. Except when applying with a Microfoil boom, a drift control agent may be added at the specified label rate. **DO NOT** side trim with this product unless death of treated tree can be tolerated.

Uniformly apply the specified amount of this product in 2 to 30 gallons of water per acre. A foam reducing agent may be added at the specified label rate.

Immediately after each use of this product thoroughly clean application equipment, including landing gear. Uncoated steel surfaces (except stainless steel surfaces) may result in corrosion and failure after prolonged exposure to the product. The maintenance of any paint (organic coating) may prevent corrosion.

GROUND APPLICATION RESTRICTIONS

Maximum Rate – Annual

- **DO NOT** apply more than 48 fluid ounces of this product (1.5 lb. (ae) Imazapyr) per acre per year.

Maximum Rate – Single Application

- **DO NOT** apply more than 48 fluid ounces of this product (1.5 lb. (ae) Imazapyr) per single application.

Maximum Number of Applications Per Year

- **DO NOT** apply more than 1 application per year.

Low Volume Foliar:

Use equipment calibrated to deliver 5 to 20 gallons of spray solution per acre. To prepare the spray solution, thoroughly mix in water 0.25 to 2.5% of this product plus surfactant (see the "ADJUVANTS" section of this label for specific instructions). A foam reducing agent may be applied at the label rate, if needed. For control of difficult species (see "AQUATIC WEEDS CONTROLLED" section and the "TERRESTRIAL WEEDS CONTROLLED" section for relative susceptibility of weed species), use the higher concentrations of herbicide and/or spray volumes but **DO NOT** apply more than 48 fluid ounces of this product (1.5 lb. (ae) Imazapyr) per acre per year in Aquatic sites and Non-agricultural lands. **DO NOT** apply more than 24 fluid ounces of this product (0.75 lb. (ae) Imazapyr) per acre per year in Pasture/Rangeland sites.

Excessive wetting of foliage is not necessary.

For low volume foliar application, select proper nozzles to avoid over-application. Proper application is critical to ensure desirable results. Best results are achieved when the spray covers the crown and approximately 70% of the plant. The use of an even flat fan tip with a spray angle of 40 degrees or less will aid in proper deposition.

Appropriate tip sizes include 4004E, or 1504E. For a straight stream and cone pattern, adjustable cone nozzles including 5500 X3 or 5500 X4 may be used. Attaching a rollover valve onto a Spraying Systems Model 30 gunjet or other similar spray guns allows for the use of both a flat fan and cone tips on the same gun.

Moisten, but **DO NOT** drench target vegetation causing spray solution to runoff.

Low Volume Foliar with Backpacks:

For low-growing species, spray down on the crown, covering crown and penetrating approximately 70% of the plant.

For target species 4 to 8 feet tall, swipe the sides of target vegetation by directing spray to at least two sides of the plant in smooth vertical motions from the crown to the bottom. Make sure to cover the crown whenever possible.

For target species over 8 feet tall, lace sides of the target vegetation by directing spray to at least two sides of the target in smooth zigzag motions from crown to bottom.

Low Volume Foliar with Hydraulic Handgun Application Equipment:

Use same technique as described above for Low Volume Foliar with Backpacks.

For broadcast applications, simulate a gentle rain near the top of target vegetation, allowing spray to contact the crown and penetrate the target foliage without falling to the understory. Herbicide spray solution which contacts the understory may result in severe injury or death of plants in the understory.

High Volume Foliar:

For optimum performance when spraying medium to high-density vegetation, use equipment calibrated to deliver up to 100 gallons of spray solution per acre (GPA). Spray solutions exceeding 100 GPA may result in excessive spray runoff, causing increased ground cover injury, and injury to desirable species.

To prepare the spray solution, thoroughly mix this product in water and add a surfactant (see "ADJUVANT" section for specifications and rates of surfactants). A foam-reducing agent may be added at the label rate, if needed. For control of difficult species (see "AQUATIC WEEDS CONTROLLED" section and the "ADDITIONAL WEEDS CONTROLLED" section for relative susceptibility of weed species), use the higher concentrations of herbicide and/or spray volumes, but **DO NOT** apply more than 48 fluid ounces of this product (1.5 lb. (ae) Imazapyr) per acre per year in Aquatic sites and Non-agricultural lands. **DO NOT** apply more than 24 fluid ounces of this product (0.75 lb. (ae) Imazapyr) per acre per year in Pasture/Rangeland sites.

Uniformly cover the foliage of the vegetation to be controlled but **DO NOT** apply to run-off. Excessive wetting of foliage is not necessary.

SIDE TRIMMING

DO NOT side trim with this product unless severe injury or death of the treated tree can be tolerated. This product is readily translocated and can result in death of the entire tree.

CUT SURFACE TREATMENTS

This product may be used to control undesirable woody vegetation by applying the product solution to the cambium area of freshly cut stump surfaces or to fresh cuts on the stem of the target woody vegetation. Applications can be made at any time of the year except during periods of heavy sap flow in the Spring. **DO NOT** over apply solution causing runoff from the cut surface.

Injury may occur to desirable woody plants if the shoots extend from the same root system or their root systems are grafted to those of the treated tree.

This product may be mixed as either a concentrate or dilute solution for stump and cut stem treatments. The dilute solution may be used for applications to the surface of the stump or to cuts on the stem of the target woody vegetation. Concentrated solutions may be used for applications to cuts on the stem. Use of the concentrated solution permits application to fewer cuts on the stem, especially for large diameter trees. Follow the application instructions to determine proper application techniques for each type of solution.

- To prepare a dilute solution, mix 4 to 6 fluid ounces of this product (0.12 to 0.2 lb. (ae) Imazapyr) with one gallon of water. If temperatures are such that freezing of the spray mixture may occur, antifreeze (ethylene glycol) may be used according to manufacturer's label to prevent freezing. The use of a surfactant or penetrating agent may improve uptake through partially callused cambiums. To prepare a concentrated solution, mix 32 fluid ounces of this product (1 lb. (ae) Imazapyr) with no more than 32 fluid ounces of water.

CUT STUMP TREATMENT

Spray or brush the solution onto the cambium area of the freshly cut stump surface. Ensure that the solution thoroughly wets the entire cambium area (the wood next to the bark of the stump).

CUT STEM TREATMENT

(injection, hack-and-squirt)

Dilute Solution - Using standard injection equipment, apply 1 milliliter of solution at each injection site around the tree with no more than one-inch intervals between cut edges. Ensure that the injector completely penetrates the bark at each injection site.

Concentrate Solution - Using standard injection equipment, apply 1 milliliter of solution at each injection site. Make at least one injection cut for every 3 inches of Diameter at Breast Height (DBH) on the target tree. For example, a 3-inch DBH tree will receive 1 injection cut and a 6-inch DBH tree will receive 2 injection cuts. On trees requiring more than one injection site place the injection cuts at approximately equal intervals around the tree.

CUT STUBBLE

This product can be applied within 2 weeks after mechanical mowing or cutting of brush. To suppress or control resprouting, uniformly apply a spray solution of this product at the rate of 8 to 16 fluid ounces (0.25 to 0.5 lb. (ae) Imazapyr) per acre to the cut area. This product may be tank-mixed with Picloram, or equivalent labeled product for this use, to aid in control or suppression of brush. The addition of 5% (v/v) or more of a penetrating agent can aid in uptake through the bark or exposed roots.

Cut stubble applications are made to the soil and cut brush stumps. This type of application may increase ground cover injury. However, vegetation will recover. Making applications of this product directly to the soil can increase potential root uptake causing injury or death of desirable trees.

Efficacy can be increased and root uptake by desirable vegetation can be decreased if the brush is allowed to regrow and the foliage is treated. See the "BRUSH CONTROL" section of this label.

FRILL OR GIRDLE TREATMENT

Using a hatchet, machete, or chain saw, make cuts through the bark and completely around the tree to expose the cambium. The cut must angle downward extending into the cambium enough to expose at least two growth rings. Using a spray applicator or brush, apply a 12.5 to 50% solution of this product into each cut until thoroughly wet. Avoid applying so much herbicide that runoff to the ground or water occurs.

BASAL APPLICATION

This product is an aqueous formulation that requires mixing with **basal oil containing at least 15% emulsifier or will require the addition of an emulsifier, for application to the basal area** of brush and trees to control undesirable vegetation in the following non-cropland areas: Access roads, Airfields, Airports, along Forest roads, around Commercial or Industrial structures or outbuildings, around Farm and Ranch structures and outbuildings, Bare ground, Construction sites, Ditch banks, Dry ditches and canals, Fences and Fencerows, Firebreaks, Gravel yards, Habitat restoration and management areas, Highways and Roadsides (including Aprons, Medians, Guardrails and Right-of-ways), Industrial plant sites, Industrial areas, Lumber yards, Natural areas, Paved areas, Petroleum and other tank farms, Pumping installations, Pipeline, Power, Telephone and Utility rights-of-way, Power stations, Railroad rights-of-way, Refineries, Resorts, Storage areas, Substations, uncropped Farmstead areas, Vacant lots, Walkways, Wastelands and Wildlife habitat areas.

Thinline Basal and Stem Application

- This product may be applied as a thinline basal or arcing application to the stems of susceptible species including Big leaf maple (*Acer macrophyllum*), Willow (*Salix* spp.) and Eucalyptus (*Eucalyptus* spp.) with a stem ground line diameter of 3 inches or less. Mix 12 to 24 fluid ounces of this product in one gallon (0.4-0.75 lb. (ae) Imazapyr) of **basal oil containing at least 15% emulsifier**. Maintain uniform mixtures with frequent agitation. Direct a thin line of the spray solution to the stems beginning a few feet from the ground and descending toward the base of the tree making a zig-zag motion. **DO NOT** over apply causing puddling.

Low Volume Basal Bark Treatments

- This product, at the rate of 4 to 6 fluid ounces per gallon (0.12 to 0.20 lb. (ae) Imazapyr) may be applied for low volume basal bark treatments.
- This product at 1.5 to 2.5% (0.06 to 1.0 lb. (ae) Imazapyr) is advised to be tank-mixed with Triclopyr or other basal products to broaden the spectrum of control. Consult the herbicide labels for rates and susceptible brush species. Mixing with basal requires compatibility tests prior to mixing large quantities. Mixing aids (including emulsifiers, etc.) and ongoing agitation are required to attain a homogenous tank-mix.
- Basal application must be made to the lower 12" to 18" of the target brush and go to the soil. Care must be taken to not puddle or over treat the stem. Basal application is best suited for low density brush sites, where stems **DO NOT** exceed 700 stems per acre.

For Basal Application – It is advisory to mix only the intended amount of mixture that is to be sprayed that day. Adequate agitation must be maintained with all emulsion mixtures to prevent phase separation. Prior to tank-mixing with other products, herbicides and oils, you must determine the compatibility of the proposed mixture (See "COMPATIBILITY" section).

SPRAY SOLUTION MIXING GUIDE						
TANK-MIXING						
AMOUNT OF SPRAY SOLUTION BEING PREPARED	THIS PRODUCT ALONE		THIS PRODUCT WHEN TANK-MIXING		TRICLOPYR	
	4.0 Fl. Oz.	6.0 Fl. Oz..	1.5%	2.5%	15%	20%
1 Gal.	4.0 fl. oz.	6 fl. oz.	1.9 fl. oz.	3.2 fl. oz.	1.2 pt.	1.6 pt.
3 Gal.	12.0 fl. oz.	18.0 fl. oz.	5.75 fl. oz.	9.6 fl. oz.	1.8 qt.	2.4 qt.
4 Gal.	1.0 pt.	1.5 pt.	7.7 fl. oz.	12.8 fl. oz.	2.4 qt.	3.2 qt.
5 Gal.	1.25 pt.	1.0 pt. + 14 fl. oz.	9.6 fl. oz.	1.0 pt.	3.0 qt.	1.0 gal.
50 Gal.	1.5 gal. + 8.0 oz.	2.0 gal. + 2.75 pt.	3.0 qt.	1.25 gal.	7.5 gal.	10.0 gal.
100 Gal.	3 gal. + 1.0 pt.	4 gal. + 2.75 qt.	1.5 gal.	2.5 gal.	15.0 gal.	20.0 gal.

16 fl. oz. = 1 pt.; 2 pt. = 1 qt.; 4 qt. = 1 gal.

FORESTRY USE RESTRICTIONS

Maximum Rate – Annual

- **DO NOT** apply more than 48 fluid ounces of this product (1.5 lb. (ae) Imazapyr) per acre per year.

Maximum Rate – Single Application

- **DO NOT** apply more than 48 fluid ounces of this product (1.5 lb. (ae) Imazapyr) per single application.

Maximum Number of Applications Per Year

DO NOT apply more than 1 application per year.

SITE PREPARATION TREATMENT

This product may be used to control labeled actively growing grasses, broadleaf weeds, vines and brambles, and woody brush and trees on Forest sites in advance of regeneration for the following Conifer crop species:

Common Name	Scientific Name	Rate (Fl. Oz./A)
Loblolly pine	<i>Pinus taeda</i>	24 to 40 (0.75 to 1.24 lb. (ae) Imazapyr)
Loblolly X pitch hybrid		
Longleaf pine	<i>Pinus palustris</i>	20 to 32 (0.62 to 1 lb. (ae) Imazapyr)
Shortleaf pine	<i>Pinus echinata</i>	
Virginia pine	<i>Pinus virginiana</i>	
Slash pine	<i>Pinus elliotii</i>	
Coastal redwood	<i>Sequoia sempervirens</i>	12 to 24 (0.40 to 0.75 lb. (ae) Imazapyr)
Douglas fir	<i>Pseudotsuga menziesii</i>	
Incense cedar	<i>Libocedrus decurrens</i>	
Western hemlock	<i>Tsuga heterophylla</i>	12 to 16 (0.40 to 0.50 lb. (ae) Imazapyr)
California red fir	<i>Abies magnifica</i>	
California white fir	<i>Abies concolor</i>	
Jack pine	<i>Pinus banksiana</i>	
Lodgepole pine	<i>Pinus contorta</i>	
Pitch pine	<i>Pinus rigida</i>	
Ponderosa pine	<i>Pinus ponderosa</i>	
Sugar pine	<i>Pinus lambertiana</i>	
White pine	<i>Pinus strobes</i>	
Black spruce	<i>Picea mariana</i>	
Red spruce	<i>Picea rubens</i>	
White spruce	<i>Picea glauca</i>	

Use the specified rate of this product per acre applied as a broadcast foliar spray for long-term control of labeled woody plants and residual control of herbaceous weeds. Within 4 to 6 weeks of treatment, grass and other herbaceous weeds will be controlled and may provide fuel to facilitate a site preparation burn, if desired, to control Conifers or other species resistant to the herbicide.

Apply the specified rate of this product per acre in 5 to 30 gallons total spray solution for helicopter applications or 5 to 100 gallons total spray solution for mechanical ground spray and backpack applications. Use a minimum of 0.25% by volume non-ionic surfactant. Use the higher label rates of this product and higher spray volumes when controlling particularly dense or multilayered canopies of hardwood stands or difficult to control species.

Tank-mixes may be necessary for chemical control of Conifers and other species resistant to this product in certain cases. Always follow the most restrictive directions for use and precautionary statements of each product in the tank-mixture. Combinations with other products labeled for Forest site preparation may kill certain plants including Legumes and Blackberry which are desirable for wildlife habitat.

Where quick initial brown out (deadening of foliage) is desired for burning, apply a tank-mixture of this product with Triclopyr or other products registered for this use at specified label rates per acre. For control of seedling pines, apply this product with Glyphosate or other products registered for this use at specified label rates. For site preparation, rates less than the specified label rates of this product will provide suppression of hardwood brush and trees, and some resprouting may occur.

DO NOT plant seedlings of Black spruce (*Picea mariana*) or White spruce (*Picea glauca*) on sites that have been broadcast treated with this product or into the treated zone of spot or banded applications for three months following application or injury may occur.

HERBACEOUS WEED CONTROL

Use this product for selective weeding in the following Conifers:

Common Name	Scientific Name	Rate (Fl. Oz./A)
Loblolly pine	<i>Pinus taeda</i>	4 to 6 (0.12 to 0.20 lb. (ae) Imazapyr)
Loblolly X pitch hybrid		6 to 10 (0.20 to 0.31 lb. (ae) Imazapyr)
Virginia pine	<i>Pinus virginiana</i>	
Longleaf pine ¹	<i>Pinus palustris</i>	4 to 6 (0.12 to 0.20 lb. (ae) Imazapyr)
Shortleaf Pine ¹	<i>Pinus echinata</i>	
Slash pine ¹	<i>Pinus elliotii</i>	
Douglas fir ¹	<i>Pseudotsuga menziesii</i>	

¹ Use of surfactant is not advised.

This product may be applied as a broadcast treatment, banded over tree rows, or as a directed spray for release of young Conifers from herbaceous weeds. To prevent possibility of Conifer injury, **DO NOT** apply this product when Conifers are under stress from drought, disease, animal or Winter injury, planting shock, or other stresses reducing Conifer vigor. Broadcast applications may be made by helicopter, ground, or backpack sprayer. For difficult to control weeds, use the higher labeled rates. Where herbaceous weeds have overtopped Conifer seedlings, a non-ionic surfactant may be added to improve weed control (except for Slash pine, Long leaf pine, and Douglas fir) at a rate not to exceed 0.25% of spray solution volume. Some minor Conifer growth inhibition may be observed when herbaceous weed control treatments are made during periods of active Conifer growth.

This product may also be applied using backpack or handheld sprayers to control herbaceous weeds around individual Conifer seedlings. Mix 0.4 to 0.6 fluid ounces of this product (0.013 to 0.02 lb. (ae) Imazapyr) and 0.2 fluid ounces non-ionic surfactant per gallon of water. Direct the spray to the weeds and minimize the amount applied to Conifer foliage for best Conifer tolerance. Ensure that maximum labeled rates per acre for listed crop species above are not exceeded. This product may be tank-mixed with a Sulfometuron-methyl product to broaden the spectrum of weeds controlled. Always follow the most restrictive directions for use and precautionary statements of each product in the tank-mixture. For Loblolly pine, apply 4 to 6 fluid ounces of this product (0.12 to 0.2 lb. (ae) Imazapyr) plus a Sulfometuron-methyl product at the specified label rate per acre. The application of this product plus a Sulfometuron-methyl product at the specified label rates on other Conifer species may cause growth suppression.

CONIFER RELEASE TREATMENTS

This product may be applied as a broadcast or directed spray application for suppression of labeled brush, tree, and herbaceous weed species. Directed spray applications may be made with low-volume applications in Conifer stands of all ages by targeting the unwanted vegetation and avoiding direct application to the Conifer. Ensure that maximum labeled rates per acre listed for crop species below are not exceeded.

Use broadcast applications of this product for release of the following Conifers from hardwood competition:

Common Name	Scientific Name	Rate (Fl. Oz./A)
Loblolly pine ³	<i>Pinus taeda</i>	12 to 20 (0.40 to 0.62 lb. (ae) Imazapyr)
Loblolly X pitch hybrid ³		
Virginia pine ³	<i>Pinus virginiana</i>	
Longleaf pine	<i>Pinus palustris</i>	12 to 16 (0.40 to 0.50 lb. (ae) Imazapyr)
Pitch pine	<i>Pinus rigida</i>	
Shortleaf pine	<i>Pinus echinata</i>	
Slash pine	<i>Pinus elliottii</i>	
White pine ¹	<i>Pinus strobes</i>	8 to 16 (0.25 to 0.50 lb. (ae) Imazapyr)
California red fir	<i>Abies magnifica</i>	8 to 12 (0.25 to 0.40 lb. (ae) Imazapyr)
California white fir	<i>Abies concolor</i>	
Lodgepole pine ²	<i>Pinus contorta</i>	
Douglas fir ²	<i>Pseudotsuga menziesii</i>	
Jack pine ²	<i>Pinus banksiana</i>	6 to 12 (0.20 to 0.40 lb. (ae) Imazapyr)
Black spruce ²	<i>Picea mariana</i>	
Red spruce ²	<i>Picea rubens</i>	
White spruce ²	<i>Picea glauca</i>	

¹ DO NOT make applications to White pine stands younger than three years old. To minimize potential White pine injury, release treatments must not be made prior to July 15.

² Applications must be made after formation of final Conifer resting buds in the Fall or height growth inhibitor may occur

³ Mid rotation release: For broadcast applications below the Pine canopy in established stands of Loblolly pine, Loblolly X pitch hybrid, and Virginia pine, use 16 to 32 fl. oz. (0.5 to 1.0 lb. (ae) Imazapyr) product per acre. For mid-rotation release of other species use rates listed above.

Apply the specified rate of this product per acre when making broadcast applications with helicopter or ground spray equipment. Refer to mixing and application instructions for proper spray volumes. A non-ionic surfactant may be added at no more than 0.25% by volume.

Use the higher label rates of this product when controlling particularly dense stands or difficult to control species. Some minor Conifer growth inhibition may be observed when release treatments are made during periods of active Conifer growth. To minimize potential Conifer height growth inhibition, DO NOT make broadcast applications to Conifer stands, except Loblolly pine, before the end of the second growing season. To minimize potential Conifer height growth inhibition, broadcast release treatments may be made late in the growing season. To prevent possibility of Conifer injury, DO NOT apply this product when Conifers are under stress from drought, diseases, animal or winter injury, or other stresses reducing Conifervigor.

This product may be used to release Loblolly pine seedlings during the first growing season following planting or for one-year-old natural Loblolly pine regeneration. For one-year-old Loblolly pine release, apply 12 to 20 fluid ounces per acre of this product (0.40 to 0.62 lb. (ae) Imazapyr) after July 15. The use of rates below 16 fluid ounces (0.50 lb. (ae) Imazapyr) per acre is intended for hardwood growth suppression and some hardwood resprouting must be expected.

FOR SLASH PINE AND LONGLEAF PINE, broadcast release treatments over the top of Pines for the purpose of woody plant control must be made after August 15 and only in stands 2 through 5 years old. For applications over the top of Slash pine and Longleaf pine, DO NOT add surfactant and use lower labeled rates on Sandy soils.

FOR THE AERIAL RELEASE TO SLASH PINE (PINUS ELLIORTII) STANDS OVER THE AGE OF 5 YEARS

This product may be applied as an aerial application for release of Slash pine stands over the age of 5 years. In addition to reading and following all directions in this product, the following precautions and restrictions are required:

- Make applications in the Fall after Slash pine height growth has stopped and buds have set.
- **DO NOT** apply before September 15 even if height growth has stopped and buds have set.
- A maximum of 12 to 14 fluid ounces per acre of this product (0.40 to 0.44 lb. (ae) Imazapyr) may be applied. Use the 12 fluid ounces per acre (0.40 lb. (ae) Imazapyr) rate on *Sandier* sites.

SPOT TREATMENT OF UNDESIRABLE HARDWOOD VEGETATION

This product may be used as a directed foliar or cut stem application to control undesirable brush and hardwoods in the management of stands of all ages for the Conifer species listed in the broadcast application section above. Refer to the mixing and application instructions in the directed foliar or cut stem sections above for proper use rates, equipment, and application techniques. Ensure that the maximum labeled rates per acre listed for crop species are not exceeded. Cut stem applications may be used for spot treatment of undesirable hardwoods in Ponderosa pine stands using 12 fluid ounces (0.40 lb. (ae) Imazapyr) or less of product per acre.

Avoid direct application to desired plant species as injury may occur. Injury may occur to non-target or desirable hardwoods or Conifers if they extend from the same root system or their root systems are grafted to those of the treated tree or if their roots extend into the treated zone.

LATE ROTATION VEGETATION CONTROL IN WESTERN CONIFER

In California, the Pacific Northwest, and Inland Northwest, broadcast aerial applications of this product up to 24 fluid ounces per acre (0.75 lb. (ae) Imazapyr) are permissible in Conifer stands that are targeted for harvesting the year following treatment. Use minimum spray volume of 15 gallons per acre. **DO NOT** use this treatment if Conifer injury or mortality cannot be tolerated.

BAG AND SPRAY APPLICATION FOR CONIFER RELEASE

In Douglas fir and Ponderosa pine stands, broadcast applications of this product up to 16 fluid ounces per acre (0.50 lb. (ae) Imazapyr) are permissible when the trees are covered by bags prior to the application. The bags must prevent the spray mix from contacting the Conifer foliage. On sites with *Coarse textured soils* (e.g., decomposed granite, pumice, sandy or rocky sites) or low levels of soil organic matter (generally 5% or less) significant Conifer growth inhibition and mortality is possible. **DO NOT** use this treatment on these types of sites if Conifer growth inhibition and mortality cannot be tolerated.

NON-AGRICULTURAL LAND USE RESTRICTIONS

Maximum Rate – Annual

- **DO NOT** apply more than 48 fluid ounces of this product (1.5 lb. (ae) Imazapyr) per acre per year.

Maximum Rate – Single Application

- **DO NOT** apply more than 48 fluid ounces of this product (1.5 lb. (ae) Imazapyr) per single application.

Maximum Number of Applications Per Year

DO NOT apply more than 1 application per year. This product may be used for woody and herbaceous weed control in non-agricultural lands including private, public and military lands. Applications are not applicable to treatment of commercial timber or other grown for sale or other commercial use or for commercial seed production or for research purposes.

BRUSH CONTROL

Use the specified rate of this product with the preferred application technique for control of undesirable brush.

Tank-Mixes and Application Rates for Low-Volume Foliar Brush Control*

Target Vegetation	Use Rate of This Product (% by volume)	Tank-Mix
Mixed hardwoods without Elm, Locust or Pine	0.5 to 0.75	Surfactant
Mixed hardwoods containing Elm, Locust and Pine	0.25 to 0.5	Glyphosate plus surfactant
Mixed hardwoods with Locust and Pine but no Elm		Ammonium salt of fosamine plus surfactant
Mixed hardwoods with Locust and Elm but no Pine		Metsulfuron-methyl plus surfactant

* Tank-mixes with 2,4-D or products containing 2,4-D could result in reduced product efficacy.

Backpack and Handheld Spray Mixing Guide

% Solution	Product Per Gallon of Mix (Fl. Oz.)	Product Per 4 Gallon Backpack (Fl. Oz.)
0.25	0.3	1.3
0.5	0.6	2.6
1.0	1.3	5.1
2.0	2.6	10.2
3.0	3.8	15.4
5.0	6.4	25.6

Measuring Chart

128 fluid ounces	=	1 gallon
16 fluid ounces	=	1 pint
8 pints	=	1 gallon
4 quarts	=	1 gallon
2 pints	=	1 quart

FOR SELECTIVE CONTROL OF UNDESIRABLE WEEDS IN UNIMPROVED BERMUDAGRASS AND BAHIAGRASS

This product may be used on unimproved industrial non-cropland Bermudagrass and Bahiagrass turf, including Roadside, Utility rights-of-way and other non-agricultural lands. The application of this product on established common and coastal Bermudagrass and Bahiagrass provides control of labeled broadleaf and grass weeds. Competition from these weeds is eliminated, releasing the Bermudagrass and Bahiagrass. Treatment of Bermudagrass with this product results in a compacted growth habit and seedhead inhibition.

Uniformly apply with properly calibrated ground equipment using at least 10 gallons of water per acre. Temporary yellowing of grass may occur when treatment is made after growth begins. **DO NOT** add surfactant in excess of the specified rate (1 fl. oz. / 25 gal. of spray solution). **DO NOT** apply to grass during its first growing season. **DO NOT** apply to grass that is under stress from drought, disease, insects or other causes.

DOSAGE RATES AND TIMING:

Bermudagrass - Apply this product at 3 to 6 fluid ounces per acre (0.10 to 0.20 lb. (ae) Imazapyr) when the Bermudagrass is dormant. Apply this product at 3 to 4 fluid ounces per acre (0.10 to 0.12 lb. (ae) Imazapyr) after the Bermudagrass has reached full green-up. Applications made during green-up will delay green-up. Include a surfactant in the spray solution.

For additional pre-emergence control of annual grasses and small seeded broadleaf weeds, add Pendimethalin at the appropriately labeled rate. Consult the Pendimethalin label for weeds controlled and for other use directions and precautions.

For control of Johnsongrass in Bermudagrass turf, apply this product at 4 fluid ounces per acre (0.12 lb. (ae) Imazapyr) plus Glyphosate at the appropriate labeled rate plus surfactant. For additional control of broadleaves and vines, Triclopyr may be added to the above mix at the rate of 8 to 16 fluid ounces per acre. Observe all precautions and restrictions on the Triclopyr and Glyphosate labels.

Bahiagrass - Apply this product at 2 to 4 fluid ounces per acre (0.06 to 0.12 lb. (ae) Imazapyr) when the Bahiagrass is dormant or after the grass has initiated green-up but has not exceeded 25% green-up. Include in the spray solution a surfactant (See "ADJUVANT" section for specific use directions for surfactants).

WEEDS CONTROLLED IN UNIMPROVED BERMUDAGRASS AND BAHIAGRASS

Bedstraw (<i>Galium</i> spp.)	Foxtail (<i>Setaria</i> spp.)
Bishopweed (<i>Ptilimnium capillaceum</i>)	Little barley (<i>Hordeum pusillum</i>)
Buttercup (<i>Ranunculus parviflorus</i>)	Seedling Johnsongrass (<i>Sorghum halepense</i>)
Carolina geranium (<i>Geranium carolinianum</i>)	Wild carrot (<i>Daucus carota</i>)
Fescue (<i>Festuca</i> spp.)	White clover (<i>Trifolium repens</i>)
	Yellow woodsorrel (<i>Oxalis stricta</i>)

GRASS GROWTH AND SEEDHEAD SUPPRESSION

This product may be used to suppress growth and seedhead development of certain Turfgrass in unimproved areas. When applied to desirable Turf, this product may result in temporary turf damage and/or discoloration. Effects to the desirable Turf may vary with environmental conditions. For optimum performance, application must be made prior to culm elongation. Applications may be made before or after mowing. If applied prior to mowing, allow at least three days of active growth before mowing. If following a mowing, allow sufficient time for the Grasses to recover before applying this product or injury may be amplified.

DO NOT apply to Turf under stress (drought, cold, insect damaged, etc.) or severe injury or death may occur.

Bermudagrass - Apply this product at 3 to 4 fluid ounces per acre (0.10 to 0.12 lb. (ae) Imazapyr) from early green-up to prior to seed head initiation. **DO NOT** add a surfactant for this application.

Cool Season Unimproved Turf - Apply this product at 1 fluid ounce per acre (0.03 lb. (ae) Imazapyr) plus 0.25% non-ionic surfactant. For increased suppression, this product may be tank-mixed with such products as Glyphosate or Mefluidide.

Tank-mixes may increase injury to desired Turf. Consult each product label for specified Turf species and other use directions and precautions. Tank-mixes with 2,4-D or products containing 2,4-D may decrease the effectiveness of this product.

TOTAL VEGETATION CONTROL WHERE BARE GROUND IS DESIRED

This product is an effective herbicide for pre-emergence or post-emergence control of many annual and perennial broadleaf and grass weeds where bare ground is desired. This product is particularly effective on hard-to-control perennial grasses. This product at 12 to 48 fluid ounces per acre (0.4 to 1.5 lb. (ae) Imazapyr) can be used alone or in tank-mix with herbicides approved for use in bare ground. The degree and duration of control are dependent on the rate of this product used, tank-mix partner, the volume of carrier, soil texture, rainfall and other conditions. Consult manufacturer's labels for specific rates and weeds controlled. Always follow the more restrictive label when making an application involving tank-mixes. Applications of this product may be made anytime of the year. Use equipment calibrated to deliver desired gallons per acre spray volume and uniformly distribute the spray pattern over the treated area.

Post-emergence Applications: Always use a spray adjuvant (See "ADJUVANT" section of this label) when making a post-emergence application. For optimum performance on tough to control annual grasses, applications must be made at a total volume of 100 gallons per acre or less. For quicker burndown or brown-out of target weeds, this product may be tank-mixed with Glyphosate. Tank-mixes with 2,4-D or products containing 2,4-D may reduce the performance of this product. Always follow the more restrictive label when tank-mixing.

Spot Treatments: This product may be used as a follow-up treatment to control escapes or weed encroachment in a bare ground situation. To prepare the spray solution, thoroughly mix in each gallon of water 0.5 to 5% of this product plus an adjuvant. For increased burndown, include Glyphosate. For added residual weed control or to increase the weed spectrum, add products containing Prodiamine or Dicamba. Always follow the more restrictive label when tank-mixing.

FOR CONTROL OF UNDESIRABLE WEEDS UNDER PAVED SURFACES

This product can be used under asphalt, pond liners and other paved areas such as under driveways, cement walk ways. ONLY in industrial sites or where the pavement has a suitable barrier along the perimeter that prevents encroachment of roots of desirable plants.

This product must be used only where the area to be treated has been prepared according to good construction practices. If rhizomes, stolons, tubers or other vegetative plant parts are present in the site, they must be removed by scalping with a grader blade to a depth sufficient to ensure their complete removal. Paving must follow applications of this product as soon as possible. **DO NOT** apply where the product may contact the roots of desirable trees or other plants.

This product is not to be used under pavement on residential properties including driveways or parking lots or for use in recreational areas including under bike or jogging paths, golf cart paths, or tennis courts, or where landscape plantings could be anticipated.

Injury or death of desirable plants may result if this product is applied where roots are present or where they may extend into the treated area. Roots of trees and shrubs may extend a considerable distance beyond the branch extremities (dripline).

Applications must be made to the soil surface only when final grade is established. **DO NOT** move soil following application of this product. Apply this product in sufficient water (at least 100 gal./A) to ensure thorough and uniform wetting of the soil surface, including the shoulder areas. Add this product at a rate of 48 fluid ounces (1.5 lb. (ae) Imazapyr) per acre (1.10 fl. oz. /1000 sq. ft.) to clean water in the spray tank during the filling operation. Agitate before spraying.

If the soil is not moist prior to treatment, incorporation of this product is needed for herbicide activation. This product can be incorporated into the soil to a depth of 4 to 6 inches using a rototiller or disc. Rainfall or irrigation of 1 inch will also provide uniform incorporation. **DO NOT** allow treated soil to wash or move into untreated areas.

SPOT TREATMENTS AND CRACK-AND-CREVICE TREATMENTS

Use this product as an initial or follow up treatment to control weed escapes or weed encroachment in bare ground situations, including cracks and crevices in paved surfaces including Parking lots, Runways and Roadways.

FOR SPOT TREATMENT WEED CONTROL IN GRASS PASTURE AND RANGELAND

For the control of undesirable vegetation in Grass pasture and Rangeland, this product may be applied as a spot treatment at a rate of 1 to 24 fluid ounces of product (0.03 to 0.75 lb. (ae) Imazapyr) per acre using any of the ground application methods as described in this label. Spot applications may not exceed more than one tenth of the area to be grazed or cut for hay in Grass pasture and Rangeland. See appropriate sections of this label for specific use directions for the application method and vegetation control desired.

RESTRICTIONS

Maximum Rate – Annual

- **DO NOT** apply more than 24 fluid ounces of this product (0.75 lb. (ae) Imazapyr) per acre per year.

Maximum Rate – Single Application

- **DO NOT** apply more than 24 fluid ounces of this product (0.75 lb. (ae) Imazapyr) per single application.

Maximum Number of Applications Per Year

- **DO NOT** apply more than 1 application per year.

Grazing and Haying Restrictions:

DO NOT cut forage grass for hay for 7 days after application of this product. There are no grazing restrictions following application of this product.

Rangeland Use Instructions:

This product may be applied to Rangeland for the control of undesirable vegetation to achieve one or more of the following vegetation management objectives:

- Control of undesirable (noxious, invasive and non-native) plant species.
- Control of undesirable vegetation for wildlife habitat improvement.
- Control of undesirable vegetation to aid in the establishment of desirable Rangeland plant species.
- Release of existing desirable Rangeland plant communities from the competitive pressure of undesirable plant species.

- Control of undesirable vegetation to aid in the establishment of desirable vegetation following a fire.

- Control of vegetation to reduce wildfire fuel.

To ensure the protection of threatened and endangered plants, when applying this product to Rangeland:

- Federal agencies must follow NEPA regulations to ensure protection of threatened and endangered plants.

- Other organizations or individuals must operate under a habitat conservation plan if threatened or endangered plants are known to be present on the land to be treated.

- State agencies must work with the Fish and Wildlife Service or the Service's designated state conservation agency to ensure protection of threatened and endangered plants.

See appropriate sections of this label for specific use directions for the desired Rangeland vegetation management control desired.

This product must only be applied to a given Rangeland acre as specific weed problems arise. Long-term control of undesirable weeds ultimately depends on the successful use of the land management practices that promote the sustainability and growth of desirable Rangeland plant species.

ROTATIONAL CROP GUIDELINE

Rotational crops may be planted 12 months after applying this product at the specified Pasture and Rangeland rate. Twelve months after an application of this product, and before planting any crop, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted in the previously treated area in the Grass pasture and Rangeland and grown to maturity. The test strip must include low areas and knolls, and include variations in soil type and pH within the treated area. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

Use of this product in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various agronomic factors and environmental factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

TERRESTRIAL WEEDS CONTROLLED

In Terrestrial sites, this product will provide pre-emergence or post-emergence control with residual control of the following target vegetation species at the rates listed. Residual control refers to control of newly germinating seedlings in both annuals and perennials. In general, annual weeds may be controlled by pre-emergence or post-emergence applications of this product. For established biennials and perennials post-emergence applications of this product are advised. The rates shown below pertain to broadcast applications and indicate the relative sensitivity of these weeds. The relative sensitivity must be referenced when preparing low volume spray solutions (see "Low Volume" section of "GROUND APPLICATION"); low volume applications may provide control of the target species with less product per acre than is shown for the broadcast treatments. This product must be used only in accordance with the "DIRECTIONS FOR USE" on this label.

The relative sensitivity of the species listed below can also be used to determine the relative risk of causing non-target plant injury if any of the below listed species are considered to be desirable within the area to be treated.

Resistant Biotypes: Naturally occurring biotypes (a plant within a given species that has a slightly different, but distinct genetic makeup from other plants of the same species) of some weeds listed on this label may not be effectively controlled. If naturally occurring resistant biotypes are present in an area, this product must be tank-mixed or applied sequentially with an appropriate registered herbicide having a different mode of action to ensure control.

TERRESTRIAL WEEDS CONTROLLED		
COMMON NAME	SCIENTIFIC NAME	GROWTH HABIT ²
GRASS WEEDS		
Apply 16 to 24 fl. oz. /A¹ (0.5 to 0.75 lb. (ae) Imazapyr)		
Annual bluegrass	<i>Poa annua</i>	A
Broadleaf signalgrass	<i>Brachiaria platyphylla</i>	A
Canada bluegrass	<i>Poa compressa</i>	P
Downy brome	<i>Bromus tectorum</i>	A
Fescue	<i>Festuca</i> spp.	A/P
Foxtail	<i>Setaria</i> spp.	A
Italian ryegrass	<i>Lotium multiflorum</i>	A
Johnsongrass ⁴	<i>Sorghum halepense</i>	P
Kentucky bluegrass	<i>Poa pratensis</i>	P
Napier grass ⁵	<i>Pennisetum purpureum</i>	P
Orchardgrass	<i>Dactylis glomerata</i>	P
Paragrass	<i>Brachiaria mutica</i>	P
Quackgrass	<i>Agropyron repens</i>	P
Sandbur	<i>Cenchrus</i> spp.	A
Smooth brome	<i>Bromus inermis</i>	P
Vaseygrass	<i>Paspalum urvillei</i>	P
Wild oats	<i>Avena fatua</i>	A
Witchgrass	<i>Panicum capillare</i>	A
Apply 24 to 32 fl. oz. /A¹ (0.75 to 1.0 lb. (ae) Imazapyr)		
Barnyardgrass	<i>Echinochloa crus-galli</i>	A
Beardgrass	<i>Andropogon</i> spp.	P
Bluegrass, Annual	<i>Poa annua</i>	A
Bulrush ⁵	<i>Scirpus validus</i>	P
Cheat	<i>Bromus secalinus</i>	A
Cogongrass	<i>Imperata cylindrica</i>	P
Crabgrass	<i>Digitaria</i> spp.	A
Crowfootgrass	<i>Dactyloctenium aegyptium</i>	A
Fall panicum	<i>Panicum dichotomiflorum</i>	A
Goosegrass	<i>Eleusine indica</i>	A
Itchgrass	<i>Rottboellia exaltata</i>	A
Lovegrass ⁴	<i>Eragrostis</i> spp.	P
Maidencane ⁵	<i>Panicum hemitomon</i>	A
Panicum, Browntop	<i>Panicum fasciculatum</i>	A
Panicum, Texas	<i>Panicum texanum</i>	A
Prairie threeawn	<i>Aristida oligantha</i>	P
Sandbur, Field	<i>Cenchrus incertus</i>	A
Signalgrass	<i>Brachiaria platyphylla</i>	A
Wild barley	<i>Hordeum</i> spp.	A
Woolly cupgrass	<i>Eriochloa villosa</i>	A
Apply 32 to 48 fl. oz. /A¹ (1.0 to 1.5 lb. (ae) Imazapyr)		
Bahiagrass	<i>Paspalum notatum</i>	P
Bermudagrass ^{3,4}	<i>Cynodon dactylon</i>	P
Big bluestem	<i>Andropogon gerardii</i>	P
Dallisgrass	<i>Paspalum dilatatum</i>	P
Feathertop	<i>Pennisetum villosum</i>	P
Guineagrass	<i>Panicum maximum</i>	P
Saltgrass ³	<i>Distichlis stricta</i>	P
Sand dropseed	<i>Sporobolus cryptandrus</i>	P
Sprangletop	<i>Leptochloa</i> spp.	A
Timothy	<i>Phleum pratense</i>	P
Wirestem muhly	<i>Muhlenbergia frondosa</i>	P
¹ Use higher rate where heavy or well-established infestations occur. ² Growth Habit: A = Annual, B = Biennial, P = Perennial ³ Use a minimum of 75 GPA. ⁴ Use higher labeled rates. ⁵ Use not permitted in California unless otherwise directed by supplemental labeling.		
(Continued)		

TERRESTRIAL WEEDS CONTROLLED (Cont.)		
COMMON NAME	SCIENTIFIC NAME	GROWTH HABIT ²
BROADLEAF WEEDS		
Apply 16 to 24 fl. oz. /A¹ (0.5 to 0.75 lb. (ae) Imazapyr)		
Burdock	<i>Arctium</i> spp.	B
Carolina geranium	<i>Geranium carolinianum</i>	A
Carpetweed	<i>Mollugo verticillata</i>	A
Clover	<i>Trifolium</i> spp.	A/P
Common chickweed	<i>Stellaria media</i>	A
Common ragweed	<i>Ambrosia artemisiifolia</i>	A
Dandelion	<i>Taraxacum officinale</i>	P
Dogfennel	<i>Eupatorium capillifolium</i>	A
Filaree	<i>Erodium</i> spp.	A
Fleabane	<i>Erigeron</i> spp.	A
Hoary vervain	<i>Verbena stricta</i>	P
Indian mustard	<i>Brassica juncea</i>	A
Kochia	<i>Kochia scoparia</i>	A
Lambsquarters	<i>Chenopodium album</i>	A
Lespedeza ³	<i>Lespedeza</i> spp.	P
Miners lettuce	<i>Montia perfoliata</i>	A
Mullein	<i>Verbascum</i> spp.	B
Nettleleaf goosefoot	<i>Chenopodium murale</i>	A
Oxeye daisy	<i>Chrysanthemum leucanthemum</i>	P
Pepperweed	<i>Lepidium</i> spp.	A
Pigweed	<i>Amaranthus</i> spp.	A
Puncturevine	<i>Tribulus terrestris</i>	A
Russian thistle	<i>Salsola kali</i>	A
Smartweed	<i>Polygonum</i> spp.	A/P
Sorrell	<i>Rumex</i> spp.	P
Sunflower	<i>Helianthus</i> spp.	A
Sweet clover	<i>Melilotus</i> spp.	A/B
Tansymustard	<i>Descurainia pinnata</i>	A
Western ragweed	<i>Ambrosia psilostachya</i>	P
Wild carrot	<i>Daucus carota</i>	B
Wild lettuce	<i>Lactuca</i> spp.	A/B
Wild parsnip	<i>Pastinaca saliva</i>	B
Wild turnip	<i>Brassica campestris</i>	B
Woollyleaf bursage	<i>Franseria tomentosa</i>	P
Yellow woodsorrel	<i>Oxalis stricta</i>	P
¹ Use higher labeled rate where heavy or well-established infestations occur. ² Growth Habit: A = Annual, B = Biennial, P = Perennial ³ Use not permitted in California unless otherwise directed by supplemental labeling. ⁴ For best results, early post-emergence applications are required.		
Apply 24 to 32 fl. oz. /A¹ (0.75 to 1.0 lb. (ae) Imazapyr)		
Broom snakeweed	<i>Gutierrezia sarothrae</i>	P
Bull thistle	<i>Cirsium vulgare</i>	B
Burclover	<i>Medicago</i> spp.	A
Chickweed, Mouseear	<i>Cerastium vulgatum</i>	A
Clover hop	<i>Trifolium procumbens</i>	A
Cocklebur	<i>Xanthium strumarium</i>	A
Cudweed	<i>Gnaphalium</i> spp.	A
Desert camelthorn	<i>Alhagi pseudalhagi</i>	P
Dock	<i>Rumex</i> spp.	P
Fiddleneck	<i>Amsinckia intermedia</i>	A
Goldenrod	<i>Solidago</i> spp.	P
Henbit	<i>Lamium amplexicaule</i>	A
Knotweed, Prostrate	<i>Polygonum aviculare</i>	A/P
Pokeweed	<i>Phytolacca americana</i>	P
Purslane	<i>Portulaca</i> spp.	A
Puslev, Florida	<i>Richardia scabra</i>	A
Rocket, London	<i>Sisymbrium irio</i>	A
Rush skeletonweed ⁴	<i>Chondrilla juncea</i>	B
Saltbush	<i>Atriplex</i> spp.	A
Shepherdspurse	<i>Capsella bursa-pastoris</i>	A
Spurge, Annual	<i>Euphorbia</i> spp.	A
Stinging nettle ⁴	<i>Urtica dioica</i>	P
Velvetleaf	<i>Abutilon theophrasti</i>	A
Yellow starthistle	<i>Centaurea solstitialis</i>	A
(Continued)		

TERRESTRIAL WEEDS CONTROLLED (Cont.)		
COMMON NAME	SCIENTIFIC NAME	GROWTH HABIT ²
BROADLEAF WEEDS		
Apply 32 to 48 fl. oz. /A¹ (1.0 to 1.5 lb. (ae) Imazapyr)		
Arrowwood	<i>Pluchea sericea</i>	A
Canada thistle	<i>Cirsium arvense</i>	P
Giant ragweed	<i>Ambrosia trifida</i>	A
Gray rabbitbrush	<i>Chrysothamnus nauseosus</i>	P
Little mallow	<i>Malva parviflora</i>	B
Milkweed	<i>Asclepias</i> spp.	P
Primrose	<i>Oenothera kunthiana</i>	P
Silverleaf nightshade	<i>Solarium elaeagnifolium</i>	P
Sowthistle	<i>Sonchus</i> spp.	A
Texas thistle	<i>Cirsium texanum</i>	P
¹ Use higher rate where heavy or well-established infestations occur.		
² Growth Habit: A = Annual, B = Biennial, P = Perennial		
³ Use not permitted in California unless otherwise directed by supplemental labeling.		
⁴ For best results, early post-emergence applications are required.		
VINES AND BRAMBLES		
Apply 8 fl. oz. /A (0.25 lb. (ae) Imazapyr)		
Field bindweed	<i>Convolvulus arvensis</i>	P
Hedge bindweed	<i>Calystegia sepium</i>	A
Apply 16 to 24 fl. oz. /A¹ (0.5 to 0.75 lb. (ae) Imazapyr)		
Wild buckwheat	<i>Polygonum convolvulus</i>	P
Apply 24 to 32 fl. oz. /A¹ (0.75 to 1.0 lb. (ae) Imazapyr)		
Greenbriar	<i>Smilax</i> spp.	P
Honeysuckle ³	<i>Lonicera</i> spp.	P
Morningglory	<i>Ipomoea</i> spp.	A/P
Poison ivy	<i>Rhus radicans</i>	P
Redvine	<i>Brunnichia cirrhosa</i>	P
Wild rose ³	<i>Rosa</i> spp.	P
Including:		
Multiflora rose	<i>Rosa multiflora</i>	P
Macartney rose	<i>Rosa bracteata</i>	P
Apply 32 to 48 fl. oz. /A¹ (1.0 to 1.5 lb. (ae) Imazapyr)		
Trumpet creeper	<i>Campsis radicans</i>	P
Virginia creeper	<i>Parthenocissus quinquefolia</i>	P
Wild grape	<i>Vitis</i> spp.	P
¹ Use higher labeled rate where heavy or well-established infestations occur.		
² Growth Habit: A = Annual, B = Biennial, P = Perennial		
³ Use higher labeled rate.		
BRUSH SPECIES		
Apply 16 to 32 fl. oz. /A¹ (0.5 to 1.0 lb. (ae) Imazapyr)		
Brazilian peppertree	<i>Schinus terebinthifolius</i>	P
Chinese tallow tree Popcorn tree	<i>Sapium sebiferum</i>	P
Russian olive	<i>Elaeagnus angustifolia</i>	P
Sumac	<i>Rhus</i> spp.	P
Willow	<i>Salix</i> spp.	P
Apply 32 to 48 fl. oz. /A¹ (1.0 to 1.5 lb. (ae) Imazapyr)		
Alder	<i>Alnus</i> spp.	P
American beech	<i>Fagus grandifolia</i>	P
Ash ³	<i>Fraxinus</i> spp.	P
Aspen	<i>Populus</i> spp.	P
Autumn olive	<i>Elaeagnus umbellata</i>	P
Bald cypress	<i>Taxodium distichum</i>	P
Big leaf maple	<i>Acer macrophyllum</i>	P
Birch ³	<i>Betula</i> spp.	P
Black gum ⁴	<i>Nyssa sylvatica</i>	P
Black oak	<i>Quercus kelloggii</i>	P
Boxelder	<i>Acer negundo</i>	P
Ceanothis	<i>Ceanothis</i> spp.	P
Cherry ^{3, 4}	<i>Prunus</i> spp.	P
Chinaberry	<i>Melia azedarach</i>	P
Chinquapin	<i>Castanopsis chrysophylla</i>	P
Cottonwood	<i>Populus trichocarpa</i> P <i>deltoides</i>	P
Cypress	<i>Taxodium</i> spp.	P
Dogwood ³	<i>Cornus</i> spp.	P
Elm	<i>Ulmus</i> spp.	P
Eucalyptus	<i>Eucalyptus</i> spp.	P

(Continued)

TERRESTRIAL WEEDS CONTROLLED (Cont.)		
COMMON NAME	SCIENTIFIC NAME	GROWTH HABIT ²
BRUSH SPECIES		
Apply 32 to 48 fl. oz. /A¹ (1.0 to 1.5 lb. (ae) Imazapyr)		
Hawthorn	<i>Crataegus</i> spp.	P
Hickory ³	<i>Carya</i> spp.	P
Huckleberry	<i>Gaylussacia</i> spp.	P
Lyonia spp. Including: Fetterbush Staggerbush	<i>Lyonia lucida</i> <i>Lyonia mariana</i>	P P
Madrone	<i>Arbutus menziesii</i>	P
Maple	<i>Acer</i> spp.	P
Melaleuca	<i>Melaleuca quinquenervia</i>	P
Mulberry ^{3, 6}	<i>Morus</i> spp.	P
Oak ⁷	<i>Quercus</i> spp.	P
Persimmon ⁴	<i>Diospyros virginiana</i>	P
Poison oak	<i>Rhus diversiloba</i>	P
Poplar	<i>Populus</i> spp.	P
Privet	<i>Ligustrum vulgare</i>	P
Red alder	<i>Alnus rubra</i>	P
Red maple	<i>Acer rubrum</i>	P
Saltcedar	<i>Tamarix pentandra</i>	P
Sassafras	<i>Sassafras albidum</i>	P
Sourwood ⁴	<i>Oxydendrum arboreum</i>	P
Sweetgum	<i>Liquidambar styraciflua</i>	P
Sycamore	<i>Platanus occidentalis</i>	P
Tanoak ³	<i>Lithocarpus densiflorus</i>	P
Titi ⁸	<i>Cyrilla racemiflora</i>	P
Tree of heaven	<i>Ailanthus altissima</i>	P
Vaccinium spp. Including: Blueberry Sparkleberry	<i>Vaccinium</i> spp. <i>Vaccinium arboreum</i>	P P
Water willow ⁹	<i>Justicia americana</i>	P
Yellow poplar ³	<i>Liriodendron tulipifera</i>	P
¹ Use higher labeled rate where heavy or well-established infestations occur.		
² Growth Habit: A = Annual, B = Biennial, P = Perennial		
³ Use higher labeled rate.		
⁴ Best control with applications before formation of Fall leaf color.		
⁵ Tank-mix with Glyphosate.		
⁶ Degree of control may be species dependent.		
⁷ For Water oak (<i>Quercus nigra</i>) Laurel oak (<i>Quercus lauriflora</i>) Willow oak (<i>Quercus phellos</i>) and Live oak (<i>Quercus virginiana</i>) use higher labeled rates.		
⁸ Suppression only.		
⁹ Use not permitted in California unless otherwise directed by supplemental labeling.		

Aquatic Sites**RESTRICTIONS****Maximum Rate – Annual**

DO NOT apply more than 48 fluid ounces of this product (1.5 lb. (ae) Imazapyr) per acre per year.

Maximum Rate – Single Application

• **DO NOT** apply more than 48 fluid ounces of this product (1.5 lb. (ae) Imazapyr) per single application.

Maximum Number of Applications Per Year

• **DO NOT** apply more than 1 application per year.

No Application to Aquatic Sites in New York State.

AQUATIC WEEDS CONTROLLED

This product may be applied for control of floating and emergent weeds (see "AQUATIC WEEDS CONTROLLED" and "TERRESTRIAL WEEDS CONTROLLED" section) in or near bodies of water which may be flowing, non-flowing or transient. This product may be applied to specified Aquatic sites that include Bays, Bogs, Brackish water, Canals, Drainage ditches, Estuaries, Lakes, Marshes, Ponds, Reservoirs, Riparian sites, Rivers, Seasonal wet areas, Seeps, Streams, Swamps and transitional areas between Terrestrial and Aquatic sites and seasonal wet areas. See "USE PRECAUTIONS AND RESTRICTIONS" section of this label for precautions, restrictions, and instructions on Aquatic uses.

Read and observe the following directions if Aquatic sites are present in terrestrial non-crop areas and are part of the intended treatment area:

This product must be applied to the emergent foliage of the target vegetation and has little to no activity on submerged aquatic weeds. Concentrations of this product resulting from direct application to water are not expected to be of sufficient concentration nor duration to control target vegetation. Application must be made in such a way as to maximize spray interception by the target vegetation while minimizing the amount of overspray that enters the water.

This product does not control plants that have a majority of their foliage underwater or plants that are completely submerged.

Product Application: This product must be applied with surface or helicopter application equipment in a minimum of 5 gallons of water per acre. When applying by helicopter, follow directions under the "AERIAL APPLICATIONS" section of this label, otherwise refer to section on "GROUND APPLICATIONS" when using surface equipment.

When applying this product to moving bodies of water applications must be made while traveling upstream to prevent concentration of this herbicide in water. **DO NOT** apply to bodies of water or portions of bodies of water where emergent and/or floating weeds **DO NOT** exist.

Large Application Areas / Oxygen Depletion: When application is to be made to target vegetation that covers a large percentage of the surface area of impounded water, treating the area in strips may avoid oxygen depletion due to decaying vegetation. Oxygen depletion may result in the suffocation of some sensitive aquatic organisms. **DO NOT** treat more than one half of the surface area of the water in a single operation and wait at least 10 to 14 days between treatments. Begin treatment along the shore and proceed outward in bands to allow aquatic organisms to move into untreated areas. Avoid washoff of sprayed foliage by recreational boat backwash or spray boat for 1 hour after application.

Apply this product at 16 to 48 fluid ounces (0.5 to 1.5 lb. (ae) Imazapyr) per acre depending on species present and weed density. **DO NOT apply more than 48 fluid ounces of this product (1.5 lb. (ae) Imazapyr) per acre per year.** Use the higher labeled rates for heavy

weed pressure. Consult the "AQUATIC WEEDS CONTROLLED" section and the "ADDITIONAL WEEDS CONTROLLED" section of this label for specific rates. This product may be applied as a draw down treatment in areas described above. Apply this product to weeds after water has been drained and allow 14 days before reintroduction of water.

This product will control the following target species as specified in the Use Rates and Application Directions section of the table. Rate instructions are expressed in terms of product volume for broadcast applications and as a percent solution for directed applications including spot treatments. For percent solution applications. **DO NOT apply more than 48 fluid ounces of this product (1.5 lb. (ae) Imazapyr) per acre per year.**

Mixing Guide

% Solution	Product Per Gallon of Mix (Fl. Oz.)
0.25	0.3 (0.008 lb. (ae) Imazapyr)
0.5	0.6 (0.016 lb. (ae) Imazapyr)
1.0	1.3 (0.04 lb. (ae) Imazapyr)
2.0	2.6 (0.08 lb. (ae) Imazapyr)
3.0	3.8 (0.12 lb. (ae) Imazapyr)
5.0	6.4 (0.20 lb. (ae) Imazapyr)

Measuring Chart

128 fluid ounces	=	1 gallon
16 fluid ounces	=	1 pint
8 pints	=	1 gallon
4 quarts	=	1 gallon
2 pints	=	1 quart

Common Name	Scientific Name	Use Rates and Application Directions
Floating Weeds		
*Floating heart	<i>Nymphodes</i> spp.	16 to 32 fl. oz./A (0.5 to 1 lb. (ae) Imazapyr) applied in 100 GPA water mix. Ensure 100% coverage of actively growing emergent foliage.
*Frogbit	<i>Limnobium spongia</i>	8 to 16 fl. oz./A (0.25 to 0.5 lb. (ae) Imazapyr) applied in 100 GPA water mix. Ensure 100% coverage of actively growing emergent foliage.
*Spatterdock	<i>Nuphar luteum</i>	Apply a tank-mix of 16 to 32 fl. oz./A (0.5 to 1 lb. (ae) Imazapyr) of this product + Glyphosate in 100 GPA water for best control. Ensure 100% coverage of actively growing emergent foliage.
*Water hyacinth	<i>Eichhornia crassipes</i>	8 to 16 fl. oz./A (0.25 to 0.5 lb. (ae) Imazapyr) applied in 100 GPA water to actively growing foliage.
*Water lettuce	<i>Pistia stratiotes</i>	8 to 16 fl. oz./A (0.25 to 0.5 lb. (ae) Imazapyr) applied in 100 GPA water mix. Ensure 100% coverage of actively growing emergent foliage.
Emerg Weed		
*Alligatorweed	<i>Alternanthera philoxeroides</i>	8 to 32 fl. oz./A (0.25 to 1 lb. (ae) Imazapyr) applied in 100 GPA water mix. Ensure 100% coverage of actively growing emergent foliage.
*Arrowhead duck potato	<i>Sagittaria</i> spp.	8 to 16 fl. oz./A (0.25 to 0.5 lb. (ae) Imazapyr) applied to 100 GPA water mix. Ensure 100% coverage of actively growing emergent foliage.
*Bacopa lemon	<i>Bacopa</i> spp.	8 to 16 fl. oz./A (0.25 to 0.5 lb. (ae) Imazapyr) applied in 100 GPA water mix. Ensure 100% coverage of actively growing emergent foliage.
*Parrot feather	<i>Myriophyllum aquaticum</i>	Foliage must be above water for sufficient product uptake. Apply 16 to 32 fl. oz./A (0.5 to 1 lb. (ae) Imazapyr) (0.2% to 0.5% solution) of this product to actively growing emergent foliage.
*Pennywort	<i>Hydrocotyle</i> spp.	8 to 16 fl. oz./A (0.25 to 0.5 lb. (ae) Imazapyr) applied in 100 GPA water mix. Ensure 100% coverage of actively growing emergent foliage.
*Pickerelweed	<i>Pontedena cordata</i>	16 to 24 fl. oz./A (0.5 to 0.75 lb. (ae) Imazapyr) applied in 100 GPA water mix. Ensure 100% coverage of actively growing emergent foliage.
*Taro wild Coco yam Dasheen Elephant's ear	<i>Colocasia esculentum</i>	32 to 48 fl. oz./A (1 to 1.5 lb. (ae) Imazapyr) applied in 100 GPA with a high quality sticker adjuvant. Ensure good coverage of actively growing emergent foliage.
*Water chestnut	<i>Trappa natans</i>	32 to 48 fl. oz./A (1 to 1.5 lb. (ae) Imazapyr) applied in 100 GPA with a high quality sticker adjuvant. Ensure good coverage of actively growing emergent foliage.
*Water lily	<i>Nymphaea odorata</i>	16 to 24 fl. oz./A (0.5 to 0.75 lb. (ae) Imazapyr) applied in 100 GPA water mix. Ensure 100% coverage of actively growing emergent foliage.
*Water primrose	<i>Ludwigia uruguayensis</i>	32 to 48 fl. oz./A (1 to 1.5 lb. (ae) Imazapyr) (0.5% to 0.75% solution). Ensure 100% coverage of actively growing emergent foliage.

(Continued)

(Cont.)		
Common Name	Scientific Name	Use Rates and Application Directions
Terrestrial/Marginal Weeds		
*Aquatic nightshade Soda apple	<i>Solanum tampicense</i>	16 fl. oz./A (0.5 lb. (ae) Imazapyr) (0.25% solution) applied to foliage.
*Bamboo Japanese	<i>Phyllostachys</i> spp.	24 to 32 fl. oz./A (0.75 to 1 lb. (ae) Imazapyr) (0.375% to 0.5% solution) applied to foliage.
*Beach vitex	<i>Vitex rotundifolia</i>	2.5% solution + MSO foliar spray. 8.5% solution stem injection (hack and squirt).
Brazilian pepper Christmasberry	<i>Schinus terebinthifolius</i>	16 to 32 fl. oz./A (0.5 to 1 lb. (ae) Imazapyr) (0.25% to 0.5% solution) applied to foliage.
Cattail	<i>Typha</i> spp.	16 to 32 fl. oz./A (0.5-1 lb. (ae) Imazapyr) (0.25% to 0.5% solution) applied to actively growing green foliage after full leaf elongation. Lower rates will control cattail in the North. Higher rates are needed in the South.
Chinese tallow tree	<i>Sapium sebiferum</i>	8 to 12 fl. oz./A (0.25 to 0.40 (ae) Imazapyr) applied to foliage.
Cogongrass	<i>Imperata cylindrical</i>	Burn foliage, till area, then fall spray 32 fl. oz./A (1 lb. (ae) Imazapyr) (0.5% solution) of this product+MSO applied to new growth.
Cordgrass prairie	<i>Spartina</i> spp.	32 to 48 fl. oz./A (1 to 1.5 lb. (ae) Imazapyr) (0.5% to 0.75% solution) applied to actively growing foliage.
*Cutgrass	<i>Zizaniopsis miliacea</i>	32 to 48 fl. oz./A (1 to 1.5 lb. (ae) Imazapyr) (0.5% to 0.75% solution) applied to actively growing foliage.
*Elephant grass Napier grass	<i>Pennisetum purpureum</i>	24 fl. oz./A (0.75 lb. (ae) Imazapyr) (0.375% solution) applied to actively growing foliage.
*Flowering rush	<i>Butomus umbellatus L</i>	16 to 24 fl. oz./A (0.5-0.75 lb. (ae) Imazapyr) (0.25% to 0.375% solution) applied to actively growing foliage.
Giant reed Wild cane	<i>Arundo donax</i>	32 to 48 fl. oz./A (1.0 to 1.5 lb. (ae) Imazapyr) (0.5% to 0.75% solution) applied in spring to actively growing foliage.
*Golden bamboo	<i>Phyllostachys aurea</i>	24 to 32 fl. oz./A (0.75 to 1 lb. (ae) Imazapyr) (0.375% to 0.5% solution) applied to foliage when plant is actively growing, before setting seedhead. More foliage will result in greater herbicide uptake, resulting in greater root kill.
Junglerice	<i>Echinochloa colonum</i>	24 to 32 fl. oz./A (0.75 to 1 lb. (ae) Imazapyr) (0.375% to 0.5% solution) applied to actively growing foliage.
Knapweed	<i>Centaurea</i> spp.	Russian knapweed: 16 to 24 fl. oz./A (0.5 to 0.75 lb. (ae) Imazapyr) (0.25% to 0.375% solution) + MSO fall applied after senescence begins.
Knotweed, Japanese	<i>Polygonum cuspidatum</i> <i>Fallopia japonica</i>	24 to 32 fl. oz./A (0.75 to 1 lb. (ae) Imazapyr) (0.375% to 0.5% solution) applied post-emergence to actively growing foliage.
Melaleuca Paperbark tree	<i>Melaleuca quinquenervia</i>	<ul style="list-style-type: none"> Established stands: apply 48 fl. oz./A (1.5 lb. (ae) Imazapyr) (0.75% solution) of this product + Glyphosate + spray adjuvant. For best results use MSO as an adjuvant. Broadcast foliar control: apply aerially in a minimum of 2 passes at 10 gal./A applied cross treatment. Spot treatment: use 12.5% of this product + Glyphosate+MSO in water applied as a frill or stump treatment.
*Nutgrass Kili'p'opu	<i>Cyperus rotundus</i>	16 fl. oz./A (0.5 lb. (ae) Imazapyr) (0.25% solution) this product + MSO applied early post-emergence.
*Nutsedge	<i>Cyperus</i> spp.	16 to 24 fl. oz./A (0.5 to 0.75 (ae) Imazapyr) (0.25% to 0.375% solution) post-emergence to foliage or pre-emergence incorporated, non-incorporated pre-emergence applications will not control.
Phragmites Common reed	<i>Phragmites australis</i>	32 to 48 fl. oz./A (1 to 1.5 lb. (ae) Imazapyr) (0.5% to 0.75% solution) applied to actively growing green foliage after full leaf elongation. Ensure 100% coverage. If stand has a substantial amount of old stem tissue, mow or burn, allow to regrow to approximately 5 feet tall before retreatment. Lower rates will control phragmites in the north, higher rates are needed in the south.
*Poison hemlock	<i>Conium maculatum</i>	16 fl. oz./A (0.5 lb. (ae) Imazapyr) (0.25% solution) this product + MSO applied pre-emergence to early post-emergence to rosette before flowering.
Purple loosestrife	<i>Lynthrum salicana</i>	8 fl. oz./A (0.25 lb. (ae) Imazapyr) (0.125% solution) applied to actively growing foliage.
Reed canarygrass	<i>Phalaris arundinacea</i>	24 to 32 fl. oz./A (0.75 to 1 lb. (ae) Imazapyr) (0.375% to 0.5% solution) applied to actively growing foliage.
Rose swamp	<i>Rosa palustris</i>	16 to 24 fl. oz./A (0.5-0.75 lb. (ae) Imazapyr) (0.25% to 0.375% solution) applied to actively growing foliage.
Russian olive	<i>Elaeagnus angustifolia</i>	16 to 32 fl. oz./A (0.5 to 1 lb. (ae) Imazapyr) (0.5% solution) applied to foliage.
Saltcedar Tamarisk	<i>Tamarix</i> spp.	Aerial application: 32 fl. oz. (1 lb. (ae) Imazapyr) this product + NIS applied to actively growing foliage during flowering. Spot treatment: Use 0.50% solution of this product+ NIS and spray to wet foliage. After application, wait at least 2 years before disturbing treated saltcedar. Earlier disturbance can reduce overall control.
Smartweed	<i>Polygonum</i> spp.	16 fl. oz./A (0.5 lb. (ae) Imazapyr) (0.25% solution) applied early post-emergence.
Sumac	<i>Rhus</i> spp.	16 to 24 fl. oz./A (0.5 to 0.75 lb. (ae) Imazapyr) (0.25% to 0.375% solution) applied to foliage.
Swamp morningglory Kangkong Water spinach	<i>Ipomoea aquatic</i>	8 to 16 fl. oz./A (0.25 to 0.5 lb. (ae) Imazapyr) (0.125% to 0.25% solution) of this product + MSO applied early post-emergence.
Torpedo grass	<i>Panicum repens</i>	32 fl. oz./A (1 lb. (ae) Imazapyr) (0.5% to 0.375% solution). Ensure good coverage to actively growing foliage.
*White top Hoary cress	<i>Cardaria draba</i>	8 to 16 fl. oz./A (0.25 to 0.5 lb. (ae) Imazapyr) (0.125% to 0.25% solution) applied in Spring to foliage during flowering.
Willow	<i>Salix</i> spp.	16 to 24 fl. oz./A (0.5 to 0.75 lb. (ae) Imazapyr) (0.25% to 0.375% solution) of this product applied to actively growing foliage. Ensure good coverage.

*Use not permitted in California unless otherwise directed by supplemental labeling.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container. Keep container tightly closed. Keep away from heat and flame.

PESTICIDE DISPOSAL: To avoid waste, use all materials in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often, such programs are run by state or local governments or by industry).

CONTAINER HANDLING:

Nonrefillable Container (plastic rigid material; ≤ 5 gallons):

Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container one-fourth full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Dispose of empty container in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. **Nonrefillable Container (plastic rigid material; > 5 gallons up to < 250 gallons):**

Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container one-fourth full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Dispose of empty container in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. **Refillable plastic Container (≥ 250 gallons and Bulk):**

Refillable container. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

WARRANTY—CONDITIONS OF SALE

OUR DIRECTIONS FOR USE of this product are based upon tests believed reliable. Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically directed and other influencing factors in the use of this product are beyond the control of the Seller. To the extent consistent with applicable laws, Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith. To the extent consistent with applicable laws, in no case shall the Manufacturer or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product when such use and/or handling is not in strict accordance with directions given herewith. The foregoing is a condition of sale by the Seller and is accepted as such by the Buyer.

THE DREXEL logo is a registered trademark of Drexel Chemical Company. All other brand names, product names or trademarks belong to their respective holders.