

TRACO

COPPER SULPHATE FUNGICIDE with ZINC

ACTIVE INGREDIENT

*Copper, Expressed as Elemental

Inert Ingredients

(Inert contains 22.0% Zinc expressed as elemental. *Copper derived from Copper Sulphate, Pentahydrate, and Zinc derived from Zinc Sulphate, Monohydrate, Zinc is for correction of Zinc deficiency.)

By Weight

9.5%

90.5%

CAUTION

Keep out of reach of children and animals

**HARMFUL IF SWALLOWED — AVOID BREATHING DUST
AVOID CONTAMINATION OF FEED AND FOODSTUFFS**

Use of this material in a manner other than suggested may produce injury to plants and undesirable results.

Dispose of wastes by burying away from water supply in non-crop sites. Burn empty bags or bury with wastes. Avoid inhaling smoke from burning bags. Avoid drifting of product onto neighboring crop or non-crop areas otherwise injury of animal, plant or crops may occur. In California, dispose of waste and empty containers in accordance with State and local regulations.

USAGE CAUTIONS

This product is toxic to fish. Keep out of lakes, streams and ponds. Do not apply when weather conditions favor drift from area treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

Apply this product only as specified on this label.

CONDITIONS OF SALE

1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for use as directed on the label. No one, other than an officer of Seller is authorized to make any other warranty, guarantee or direction concerning this product.

2. Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's liability from the handling, storage and use of this product is limited to replacement of product or refund of the base price.

MIXING INSTRUCTIONS

Keep agitator running at all times.

Mechanically Agitated Tank:

Fill tank half full with water, sprinkle product into water when tank is 75% full, add hydrated lime and fill tank.

By-Pass Agitated Tank:

Prepare stock solutions of product and hydrated lime. Fill tank 65% full with water, add product then hydrated lime, continue agitation and fill tank.

Product is corrosive to steel.

DIRECTIONS FOR USE

Citrus — Central California: To control Brown Rot and Septoria mix 5 lbs. of Traco Copper Sulphate Fungicide with Zinc with 6 lbs. of hydrated lime in 100 gallons of water. Use as a full coverage spray. Apply 500 to 700 gallons of finished spray per acre to mature trees in Fall prior to wet season.

Citrus — Southern California: To control Brown Rot mix 3 lbs. of Copper Sulphate Fungicide with Zinc with 4 lbs. of hydrated lime in 100 gallons of water. Use as a skirt spray on soil, trunk and lower 3-4 feet of foliage. Apply in Fall just prior to wet season. In Coastal areas or other areas where no Copper injury has been found use a mixture of 5 lbs. of Copper Sulphate Fungicide with Zinc with 6 lbs. of hydrated lime in 100 gallons of water as a full coverage spray.

EPA Reg. No. 20004-4

NET WEIGHT

LBS.

Manufactured For:

Traylor Chemical & Supply Co., Inc.

Orlando, Florida

Royal Center, Indiana

Fresno, California

ACCEPTED