

PM-15 20552-2 P 1/4

US ENVIRONMENTAL PROTECTION AGENCY  
OFFICE OF PESTICIDES PROGRAMS  
REGISTRATION DIVISION (TS-767)  
WASHINGTON, DC 20460

EPA REGISTRATION  
20552-2

DATE OF ISSUANCE

MAR 30 1987

TERM OF ISSUANCE  
Until Reregistration

NOTICE OF PESTICIDE:  REGISTRATION  
 REREГИSTRATION  
*(Under the Federal Insecticide, Fungicide  
and Rodenticide Act, as amended)*

NAME OF PESTICIDE PRODUCT  
Evergreen Fly Block (with Rabon  
Oral Larvicide)

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

Evergreen Mills, Inc.  
221 North Rennie  
P.O. Box 542  
Ada, OK 74820

160/178786  
18/1

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally reregistered in accordance with FIFRA section 3(c)(7)(A) provided that you:

1. Submit/cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data.

2. Make the labeling changes listed below before you release the product for sale:

- a. On the front panel change "EPA Registration No. 20552-2."
- b. Under the "Precautionary Statements" section delete the sentence "Toxic to Bees."
- c. On the front panel change "animals" to read "cattle."

3. The following comments pertain to product chemistry.

- a. Provide certified limits on all ingredients listed on the Confidential Statement of Formula (CSF). The lower limit for the actives must not fall below label claim. These limits are based on pure active ingredients rather than the technical or concentrate (CSF Form enclosed).

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

DATE

3/23/87  
4/1/87

BEST AVAILABLE COPY

b. Please provide a description of the formulation process.

4. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

George T. LaRocca  
Product Manager (15)  
Insecticide-Rodenticide Branch  
Registration Division (TS-767C)

Enclosures

**BEST AVAILABLE COPY**

33.3 LBS (NET)

12 2 14

EVERGREEN FLY BLOCK  
(WITH RABON® ORAL LARVICIDE)

To prevent the development of horn flies, face flies, house flies and stable flies  
in the manure of treated animals.

ACTIVE INGREDIENT:

2-chloro-1-(2, 4, 5-trichlorophenyl)

vinyl dimethyl phosphate . . . . . 0.30%

INERT INGREDIENTS\* . . . . . 99.70%

TOTAL . . . . . 100.00%

\*Refers only to ingredients which are not larvicidal.

\*\*RABON Insecticide - Contains 1362 mg of RABON per pound

RABON is the registered trademark of Shell Oil Company.

MADE IN U.S.A.  
NET WT. 33.3 LBS

MAR 3 1967

EPA REG. NO. 20552

EPA EST. NO.

Under the  
Federal, and  
as amended, for  
display under 101

GUARANTEED FEED ANALYSIS

20552-2

Crude Protein, minimum . . . . .	12.00%
Crude Fat, minimum . . . . .	1.50%
Crude Fiber, maximum . . . . .	8.00%
Calcium (Ca), maximum . . . . .	5.00%
Calcium (Ca), minimum . . . . .	4.00%
Phosphorus (P), minimum . . . . .	1.25%
Salt (NaCl), maximum . . . . .	21.00%
Salt (NaCl), minimum . . . . .	18.00%
Iodine (I), minimum . . . . .	0.001%
Vitamin A, minimum . . . . .	35,000 USP Units/lb.
Vitamin D <sub>3</sub> , minimum . . . . .	7,500 USP Units/lb.

FEED INGREDIENTS

Soybean meal, cottonseed meal, dried cane molasses, dehydrated alfalfa meal, cane molasses, salt, calcium carbonate, monocalcium phosphate and dicalcium phosphate, magnesium oxide, iron oxide, iron sulfate, manganese oxide, zinc oxide, ethylenediamine dihydride, copper sulfate, cobalt sulfate, vitamin A supplement, d-activated animal sterol (source of vitamin D<sub>3</sub>) vitamin E supplement.

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

SEE REVERSED SIDE OF TAG FOR ADDITIONAL  
PRECAUTIONARY STATEMENTS AND USE DIRECTIONS.

MANUFACTURED BY:  
EVERGREEN MILLS, INC.  
P. O. Box 548 Ada, Oklahoma 74820

BEST BEFORE DATE

11/4

# PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS CAUTION

Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

## ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in a dry place in original container.

Container Disposal: Completely empty bag into feeding equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on-site, or an approved waste disposal facility.

---

## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Allow free choice to cattle. Cattle should consume an average of 0.8 ounces of the Fly Block per 100 pounds of body weight per day, which is equivalent to 8 ounces for a 1000 lb. animal daily. This will supply the recommended average daily intake of 70 mgs. RABON ORAL LARVICIDE per hundred pounds of body weight. Animals should not be starved for minerals before initiation of the EVERGREEN Fly Block with Rabon program but OMIT all salt from other feed after the program is started. Do not feed other free choice sources of salt or mineral containing salt in any form. Allow one block for each 10 head of cattle and place near watering and loafing areas. Put out no more than two blocks supply at a time.

If consumption of EVERGREEN Fly Block with Rabon above 0.8 ounce per 100 lbs. of body weight per day, reduce the number of feed locations or relocate. If consumption is below 0.8 ounce per 100 lbs. of body weight per day, increase the number of feeding locations. Start feeding EVERGREEN Fly Block with Rabon early in the spring before flies begin to appear, and continue feeding throughout the summer and into the fall.

EVERGREEN Fly Block with Rabon may be fed up to slaughter and to lactating dairy cows.

EVERGREEN Fly Block with Rabon prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle, but is not effective against existing adult flies. When starting a feeding program during the fly season, it may be necessary to use other control measures to reduce the population of existing adult flies.

Supplemental control measures may be needed in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle manure but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control EVERGREEN Fly Block with Rabon should be used in conjunction with other good management and sanitation practices.

