

We make it right.

33.3 LBS (NET)



EVERGREEN FLY BLOCK

It's more than a promise. It's a guarantee.

(WITH RABON® ORAL LARVICIDE)

To prevent the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle

ACTIVE INGREDIENTS

2-chloro-1-(2,4,5-trichlorophenyl)

ethyl dimethyl phosphate

0.30%

INERT INGREDIENTS

99.70%

TOTAL 100.00%

*Refers only to ingredients which are not larvicidal

**RABON Insecticide - Contains 1362 mg. of RABON per pound


RABON is the registered trademark of Shell Oil Company

EPA REG. NO. 20552-2

EPA EST. NO. 20552-OK-001

GUARANTEED ANALYSIS

Crude Protein, minimum	12.00%
Crude Fat, minimum	1.50%
Crude Fiber, maximum	8.00%
Calcium (Ca), maximum	5.00%
Calcium (Ca), minimum	4.00%
Phosphorus (P), minimum	1.25%
Salt (NaCl), maximum	21.00%
Salt (NaCl), minimum	18.00%
Moisture (H), minimum	0.001%
Vitamin A, minimum	35,000 USP Units/Lb.
Vitamin D ₃ , minimum	7,500 USP Units/Lb.



FEED INGREDIENTS

ground grain sorghum, soybean meal, cottonseed meal, ground yellow corn, dried cane molasses, ...

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE REVERSED SIDE OF TAG FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND USE DIRECTIONS.

EVERGREEN MILLS, INC. Box 548, Ada, OK 74820

BEST AVAILABLE COPY

2 of 3

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS CAUTION

Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in a dry place in original container.

Container Disposal: Completely empty bag into feeding equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on-site, or at an approved waste disposal facility.

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Allow free choice to cattle. Cattle should consume an average of 0.8 ounces of the Fly Block per 100 pounds of body weight per day, which is equivalent to 8 ounces for a 1000 pound animal daily. This will supply the recommended average daily intake of 70 mgs. RABON ORAL LARVICIDE per hundred pounds of body weight. Animals should not be starved for minerals before initiation of the EVERGREEN Fly Block with Rabon program but OMIT all salt from other feed after the program is started. Do not feed other free choice sources of salt or mineral containing salt in any form. Allow one block for each 10 head of cattle and place near watering and loafing areas. Put out no more than two weeks supply at a time.

If consumption of EVERGREEN Fly Block with Rabon above 0.8 ounce per 100 pounds of body weight per day, reduce the number of feed locations or relocate. If consumption is below 0.8 ounce per 100 pounds of body weight per day, increase the number of feeding locations. Start feeding EVERGREEN Fly Block with Rabon early in the spring before flies begin to appear, and continue feeding throughout the summer and into the fall.

EVERGREEN Fly Block with Rabon may be fed up to slaughter and to lactating dairy cows.

EVERGREEN Fly Block with Rabon prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle, but is not effective against existing adult flies. When starting a feeding program during the fly season, it may be necessary to use other control measures to reduce the population of existing adult flies.

Supplemental fly control measures may be needed in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle manure but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control EVERGREEN Fly Block with Rabon should be used in conjunction with other good management and sanitation practices.

61

BEST AVAILABLE COPY