

mosquito
mosquito
mosquito

Altosid SR-10


Mosquito Growth Regulator

Prevents Emergence of Adult Floodwater Mosquitoes

CAUTION: KEEP OUT OF REACH OF CHILDREN

See Guide to Product Application

NET CONTENTS 1 GALLON (3.78 LITERS)



DIRECTIONS FOR USE

INSTRUCTIONS FOR USE

CAUTION: KEEP OUT OF REACH OF CHILDREN

STORAGE AND DISPOSAL

STORAGE
DISPOSAL

WARRANTY AND CONDITIONS OF SALE

RECOMMENDED APPLICATIONS

INTRODUCTION

floodwater mosquitoes

mosquitoes

no effect when applied to pupae or adult

Guide to Product Application

METHODS OF APPLICATION

Aerial

Ground


APPLICATION RATE

APPLICATION SITES

Pastures

Rice

Intermittently Flooded Noncrop Areas



dense vegetation or canopy

Guide to Product Application

150

zoëcon

Altosid® SR-10

Mosquito Growth Regulator

Prevents Emergence of Adult Floodwater Mosquitoes


ACTIVE INGREDIENT methoprene (isopropyl (E)-11-methoxy-3,7,11-trimethyl-2,4-dodecadienoate)	10.0%
INERT INGREDIENTS	90.0
TOTAL	100.0

Altosid SR-10 is a registered trademark of Zoëcon, Inc.
 Zoëcon, Inc., 10000 E. 1st Ave., Denver, CO 80231
 Altosid SR-10 is a registered trademark of Zoëcon, Inc.

CAUTION: KEEP OUT OF REACH OF CHILDREN

Because of the danger to children from ALTOSID SR-10, it is recommended that this product be stored in a secure container. See Guide to Product Application.

NET CONTENTS 1 GALLON (3.78 LITERS)



DIRECTIONS FOR USE

Follow all directions on label and on product container for use.

CAUTION: KEEP OUT OF REACH OF CHILDREN

STORAGE AND DISPOSAL

STORAGE: Store in a cool, dry place.
DISPOSAL: Do not empty container into water. Dispose of contents and container as directed on label.

INSTRUCTIONS FOR USE:

1. Shake Well. Before using ALTOSID SR-10, shake container thoroughly to mix contents.
 2. Apply ALTOSID SR-10 to water in a uniform manner.
 3. Do not apply ALTOSID SR-10 to water in a pond or lake where there are fish or other aquatic life.
 4. Do not apply ALTOSID SR-10 to water in a pond or lake where there are birds or other wildlife.
 5. Do not apply ALTOSID SR-10 to water in a pond or lake where there are mammals or other wildlife.

WARRANTY AND CONDITIONS OF SALE

This warranty is made by Zoëcon, Inc. on the condition that the user of this product will follow the directions on the label and on the product container for use.

BEST DOCUMENT AVAILABLE

zoëcon

RECOMMENDED APPLICATIONS

INTRODUCTION

ALTOSID SR-10 is a registered trademark of Zoëcon, Inc. and is used to prevent the emergence of adult floodwater mosquitoes. ALTOSID SR-10 is a growth regulator that prevents the development of the pupae stage of the mosquito. ALTOSID SR-10 has no effect when applied to pupae or adult mosquitoes. ALTOSID SR-10 is a registered trademark of Zoëcon, Inc. and is used to prevent the emergence of adult floodwater mosquitoes. ALTOSID SR-10 is a growth regulator that prevents the development of the pupae stage of the mosquito. ALTOSID SR-10 has no effect when applied to pupae or adult mosquitoes. ALTOSID SR-10 is a registered trademark of Zoëcon, Inc. and is used to prevent the emergence of adult floodwater mosquitoes.

METHODS OF APPLICATION

Aerial: Use a backpack sprayer or a fixed-wing aircraft to apply ALTOSID SR-10 to water. Apply ALTOSID SR-10 to water in a uniform manner. Do not apply ALTOSID SR-10 to water in a pond or lake where there are fish or other aquatic life.

Ground: Use a backpack sprayer or a fixed-wing aircraft to apply ALTOSID SR-10 to water. Apply ALTOSID SR-10 to water in a uniform manner. Do not apply ALTOSID SR-10 to water in a pond or lake where there are fish or other aquatic life.

APPLICATION RATE

Apply 4.0 fl. oz. of ALTOSID SR-10 per acre of water.

APPLICATION SITES

Pastures: ALTOSID SR-10 may be applied to pastures where there are floodwater mosquitoes.

Rice: ALTOSID SR-10 may be applied to rice fields where there are floodwater mosquitoes. Do not apply ALTOSID SR-10 to rice fields where there are fish or other aquatic life.

Intermittently Flooded Noncrop Areas: ALTOSID SR-10 may be applied to intermittently flooded noncrop areas where there are floodwater mosquitoes.

Typical sites include: tree water, swamps, marshes, wood and pools, and meadows. Do not apply ALTOSID SR-10 to areas where there are fish or other aquatic life.

In dense vegetation or canopy areas, ALTOSID SR-10 may be applied to the water surface. Do not apply ALTOSID SR-10 to areas where there are fish or other aquatic life.

ZORCON
EZ-515876

AN INSECT GROWTH REGULATOR FOR THE CONTROL OF INSECT PESTS IN TOBACCO PROCESSING AND FOOD HANDLING ESTABLISHMENTS.

ACTIVE INGREDIENTS:

Methoprene (Isopropyl 2,2-11-Methoxy 3,7,11-trimethyl-2,4 dodecadienate)	65.13
Xylene Range Aromatic Solvents	30.42
INERT INGREDIENTS	13.91
TOTAL	100.00

This product contains 5 lb/gallon (600 grams/liter) methoprene active ingredient.
U.S. Patents 3,904,462 & 3,912,815
EPA Reg. No. 20956-101

EPA Est. No. 20956-CA-1

KEEP OUT OF REACH OF CHILDREN
CAUTION
See Side Panel
For Additional Precautionary Statements.

STORAGE AND DISPOSAL

Storage: Store in a cool dry place. Do not contaminate water, food or feed by storage or disposal.
Disposal: Pesticide, spray mixture or residue that cannot be used or chemically reprocessed should be disposed of according to procedures approved by Federal, State or local disposal authorities.
Triple rinse container and offer for recycling, reconditioning, or disposal in approved landfill or burn in a safe place.
Consult Federal, State or local disposal authorities for approved alternative procedures.
Prohibition: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

INTRODUCTION

EZ-515876 is an emulsifiable formulation of the insect growth regulator methoprene. It is designed for use in mechanical foggers and as a concentrate for an aqueous emulsive spray. Methoprene, unlike traditional pesticides, does not result in immediate insect kill. Instead, it interferes with the normal process of insect development and prevents the emergence of adults from larvae infesting treated accumulations of dust and litter. EZ-515876, when used according to label directions, will control the following insects:

Almond moth (<i>Ephestia cautella</i>)	Indian meal moth (<i>Plodia interpunctella</i>)
Cigarette beetle (<i>Lasioderma serricorne</i>)	Lesser grain weevil (<i>Rhyzopertha dominica</i>)
Sawtooth grain beetle (<i>Uryzodiphus surinamensis</i>)	Merchant grain beetle (<i>Oryzaephilus mercator</i>)
Red flour beetle (<i>Tribolium castaneum</i>)	Confused flour beetle (<i>Tribolium confusum</i>)
Tobacco moth (<i>Plodia interpunctella</i>)	German Cockroach (<i>Blattella germanica</i>)

PRECAUTION STATEMENTS
HAZARDOUS TO HUMANS
CAUTION

Avoid contact with eyes. In case of contact, immediately flush eyes with plenty of water. Get medical attention if irritation persists.

BEST DOCUMENT AVAILABLE

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Method, Rate and Timing of Application: Do not apply directly to food or tobacco being processed. Cover or remove any food or tobacco and processing equipment during application. After spraying, wash all equipment, benches, shelving, etc. where food or tobacco will be handled, with potable water.

Application with Mechanical Chemical Foggers: Apply EZ-515876 as a mist or aerosol. Dilute as required, following the instructions of the spray equipment manufacturer. Apply EZ-515876 at the rate of 10 ml per 10,000 square feet. Use spray equipment to achieve uniform treatment.

Close room or warehouse and shut off all ventilating systems. Apply the product as a fine mist in the air above the trays, shelves, and in areas where dust may accumulate. Leave the room closed for 30 minutes to allow spray mist to settle on accumulated food or tobacco deposits. Do not remain in treated area and ventilate before entry. Repeat application as required.

Application as An Aqueous Spray: Prepare a diluted spray solution by adding 10 milliliters of EZ-515876 to 10 gallons of water. Apply at a rate of 1 gallon of diluted solution per 1,000 square feet of surface area with any low pressure sprayer typically used for indoor applications. Treat all areas which may harbor insect pests. Repeat application as required.

To Control German Cockroaches: Prepare a diluted spray solution by adding 100 milliliters of DIAMEX to 10 gallons of water. Apply at a rate of 10 gallons of diluted solution per 10,000 square feet. Apply DIAMEX as a fine mist in cracks and crevices, along baseboards, under sinks and other areas where roaches hide.

Mixing Instructions: Partially fill the mixing container with water. Add EZ-515876, mix, and complete filling with recommended quantity of water. Use diluted spray solution within 24 hours of mixing. Agitate before each use.

WARRANTY AND CONDITIONS OF SALE

Zorcon Corporation warrants that this product conforms to the chemical description on the label and is useful for the control of insect pests in tobacco processing and food handling establishments as described herein. Neither Zorcon nor the seller makes any other warranty, express or implied.

02930

Zorcon Corporation
Agricultural Chemicals Division
Palo Alto, California 94304
NET CONTENTS: 2.5 LITERS

10-1-82

20954-101