

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS

CAUTION

Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes, or clothing. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. Wash thoroughly after handling. Do not store near or contaminate food or feedstuffs.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE
GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. STORAGE: Store in a cool place.

DISPOSAL: Bury empty container or discard according to local regulations. Never reuse empty container.

INSTRUCTIONS FOR USE

1. Partially fill spray tank with water, then add the recommended amount of ENSTAR 5E, agitate and complete filling.
2. Diluted spray solution should be used within 24 hours; always agitate before using.

WARRANTY AND CONDITIONS OF SALE

Zoecon warrants that this product conforms to the chemical description on the label, and is useful for the control of greenhouse pests as described herein. Neither Zoecon nor the seller make any other warranty, express or implied.

MFG. BY

Zoecon Corporation, 975 California Ave., Palo Alto, CA 94304

Zoecon

ENSTAR™ 5E

INSECT GROWTH REGULATOR
For Control of Whiteflies and Aphids in Greenhouses on
Ornamental Plants and Vegetable Seed Crops

ACTIVE INGREDIENTS: kinoprene
[2-propynyl (E,E)-3,7,11-trimethyl-2,4-
dodecadienoate] * 65.3%

INERT INGREDIENTS: 34.7%
TOTAL 100.0%

Formulation contains 0.625 lb. active ingredient per pint

*Patent Pending E.P.A. REG. NO. 20354-4-AA
ENSTAR is a trademark of Zoecon Corporation
E.P.A. Est. No. 20954-CA-2

Keep out of reach of children

CAUTION

See side panel for additional precautionary statements.

NET CONTENTS 1 PINT

RECOMMENDED GREENHOUSE
APPLICATIONS

RATES OF APPLICATION

Preventative Maintenance Program: Before whitefly and aphid populations reach damaging levels, mix ENSTAR 5E at the rate of 6 to 10 oz per 100 gallons of water (1 tsp/2 gal) and apply to runoff every 2 weeks. Generally, 1 oz of ENSTAR 5E per 2,000 sq ft every two weeks is required for Preventative Maintenance.

Heavy Population Control Program: To a heavy population of adult whiteflies or aphids, make 2 applications 7 days apart of ENSTAR 5E to runoff at a rate of 20 oz per 100 gallons of water (1 tsp/gal). Two weeks later, begin the Preventative Maintenance Program.

METHODS OF APPLICATION

Complete coverage of foliage is essential for control of whiteflies and aphids. For conventional spray equipment, mix the recommended amount of ENSTAR 5E and water and spray both top and bottom surfaces of leaves until runoff. For other applications, follow the equipment manufacturer's specifications.

EFFECTS ON GREENHOUSE PLANTS

Slight to moderate injury has occasionally been noted on blooms under certain conditions. For whitefly and aphid control, it is suggested that applications be made in the pre-bloom stage. Before making large scale applications to any crop, a grower should test spray small numbers of plants to determine plant safety under his own cultural conditions.

Applications of ENSTAR 5E insect growth regulator have been made to the following greenhouse plants without adverse effects to the foliage:


Azalea, Chrysanthemum, Dianthus, Fuchsia, Geranium,
Hydrangea, Marigold, Pelargonium, Petunia, Snapdragon


Application of ENSTAR 5E to certain varieties of poinsettia has produced bract damage.

Applications of ENSTAR 5E insect growth regulator have been made to the following vegetable seed crops in greenhouses without adverse effects:

Fava Bean, Green Bean, Spinach, Sugar Beet, Tomato

Seeds from treated vegetable seed crops must be used only for subsequent planting. Portions of treated plants other than seeds must be disposed of according to local regulations and must not be used for food or feedstuffs.





zoëcon

ENSTAR™ 5E

INSECT GROWTH REGULATOR

For Control of Whiteflies and Aphids in Greenhouses
on Ornamental Plants and Vegetable Seed Crops

What is ENSTAR 5E insect growth regulator?

ENSTAR 5E insect growth regulator (IGR) is a chemical analog of a natural insect hormone. Unlike traditional pesticides the presence of ENSTAR 5E does not produce an immediate insect kill, but rather it interferes with the normal process of insect development. Because ENSTAR 5E is specific for insects of the order Homoptera, it is recommended for control of whiteflies and aphids in greenhouses.

What does ENSTAR 5E IGR do?

At the preventative maintenance rate, ENSTAR 5E causes morphogenetic, cecidial, and sterilizing effects which result in a gradual reduction of insect populations to below damaging levels. Evaluation methods should allow sufficient time (two weeks) after application for the full control potential of ENSTAR 5E to be realized. At the higher rate for heavy populations, ENSTAR 5E kills adult whiteflies and aphids.

How should ENSTAR 5E IGR be used?

A preventative spray schedule along with good management practices (such as keeping doors closed and not introducing heavily infested plants into the greenhouses) will provide effective whitefly and aphid control. In the case of heavy adult populations, ENSTAR 5E should be used at the higher rate to quickly reduce the whitefly or aphid population.

TAKE TIME



OBSERVE LABEL
DIRECTIONS

— SCORE

