

**COMPREHENSIVE WINTER SPRAY**

Directions: Use 1 1/2 pints of spray and 1/4 pint of 50% DDT Spray in 6 gallons of water (7 1/2 tablespoons KORREKT Lime Sulfur Spray and 2 1/4 tablespoons DDT Spray per gallon). Keep agitated during spraying.

Roses and Ornamentals (Tormax deciduous only): Rose-rose aphid, rose thrips, brown eye, scale, black scale, overwintering spider mite and aphid eggs, many overwintering fungus spores or plant diseases.

Fruit Trees: Apples, Cherries, Nectarines, Peaches, Plums, Prunes, Quinces (Prunus deciduous only): San Jose scale, brown aphid, scale, overwintering spider mite and aphid eggs, many overwintering fungus spores or plant diseases.

**GENERAL SEASON SPRAYING**

Powdery Mildew: Roses, delphiniums, Black-eyed Susans, columbine, crepe myrtle, sweet peas, zinnias, western grapes, holly, ornamentals, tuberosa fragrant. Spray of 2 tablespoons per gallon of water on tender new growth and 1 tablespoon per gallon of water on mature foliage. Apply every 1 to 2 weeks.

**LIME AND SOIL TREATMENTS**

Lime: Gardens. To condition alkaline lawn and garden soils and improve water penetration on moisture retention in heavy soils, apply 1 pint in 7 1/2 to 15 gallons of water on 500 sq ft of area. Apply with sprayer or watering can. After application, water lawn immediately. Apply every 60 days throughout spring and summer seasons as needed.

Soil Correction in Acid-Loving Plants: Rhododendrons, Azaleas, Camellias. Spray of 1 tablespoon per gallon of water. Apply 1 1/2 gallons diluted material per plant's surface of soil around plant. Repeat treatment every 2 to 3 months until plants regain proper growth.

NOTICE: Seller warrants that this product consists of the ingredients specified and is recommended for use as directed on the label. Seller is authorized to make any other warranty, guarantee, or direction concerning this product. Because the time, place, rate of application, and other conditions of use are beyond Seller's control, Seller's liability from the handling, storage, and use of this product is limited to replacement of product or refund of purchase price.



BEST PRODUCTS  
P.O. Box 267, Lathrop, Ore. 97130  
P.O. Box 5356, Houston, Tex. 77212  
Division of Zinco Corp.  
Label No. 979  
EPA Reg. No. 27954-E-7  
EPA Est. No. 20054-CA-1



**BEST**  
**Korrek**

**LIME-SULFUR SPRAY**

For growing season and dormant spraying to control certain plant diseases. Controls Rose powdery mildew and Peach leaf curl.



ACTIVE INGREDIENT	Likum Disulfide	29.0%
INERT INGREDIENTS		71.0%
		TOTAL
		100.0%
Net Weight	1.00 lb. per gallon of AA-1	10.0 lb.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

SEE BACK PANEL FOR ADDITIONAL CAUTIONS

**Net Contents: 1 Pint (473 ml)**

READ ENTIRE LABEL USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS AND DIRECTIONS.

**PRECAUTIONARY STATEMENTS - HAZARDS TO HUMANS  
CAUTION**

KEEP OUT OF REACH OF CHILDREN: Avoid contact with skin. May cause skin irritation. Wash hands after use. If material gets into eyes, immediately flush with water. If eye irritation persists, contact doctor at once.

**ENVIRONMENTAL HAZARDS**

Pick open flowers before application to avoid discoloration. Avoid spraying on painted surfaces to avoid staining. However, if accidentally sprayed on painted surfaces, considerable bleaching out will occur. If dust or dirt on surface, wash with a stick. Crystallization will not impair product quality. Do not use 30 days before or after using an oil spray during growing season or on plants in leaf. Lime sulfur should not be used within 10 to 14 days of an oil spray of any type.

**DIRECTIONS FOR USE**

If in a violation of Federal law to use this product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**

KEEP CONTAINER TIGHTLY SEALED BEFORE STORAGE. Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Do not reuse empty container. Wrap container and put in trash collection.

For best results, spray to cover all exposed plant parts and repeat as necessary. Do not spray when temperature is over 80°F. All spraying should be done early in the morning to avoid burning.

AGRICULTURAL CAUTION: Do not use on crops or plants if they are susceptible to sulfur injury. Do not use on evergreen plants. Do not apply in freezing weather.

**DORMANT SPRAYING**

Roses: Rose Rust, Black Spot, Prune and leafy off old leaves. Spray bushes of 10 tablespoons (5 oz.) per gallon of water.

Peaches: Peach Leaf Curl, peach blight. Spray at 10 tablespoons (5 oz.) per gallon of water in late fall before heavy rains, also before buds begin to swell in late winter.

Pears: Pear Leaf Rust. Spray at 10 tablespoons (5 oz.) per gallon of water just as buds begin to swell in spring and late fall before leaves start to turn color.

Caneberries (Blackberries, Raspberries, Rubus spp.): Leaf Spot, Cane Rust, Red Berry Trouble (Recognized by Heart Berries). Spray at 10 tablespoons (5 oz.) per gallon of water in late fall, followed by another treatment in spring as buds begin to swell.

ACCEPTED  
JUN 9 1980  
Under the Federal Insecticide, Fungicide, and Rodenticide Act, this product is hereby certified for sale and use in the State of Texas.  
EPA Reg. No. 27954-E-7