

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES, flush with plenty of water for 15 minutes and get medical attention

IF SWALLOWED, drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger or blunt object. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention

IF INHALED, remove patient from contaminated area and get medical attention

IF ON SKIN, remove contaminated clothing and wash skin with soap and water

WARRANTY

EM INDUSTRIES warrants that this product (1) conforms to the ingredient statement on the label and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. EM INDUSTRIES MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or in combination with other materials.

Net Contents 1 U.S. Gallon

FUNGINEX

EMULSIFIABLE CONCENTRATE

ACC
MAR 6 1981
Use for the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 21137-4

ACTIVE INGREDIENT:	
Triforine (N,N-(1,4-piperazinediyl)bis (2,2,2-trichloroethylidene) bis (formamide))	18.2%
INERT INGREDIENTS	81.8%
TOTAL	100.0%

THIS PRODUCT CONTAINS 1.1 LBS OF TRIFORINE PER GALLON

DANGER KEEP OUT OF REACH OF CHILDREN
See Side Panels For Additional Precautions



Manufactured for
EM INDUSTRIES, INC.
Plant Protection Division
5 Skyles Drive
Highland Park, N.J. 07624

EPA Registration Number 21137-4
EPA Establishment Number 279 FL 1
FUNGINEX is a product and trademark of
CELAMERCK S.A.B.M. & Co., K.G. Ingelheim, Germany
Aquate (CEL) GMBH, Ingelheim and E. MERCK, Darmstadt, Germany

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND
DOMESTIC ANIMALS**

DANGER

Corrosive, causes eye damage. Do not get in eyes. Wear goggles or face shield. Harmful if swallowed. Do not inhale spray mist. Avoid contact with skin. After handling, wash thoroughly with soap and water.

ENVIRONMENTAL HAZARDS

Keep out of lakes, ponds and streams. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

STORAGE AND DISPOSAL

Store in a cool, dry place. Do not store below 32° F. Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not re-use empty container.

PESTICIDE DISPOSAL Pesticide or rinse liquid that cannot be used or chemically reprocessed, should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

CONTAINER DISPOSAL Dispose of in an incinerator or landfill approved for pesticide containers or bury in a safe place. Consult federal, state or local disposal authorities for approved alternative procedures such as limited open burning.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PEACHES:

BROWN ROT BLOSSOM BLIGHT (*Monilinia fructicola* and *M. laxa*)

Apply 12 to 16 fl. oz. of FUNGINEX per 100 gallons of water or 36 to 48 fl. oz. per acre dilute. Make 2 or 3 applications depending upon the length of the bloom period and conditions favoring brown rot development.

Apply the first spray at pink bud (blossom buds just before opening) and repeat at 75-100% bloom. Make a third application at petal fall. If warm, wet weather prevails, apply 2 to 4 day intervals.

In California Only

Apply two sprays of FUNGINEX at 12 fl. oz. per 100 gallons of water or 36 fl. oz. per acre dilute. Make the first application at pink bud, followed by a second application at full bloom.

HIGHBUSH BLUEBERRIES:

MUMMYBERRY DISEASE (*Monilinia vaccinii-corymbosi*)

Pacific and Mid Western States:

Apply 24 fl. oz. of FUNGINEX per acre in 20 to 50 gallons of water for ground application or in 5 gallons of water for aerial application.

Make the first application at leaf bud break and repeat in 7 to 10 days. Make the third application at pink bud stage and repeat in 7 to 10 days at early bloom.

For the last application, apply 16 fl. oz. of FUNGINEX per acre in 20 to 50 gallons of

water for ground application or in 5 gallons of water for aerial application.

Make the last application between full bloom and early petal fall.

Do not make more than five applications from leaf bud break to early petal fall.

Eastern Seaboard States (For primary infection only)

Apply 24 fl. oz. of FUNGINEX per acre in 20 to 50 gallons of water for ground application or in 5 gallons of water for aerial application.

Make the first application at leaf bud break and repeat in 7 to 10 days. Make the last application at pink bud stage.

Do not make more than three applications from leaf bud break to pink bud stage. Application of FUNGINEX during or beyond early bloom may result in fruit russetting.

USE PRECAUTIONS

Do not graze animals in treated orchards.

As timing of fungicide applications for disease control vary due to climatic and other conditions, consult official local or state recommendations.

Do not use this material if it cannot be applied according to the use patterns on this label.

Do not mix FUNGINEX with wetting agents, spreader-stickers, or other adjuvants.

Do not let spray mixture stand in tank overnight.

Printed in USA 1980

21137-4

75

76