

7

21137-11

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS  
AND DOMESTIC ANIMALS**

**CAUTION**

Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing dust or spray mist. Avoid contact with eyes or clothing. If swallowed, drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger or blunt object. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention. In case of contact immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists.

**STORAGE AND DISPOSAL**

Store in a cool, dry place.

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse container.

**PESTICIDE DISPOSAL:** Pesticide or rinse liquid that cannot be used or chemically reprocessed, should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

**CONTAINER DISPOSAL:** Dispose of in an incinerator or landfill approved for pesticide containers or bury in a safe place. Consult federal, state, or local disposal authorities for approved alternative procedures such as limited open burning.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**STONEFRUIT:** APRICOTS, CHERRIES, NECTARINES, PEACHES, PLUMS (FRESH PRUNES)

DO NOT USE ON VARIETIES THAT WILL BE DRIED

**BROWN ROT POST-HARVEST** (*Monilinia fructicola* and *M. laxa*)

In California Only -

**DIP OR HYDROCOOLER:** Dip fruit thoroughly for at least 30 seconds but not more than 60 seconds as soon as possible after harvest. Use 1 lb FUNGINEX WP per 100 gallons of water. Treat 200,000 lb of fruit with 100 gallons of Funginex solution. Replenish dip solution to maintain adequate dip level. Drain and flush system daily.

**SPRAY:** Spray fruit as soon as possible after harvest. Use 1 lb Funginex WP per 100 gallons of water. Treat 200,000 lb of fruit with 100 gallons of Funginex solution. Do not recirculate used spray solution.

Fruit may be treated with suitable post-harvest wax for shrinkage control.

**WAX:** Use 1 lb Funginex WP per 100 gallons of wax/water mixture. Use one part wax per seven parts water or follow manufacturers recommendations. Treat 200,000 lb of fruit with 100 gallons of Funginex-Wax solution. Do not reuse run-off solution from brushes or spray.

**WARRANTY:** EM Industries warrants that this product, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT SO WARRANTED, THE PRODUCT IS SOLD AS IS. EM INDUSTRIES MAKES NO OTHER WARRANTY EXPRESSED OR IMPLIED.

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or in combination with other materials.

# FUNGINEX®

WETTABLE POWDER

**Active Ingredient:**

Triforine (N,N'-[1,4-piperazinediyl]bis (2,2,2-trichloroethylidene)) bis (formamide)..... 50 %

Inert Ingredients:..... 50 %

Total: 100%

## CAUTION

### KEEP OUT OF REACH OF CHILDREN

See Side Panel For Additional Precautions

Manufactured for:



**INDUSTRIES, INC.**

Plant Protection Division  
5 Skyline Drive  
Hawthorne, New York 10532

Funginex® is a product and trademark of  
CELANERCK GMBH & CO., KG.,  
Ingelheim, Germany

EPA Registration Number: 21137-11  
EPA Establishment Number: 6967-WG-3

Net Contents: 2 LBS.

12/80