

32
oz.
30
oz.
28
oz.
26
oz.
24
oz.
22
oz.
20
oz.
18
oz.
16
oz.
14
oz.
12
oz.
10
oz.
8
oz.
6
oz.
4
oz.
2
oz.

HY-GUARD "100"

DIRECTIONS:

HOW MUCH TO USE:
When Algae Is Not Visible. In freshly filled pools or pools showing no visible algae growth, add one quart **HY-GUARD "100"** per 12,500 gallons of water.

When Algae Is Visible. If algae growth is visible, add 2.5 quarts **HY-GUARD "100"** per 12,500 gallons of water to kill and control algae species most commonly found in swimming pools. If algae growth is just visible the initial dose must be 2.5 quarts per 12,500 gallons, followed by cleaning out settled algae and applying 1 quart per 12,500 gallons of water.

MAINTENANCE DOSAGE
 Add 1 quart **HY-GUARD "100"** every 5 to 7 days per 12,500 gallons of water. If warm weather prevails or pool has unusually heavy use, maintenance dosage may have to be repeated.

HOW TO USE:
 Mix required amount of **HY-GUARD "100"** into a bucket of water and distribute evenly throughout pool water.

THIS PRODUCT IS TOXIC TO FISH. Treated effluent should not be discharged where it will drain into lakes, streams, ponds or public water.



HY-GUARD "100" ALGAE CONTROL FOR SWIMMING POOLS

**HELPS KEEP POOL WATER CLEAR
AND FREE OF GROWING ALGAE**

ACTIVE INGREDIENTS
 n-Alkyl (50% C₁₄, 40% C₁₂, 10% C₆) dimethyl
 benzyl ammonium chloride 10%
INERT INGREDIENTS 90%

DANGER
KEEP OUT OF REACH OF CHILDREN
 Rinse empty container thoroughly with water and discard it. See antidote statement and other required statements on the side panels.

NET CONTENTS: 32 FL. OZ. (1 U.S. QT.)
 EPA Registration No. 21144-E

HY-GUARD "100"

DANGER:

Keep out of reach of children. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, **PROMPTLY** drink a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

**MANUFACTURED BY
HYPO CHEMICAL CORP
MILWAUKEE, WIS. 53225**

ACCEPTED
 APR 27 1973