91975:I:Pringle:P-3:KENCO:11/3/87:11/16/87:rw:VO:EK:aw

PREVIOUS EDITION MAY BE USED UNTIL SUPPLY IS EXHAUSTED.

ACCEPTED
with COMMEXTS
in EPA Latter Details

RLCA-W12-0

NOV1 2 1987

Green the Feneral Insecticide, Fungi Te, and Rodenticide Act of for the posticide and EPA Reg. No. 2/164-10

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid breathing vapors. Avoid contact with skin, eyes, or clothing.

FIRST AID: If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Keep out of lakes, streams, or ponds. Permits may be required for discharges containing this pesticide into fakes, streams, ponds, or public water. For guidance, contact the regional office of the Environmental Protection Agency.

DIRECTIONS FOR

It is a violation of Federal law to use this proc
its labeling.
RLCA-W12-0 is
and fungi in recirculating commercial and indu
to its use, systems must be cleaned to remo
slime, and other deposits. An initial slug addit
fluid ounces of RLCA-W12-0
of water to provide a concentration of40
RLCA-W12-O, b
in the system, is recommended. Repeat initial
Subsequent slug additions of1.20
RLCA-W12-0 per 1000
100 parts per million ofRLCA-W12-(
employed every 2 to 5 days, or as needed. Th
upon the relative amount of bleedoff and the se
lem. Slug additions should be made in the sur
RLCA-W12-Ois use
air-washing systems that maintain effective m
to its use, systems should be cleaned to re
deposits. An initial slug dose of21.37
RLCA-W12-0 pe
mended. Repeat initial dosage until control is
tions of14_44 to35_63 fluid ounces of _
per 1000 gallons of water should be employed e
frequency of addition depends upon the relativ
of the bacterial problem. Slug additions may be
collection trays of the airwash system.

STORAGE AND DI

On not contaminate water, food, or feed by storage or disposal. STORAGE: Do not stack more than four drums high. Leaking of drums for disposal. Spills should be absorbed in sawdust or sand tainer closed when not in use.

PESTICIDE DISPOSAE: Wastes resulting from the use of this proved waste disposal facility.

CONTAINER DISPOSAL

- PLASTIC Triple rinse for equivalent! Then offer for recycling of sandary landfill, by incineration, or, if allowed by state and loc smoke.
- METAL. Triple ruse for equivalent. Then offer for recycling of samtary landfill, or try other procedures approved by state and

		lanufactured
_	RIO LINDA CHEMI	CAL CO.,
_	410 NORTH 10th	
_	SACRAMENTO, CA	95814
EPA REC	NO	EP

NET CONTENTS .