

### PRECAUTIONS

When exposed to Larvacide 100 vapors, use full face gas mask with canister designed for acid gases and organic vapors.  
 Do not fumigate valuable planting seed. Always aerate thoroughly after exposure period. Do not direct spray toward open flame.  
 Larvacide 100 is toxic to animals and plants. Remove animals, birds, plants, bulbs from any exposure to fumes. Do not fumigate such materials as fresh fruits, fresh vegetables, fresh meat, yeast, eggs, butter or cheese. Larvacide 100 may cause corrosion on some metals. Avoid fumigating in humid atmosphere.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### SPACE FUMIGATION IN EMPTY MILLS, WAREHOUSES, EMPTY STORAGE BUILDINGS, VAULTS:

- Remove all processed food or feeds such as flour, mixed feed or processed food products.
- Seal all openings as carefully as possible using tape and polyethylene sheets or other convenient material. Distribute empty, burlap sacks on the floor and sprinkle Larvacide over them. Use gas mask.

**"DO NOT MOVE TRUCKS, VANS, OR TRAILERS DURING FUMIGATION. THEY MUST BE COMPLETELY AERATED BEFORE MOVEMENT IS ALLOWED."**

Fumigated areas must be placarded on all entrances with signs containing at least the signal word **DANGER** and the "Skull and Crossbones" and the words "Area Under Fumigation. Do Not Enter Until Fully Aerated" the date of fumigation, name of the fumigant used, emergency telephone number for contact, and the name and address of the fumigator. Do not remove warning signs until the fumigated area is completely aerated and safe for entry as indicated by a suitable detector.

### STORAGE AND DISPOSAL

Store in dry place.

Return empty cylinder to manufacturer.

Triple rinse (or equivalent) and dispose of in a sanitary landfill or by other approved State and local procedures.

#### DIRECTIONS FOR EMPTYING CYLINDER

**CAUTION** Whiplash most violent when cylinder becomes empty. To avoid tear gas and over-ence in area, close valve as soon as gas is exhausted and tubing begins to whip.

It controls insects like rice weevils, granary weevil, confused flour beetle, coffee berry borer, grain borer, saw-toothed grain beetle, flour moths and grain moths. It also controls rats and mice and some mold causing fungi.

**ACCEPTED**

JUL 29 1981

21327-4

### RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's application.

# Larvacide

100 • CHLOROPICRIN

## GRAIN FUMIGANT SPACE FUMIGANT SOIL FUMIGANT

Active Ingredient Chloropicrin 99%  
Inert Ingredients.....1%

This product contains 13.75 lbs. of Chloropicrin per gallon.

KEEP OUT OF REACH OF CHILDREN

**DANGER**  **POISON**

#### STATEMENT OF PRACTICAL TREATMENT

- IF SWALLOWED: Call a physician or Poison Control Center immediately. If possible, vomiting should be induced under medical supervision. Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Remove victim to fresh air. Apply artificial respiration if indicated.
- IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated.
- IF ON SKIN: Remove contaminated clothing and wash affected areas with soap and water.
- IF IN EYES: Flush eyes with plenty of water. Call a physician immediately.

Mfg. by:



411 S. PARKER  
AMARILLO, TEXAS 79106

NET CONTENTS

EPA No. 21327-00004


EPA Establishment No. 21327-1X-1

**RESTRICTED USE PESTICIDE**  
 Available for use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's application.

**Larvacide**  
 100 • CHLOROPICRIN

**GRAIN FUMIGANT**  
**SPACE FUMIGANT**  
**SOIL FUMIGANT**

Active Ingredient Chloropicrin 99%  
 Inert Ingredients.....1%  
 This product contains 13.75 lbs.  
 of Chloropicrin per gallon.

**OUT OF REACH OF CHILDREN**  
**DANGER**  **POISON**

**STATEMENT OF PRACTICAL TREATMENT**  
**REMEDY:** Call a physician or Poison Control Center immediately. If possible, vomiting should be induced under medical supervision. Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Remove victim to fresh air. Apply artificial respiration if indicated.  
**REMARKS:** Remove contaminated clothing and wash affected areas with soap and water. Flush eyes with plenty of water. Call a physician immediately.

Mfg. by:  
 **411 S. PARKER**  
**AMARILLO, TEXAS 79106**

NET CONTENTS

21327 0004 EPA Establishment No. 21327 TX 1

**TREATS HOT SPOTS**  
 For control of insects, rodents and fungi in grain or in grain bins. For control of fungi, insects, nematodes and weeds in soil.

**CONTROLS**  
**INSECTS,**  
**RODENTS,**  
**FUNGI,**  
**IN GRAIN**  
**OR IN**  
**GRAIN**  
**BINS**

**GENERAL INFORMATION**  
**Larvacide 100** is a fumigant for control of insects, rodents, and fungi in grain and grain bins.  
**Larvacide 100** is a fumigant for control of insects, rodents, and fungi in grain and grain bins. It is also effective against weeds in soil.  
**Larvacide 100** is a fumigant for control of insects, rodents, and fungi in grain and grain bins. It is also effective against weeds in soil.

**DOSAGE IN**  
**Warehouses** 2 lbs. per 1000 cu. ft. of space.  
**Vaults** 2 lbs. per 1000 cu. ft. of space.  
**EXPOSURE PERIOD** 48 hours.  
**GRAIN TREATMENT** Apply Larvacide 100 to grain at a rate of 2 lbs. per 1000 bu. of grain.  
**DOSAGES FOR CLEAN DRY GRAIN:**  
 Wheat 2 lbs. per 1000 bu.  
 Rice 2 lbs. per 1000 bu.  
 Corn 3 lbs. per 1000 bu.  
 Soybeans 4 lbs. per 1000 bu.  
**Seal treated bin.** Exposure period at least 72 hours.

**AERATE THOROUGHLY BEFORE UNLOADING BINS**  
**SOIL FUMIGATION:** Apply Larvacide 100 to soil at a rate of 2 lbs. per 1000 sq. ft. of area.  
**DOSAGE:** Normal dosage is 2 lbs. per 1000 cu. ft. of space.  
**EXPOSURE PERIOD:** 48 hours.  
**AERATION:** Aerate grain thoroughly before unloading bins.  
**NON WARRANTY:** This product is sold as is, without warranty.