

DANGER:

Keep Out of Reach of Children. Corrosive. Causes eye damage and skin irritation. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

FIRST AID:

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution; or if these are not available, drink large quantities of water. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

USE DIRECTIONS FOR DISINFECTION

Disinfection in Hospitals, Nursing Homes and Other Health Care Institutions

For disinfecting floors, walls, countertops, bathing areas, lavatories, bedframes, tables, chairs, garbage pails and other hard surfaces.

Add 1 1/2 oz. Sharp Clean L 55 to 4 gallons water. Apply to previously cleaned hard surface with mop or cloth.

At this use level, Sharp Clean L 55 is effective against *Pseudomonas aeruginosa*.

Disinfectant in Institutions, Industry, Schools and Homes

For disinfecting floors, walls, bedframes, countertops, tables, chairs, garbage pails, bathroom fixtures and other hard surfaces.

SHARP CLEAN L55

Disinfectant Sanitizer-Fungicide-
Deodorizer for Hospital, Institutional,
Industrial, School, Dairy and Other
Farm and Home Use
A.O.A.C. Phenol Coefficients

Staphylococcus aureus, ATCC #6538	77.5
Salmonella typhosa, ATCC #6539	77.5

Active Ingredient

Didecyl dimethyl ammonium chloride 7.5%

Inert Ingredients 92.5%

100.0%

EPA Registration No. 22058-6

EPA EST. No. 22058-Tx-1

ACCEPTED

JUN 29 1976

the Federal Insecticide

22058-6

MANUFACTURED BY:

SHARP CHEMICAL CO.

5921 PLAINVIEW

HOUSTON, TEXAS 77017

DANGER:

KEEP OUT OF REACH OF CHILDREN. SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

Disinfection of Barber Tools

Barber tools (such as combs, brushes, razors, and scissors) can be disinfected by immersing in a 1/2 oz/gallon solution of Sharp Clean L 55.

Disinfection of Poultry Equipment, Animal Quarters and Kennels

Poultry brooders, watering founts, feeding equipment and other animal quarters (such as stalls and kennel areas) can be disinfected after thorough cleaning by applying a solution of 2 oz. Sharp Clean L 55 in 4 gallons of water with mop, cloth or brush. Small utensils should be immersed in this solution.

Prior to disinfection, all poultry, other animals and their feeds must be removed from the premises. This includes emptying all troughs, racks and other feeding and watering appliances. Remove all litter and droppings from floors, walls and other surfaces occupied or traversed by poultry or other animals.

After disinfection, ventilate buildings, coops and

USE DIRECTIONS FOR SANITIZING

Sanitizing of Food Processing Equipment and other Hard Surfaces in Food Contact Locations

For sanitizing food processing equipment, dairy equipment, food utensils, dishes, silverware, glasses, sink tops, counter tops, refrigerated storage and display equipment and other hard surfaces.

Wash and rinse all articles thoroughly, then apply a solution of 1 oz. Sharp Clean L 55 in 4 gallons of water.

Apply to sink tops, countertops, refrigerated storage and display equipment and other stationary hard surfaces by cloth or brush.

Dishes, silverware, glasses, cooking utensils and other similar size food processing equipment can be sanitized by immersion in a 1 oz./4 gallon dilution of Sharp Clean L 55.

Food contact surfaces must be rinsed with potable water before use.

At 1 oz./4 gallons, Sharp Clean L 55 fulfills the criteria of Appendix F of the Grade A Pasteurized Milk Ordinance 1965 Recommendations of the U.S. Public Health Service in waters up to 1000 ppm of hardness calculated as

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution; or if these are not available, drink large quantities of water. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

(front panel)

USE DIRECTIONS FOR DISINFECTION

Disinfection in Hospitals, Nursing Homes and Other Health Care Institutions

For disinfecting floors, walls, countertops, bathing areas, lavatories, bedframes, tables, chairs, garbage pails and other hard surfaces

Add 3 1/2 oz. Sharp Clean L 55 to 4 gallons water. Apply to previously cleaned hard surface with mop or cloth.

At this use level, Sharp Clean L 55 is effective against *Pseudomonas aeruginosa*.

Disinfectant in Institutions, Industry, Schools and Homes

For disinfecting floors, walls, bedframes, counter tops, tables, chairs, garbage pails, bathroom fixtures and other hard surfaces

Add 2 oz. of Sharp Clean L 55 to 4 gallons of water. Apply to previously cleaned hard surface with mop or cloth.

At 2 oz./4 gallon use level, Sharp Clean is effective against *Staphylococcus aureus*, *Salmonella choleraesuis* and *Trichophyton interdigitale* (the athlete's foot fungus)

NET CONTENTS: 1 gallon

Sharp Clean L 55

Active Ingredient

Didecyl dimethyl ammonium chloride

7.5%

Inert Ingredients

92.5%

100.0%

EPA Registration No. 22058-6

EPA EST. No. 22058-Tx-1

ACCEPTED
 JUN 20 1976
 The Federal Insecticide,
 Fungicide, and
 Rodenticide Act
 22058-6

MANUFACTURED BY:
SHARP CHEMICAL, CO.
 5921 PLAINVIEW
 HOUSTON, TEXAS 77017

DANGER:

KEEP OUT OF REACH OF CHILDREN. SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

Disinfection of Barber Tools

Barber tools (such as combs, brushes, razors, and scissors) can be disinfected by immersing in a 1/2 oz./gallon solution of Sharp Clean L 55.

Disinfection of Poultry Equipment, Animal Quarters and Kennels

Poultry brooders, watering founts, feeding equipment and other animal quarters (such as stalls and kennel areas) can be disinfected after thorough cleaning by applying a solution of 2 oz. Sharp Clean in 4 gallons of water with mop, cloth or brush. Small utensils should be immersed in this solution.

Prior to disinfection, all poultry, other animals and their feeds must be removed from the premises. This includes emptying all troughs, racks and other feeding and watering appliances. Remove all litter and droppings from floors, walls and other surfaces occupied or traversed by poultry or other animals.

After disinfection, ventilate buildings, coops and other closed spaces. Do not house poultry or other animals, or employ equipment until treatment has been absorbed, set or dried.

All treated equipment that will contact feed or drinking water must be rinsed with potable water before reuse.

Rinse empty container thoroughly with water and discard.

USE DIRECTIONS FOR SANITIZING

Sanitizing of Food Processing Equipment and other Hard Surfaces in Food Contact Locations

For sanitizing food processing equipment, dairy equipment, food utensils, dishes, silverware, glasses, sink tops, counter tops, refrigerated storage and display equipment and other hard surfaces.

Wash and rinse all articles thoroughly, then apply a solution of 1 oz. Sharp Clean L 55 in 4 gallons of water.

Apply to sink tops, countertops, refrigerated storage and display equipment and other stationary hard surfaces by cloth or brush.

Dishes, silverware, glasses, cooking utensils and other similar size food processing equipment can be sanitized by immersion in a 1 oz./4 gallon dilution of Sharp Clean L 55.

Food contact surfaces must be rinsed with potable water before use.

At 1 oz./4 gallons, Sharp Clean L 55 fulfills the criteria of Appendix F of the Grade A Pasteurized Milk Ordinance 1965 Recommendations of the U.S. Public Health Service in waters up to 1000 ppm of hardness calculated as CaCO₃ when evaluated by the AOAC Germicidal and Detergent Sanitizer Method against *Escherichia coli* and *Staphylococcus aureus*.

The udders, flanks, and teats of dairy cows can be sanitized by washing with a solution of 1 oz. Sharp Clean L 55 in 4 gallons of warm water.

Use a fresh towel for each cow. Avoid contamination of sanitizing solution by dirt and soil. Do not dip used towel back into sanitizing solution. When solution becomes visibly dirty, discard and provide fresh solution.