

NET CONTENTS 5 GALLONS

CAUTION: Some fruits and other plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur on any crop unless local use has proven that sulfur is safe in that locality.

During periods of high temperature sulfur may burn foliage and fruit. Do not make sulfur applications at excessively high temperatures. Do not use THAT_® FLOWABLE SULFUR with oil or within 4 weeks of an oil application.

When THAT_® FLOWABLE SULFUR is used with Azeonate_®, lime should be added to prevent plant injury.

DIRECTIONS FOR MIXING BEFORE USING, STIR UNTIL SMOOTH

Pour recommended amount in partially filled spray tank. Keep agitator running during filling and spraying operation. **DO NOT ALLOW MIXTURE TO STAND.** Failure to maintain agitation will cause THAT_® FLOWABLE SULFUR to settle and may necessitate manual stirring to redispense.

The strong adhesive properties of THAT_® FLOWABLE SULFUR acts as a sticker on the plant, and the sticking characteristic necessitates the flushing of equipment with water after each day's use. Sulfur in any form is corrosive material. **TO REDUCE THE EFFECT, EQUIPMENT SHOULD BE FLUSHED DAILY.**

RECOMMENDATIONS

Use of this material in a manner other than in accordance with this label may cause serious plant, crop, or personal injury, excessive residue, unsatisfactory control or other unintended consequences.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty or any other warranty of merchantability of fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions not reasonably foreseeable to the seller, and buyer assumes the risk of any such use.

Consult your State Agricultural Experiment Station or State Extension Service Specialist for complete details on the spraying program best suited to your local conditions.

EPA Reg. No. 22534-2

That FLOWABLE SULFUR

ACTIVE INGREDIENT

Sulfur 52%

INERT INGREDIENTS

48%


Total 100%

(Contains 6.0 pounds Sulfur per gallon)

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid breathing spray mist. May cause irritation of nose, throat or skin. Avoid contact with eyes, skin and clothing. Do not contaminate feed and foodstuffs. Do not apply to any area not specified on label. Do not reuse empty container. Wash thoroughly with water and detergent. Puncture, make unusable. Dispose of in a safe place.



MANUFACTURED BY
STOLLER CHEMICAL COMPANY, INC.
8705 KATY FREEWAY
HOUSTON, TEXAS 77024

Unless otherwise specified for specific tree fruits, dosage rates are given in parts of THAT_® FLOWABLE SULFUR per 100 gallons water for use in the average spray.

APPLE, PEACH, PLUM

Thorough through foliage sprays 1½ to 3½ pints
Cover sprays 1 to 2 pints

CHERRY, MANDARIN

Thorough through foliage sprays 2½ to 3 pints
Cover sprays 2 pints

PEACH, PEAR, MANDARIN, BROWN ROT, SCAB

Link and bloom sprays 1½ to 3½ pints
Crust fall, crack and cover sprays 1 to 1½ pints

CHERRY, PLUM, PRUNE, BROWN ROT, LEAFSPOT

Link and bloom sprays 1½ to 1¾ pints
Crust fall, crack and cover sprays 2/3 to 1½ pints

CHERRY, GRAPEFRUIT, LEMONS, LIMES, ORANGES,

LEMONS, and LARGEMAN
Crust fall 1 to 4 pints

Do not apply within 4 weeks of an oil application.

PEAR, PLUM, PEACH

Apply 2 to 4 pints per acre in sufficient water for thorough coverage. Begin application approximately 6 weeks after planting. Repeat at 7 to 10 day intervals as long as rust control is required. Do not apply within 4 weeks of an oil application.

CHERRY, PLUM, PEACH

2 pints for early season control and up to 1 gallon for mid to late season control.

CHERRY, PLUM, PEACH, BROWN ROT, LEAFSPOT

Use 1 to 2 pints per 100 gallons water, maximum of 8 pints THAT_® FLOWABLE SULFUR per acre in spring and summer. Start when shoots are 6 to 8 inches long, and repeat at 3 week intervals.

For varying rates and number of applications, consult your State Agricultural Service or State Extension Experiment Station in your report for specific information. Follow the USE PRECAUTIONS given on this label.

PEACH, PLUM, PEACH

Use 3 to 4 pints in 8 gallons water per acre by air. Apply at the first sign of infection and repeat at 3 week intervals.

CHERRY, PLUM, PEACH

Use 1 pint. Apply at first sign of infection and repeat at 3 week intervals. For strawberries that will be canned, discontinue application of sulfur well before harvest in accordance with local commercial requirements.

PEACH, PLUM, PEACH

Use 1 to 1 gallon per acre depending upon size of plants. Use before bloom spray.