

FOLIAR SPRAY RECOMMENDATIONS

PEANUTS

To control rust and leaf spot, use 2 to 3 quarts in at least 10 gallons of water to maintain good coverage. Begin applying the above mixture 40 to 60 days after planting. Maintain a schedule of 10 to 14 days through the growing season.

The following recommendations are made on the basis of product per 100 gallons of water. These recommendations are for sprays to be applied by conventional spray equipment, or concentrated sprayers, consult the manufacturers' recommendations.

CITRUS:

Brown Rot - use 1 gallon; do not add a spreader. Apply at the first indication of rain or first appearance of Brown Rot. Repeat as needed to protect during wet weather. Melanose - use 3 gallons. Apply 1 to 3 weeks after petal fall. Scab - use 3 gallons. Apply two sprays, use before trees begin to flush and at 2/3 petal fall.

Caution: Do not apply in areas where copper injury is known to occur.

Florida Only: Melanose - use 1 1/2 gallons. Apply 1 to 3 weeks after petal fall. Repeat in 4 weeks if there has been a history of the disease, a wet spring or late or scattered bloom.

PEARS:

Fire Blight: use 0.5 gallons in 400 gallons per acre to give 1/2 lb. metallic copper per acre. Apply at 10% bloom and continue every 5 to 7 days throughout bloom. Repeat especially after showers or rain. Use only where State Disease Control officials recommend such use on specific varieties.

ALMONDS:

Brown Rot and Shot Hole - use 0.5 to 1.0 gallons. As a dormant spray for control of Shot Hole Disease apply Dec. 15 to Jan. 15. Bud sprays for control of Brown Rot should be applied through swelling bud stage to popcorn stage. Do not apply after 50% bloom or when trees are in leaf.

APRICOTS:

Brown Rot and Shot Hole - use 1 to 1.5 gallons. As a dormant spray for the control of Shot Hole disease apply Nov. 15 to Jan. 15. Bud sprays for control of Brown Rot should be applied through swelling bud stage to popcorn stage. Do not spray when trees are in leaf.

For the following crops begin application when the disease normally appears and repeat at 7 to 10 day intervals unless stated otherwise.

SUGAR BEETS:

Cercospora Leaf Spot - use 4 gallons.

BEETS:

Downy Mildew - use 3 gallons.

BROCCOLI, CABBAGE, CAULIFLOWER:

Downy Mildew, Leaf Spot - use 3 gallons.



**TOP COP
with sulfur**

ACTIVE INGREDIENTS:

Sulfur as S 50.0%
Copper as Metallic 4.4%

INERT INGREDIENTS 45.6%


TOTAL 100.0%

Derived from Elemental Sulfur and Tribasic Copper Sulfate, 8.4%

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid breathing spray mist. May cause irritation of nose, throat or skin. Avoid contact with eyes, skin and clothing. Do not contaminate feed and foodstuffs. Do not apply to any area not specified on label. Do not re-use empty container. Wash thoroughly with water and detergent. Puncture, make unusable. Dispose of in a safe place.



NET WEIGHT OF CONTENTS: 55 POUNDS

MANUFACTURED BY:

STOLLER CHEMICAL COMPANY, INC.
8705 KATY FREEWAY, SUITE 400
HOUSTON, TEXAS 77024

CARROTS:

Alternaria Leaf Blight, Cercospora Blight, Downy Mildew - use 3 gallons.

EGGPLANTS:

Alternaria Blight, Anthracnose, Phomopsis - use 3 gallons.

PEPPERS:

Anthracnose, Bacterial Spot, Cercospora Leaf Spot - use 3 gallons.

POTATOES:

Early and Late Blight - use 4 gallons.

SPINACH:

Downy Mildew (Blue Mold), White Rust - use 3 gallons.

CAUTION

Some plants are susceptible to injury from sulfur under climatic conditions. The user is advised not to use sulfur on a crop unless local use has proven that sulfur is safe.

During periods of high temperature, sulfur may injure fruit. Do not make sulfur applications at excessive temperatures. Do not use Top Cop with Sulfur with oil or oil application.

This product is toxic to fish. Keep out of lakes, streams, and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS FOR MIXING

BEFORE USING, STIR UNTIL SMOOTH

Pour recommended amount into partially filled sprayer with agitator running during filling and spraying operation. Allow mixture to stand. Failure to maintain agitation will result in sulfur settling and may necessitate manual stirring.

The strong adhesive properties of Top Cop will stick to the plant, and the sticking characteristic of sulfur flushing of equipment with water after each day's use. Sulfur, in its solid form, is corrosive material. To reduce the effect, equipment should be flushed daily.

WEIGHT: 11.0 POUNDS PER GALLON

CONDITIONS OF SALE

1. Seller warrants that this product consists of the ingredients specified. No one, other than an officer of Seller, is authorized to make any warranty, guarantee or direction concerning this product.
2. Because the time, place, rate of application and conditions of use are beyond Seller's control, Seller's liability for damage to crops, animals, or property from storage and use of this product is limited to replacement or refund of purchase price.

EPA

SC-84

ACCEPTED

23 APR 1975

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 22555-3

RECOMMENDATIONS

Use 2 to 3 quarts in at least 10 good coverage. Begin applying the after planting. Maintain a schedule growing season.

are made on the basis of product recommendations are for sprays to ray equipment. For concentrated ers' recommendations.

Do not add a spreader. Apply at the appearance of Brown Rot. Repeat wet weather. Melanose - use 3 gal- ter petal fall. Scab - use 3 gallons. rees begin to flush and at 2/3 petal

as where copper injury is known to

1 1/2 gallons. Apply 1 to 3 weeks reeks if there has been a history of e or scattered bloom.

400 gallons per acre to give 1/2 lb. ply at 10% bloom and continue ut bloom. Repeat especially after ere State Disease Control officials fic varieties.

se 0.5 to 1.0 gallons. As a dormant e Disease apply Dec. 15 to Jan. 15. wn Rot should be applied through n stage. Do not apply after 50% eaf.

se 1 to 1.5 gallons. As a dormant Hole disease apply Nov. 15 to Jan. of Brown Rot should be applied popcorn stage. Do not spray when

lication when the disease normally y intervals unless stated otherwise.

allions.

LOWER: a 3 gallons.



TOP COP with sulfur

ACTIVE INGREDIENTS:

Sulfur as S	50.0%
Copper as Metallic	4.4%

INERT INGREDIENTS..... 45.6%

TOTAL.....100.0%

Derived from Elemental Sulfur and Tribasic Copper Sulfate, 8.4%


CAUTION KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid breathing spray mist. May cause irritation of nose, throat or skin. Avoid contact with eyes, skin and clothing. Do not contaminate feed and foodstuffs. Do not apply to any area not specified on label. Do not re-use empty container. Wash thoroughly with water and detergent. Puncture, make unusable. Dispose of in a safe place.

NET WEIGHT OF CONTENTS: 55 POUNDS

MANUFACTURED BY:

STOLLER CHEMICAL COMPANY, INC.
8705 KATY FREEWAY, SUITE 400
HOUSTON, TEXAS 77024



CARROTS:
Alternaria Leaf Blight, Cercospora Blight, Downy Mildew - use 3 gallons.

EGGPLANTS:
Alternaria Blight, Anthracnose, Phomopsis - use 3 gallons.

PEPPERS:
Anthracnose, Bacterial Spot, Cercospora Leaf Spot (Frogeye) - use 3 gallons.

POTATOES:
Early and Late Blight - use 4 gallons.

SPINACH:
Downy Mildew (Blue Mold), White Rust - use 3 gallons.

CAUTION

Some plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur on any crop unless local use has proven that sulfur is safe in that locality.

During periods of high temperature, sulfur may burn foliage and fruit. Do not make sulfur applications at excessively high temperatures. Do not use Top Cop with Sulfur with oil or within 4 weeks of an oil application.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS FOR MIXING

BEFORE USING, STIR UNTIL SMOOTH

Pour recommended amount into partially filled spray tank. Keep agitator running during filling and spraying operations. Do not allow mixture to stand. Failure to maintain agitation will cause Top Cop with Sulfur to settle and may necessitate manual stirring to redisperse.

The strong adhesive properties of Top Cop with Sulfur act as a sticker on the plant, and the sticking characteristic necessitates the flushing of equipment with water after each day's use. Sulfur, in any form, is corrosive material. To reduce the effect, equipment should be flushed daily.

WEIGHT: 11.0 POUNDS PER GALLON

CONDITIONS OF SALE

1. Seller warrants that this product consists of the ingredients specified. No one, other than an officer of Seller, is authorized to make any warranty, guarantee or direction concerning this product.
2. Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's Liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price.

EPA Reg No 22555-3

ACCEPTED

23 APR 1975

REGISTERED UNDER THE FEDERAL INSECTICIDE AND RODENTICIDE ACT ECONOMIC POISON REGISTER UNDER NO. 22555-3

Job# 5741

FOLIAR SPRAY RECOMMENDATIONS

PEANUTS

To control rust and leaf spot, use 2 to 3 quarts in at least 10 gallons of water to maintain good coverage. Begin applying the above mixture 40 to 60 days after planting. Maintain a schedule of 10 to 14 days through the growing season.

The following recommendations are made on the basis of product per 100 gallons of water. These recommendations are for sprays to be applied by conventional spray equipment. For concentrated sprayers, consult the manufacturers' recommendations.

CITRUS:

Brown Rot - use 1 gallon; do not add a spreader. Apply at the first indication of rain or first appearance of Brown Rot. Repeat as needed to protect during wet weather. Melanose - use 3 gallons. Apply 1 to 3 weeks after petal fall. Scab - use 3 gallons. Apply two sprays, use before trees begin to flush and at 2/3 petal fall.

Caution: Do not apply in areas where copper injury is known to occur.

Florida Only: Melanose - use 1 1/2 gallons. Apply 1 to 3 weeks after petal fall. Repeat in 4 weeks if there has been a history of the disease, a wet spring or late or scattered bloom.

PEARS:

Fire Blight: use 0.5 gallons in 400 gallons per acre to give 1/2 lb. metallic copper per acre. Apply at 10% bloom and continue every 5 to 7 days throughout bloom. Repeat especially after showers or rain. Use only where State Disease Control officials recommend such use on specific varieties.

ALMONDS:

Brown Rot and Shot Hole - use 0.5 to 1.0 gallons. As a dormant spray for control of Shot Hole Disease apply Dec. 15 to Jan. 15. Bud sprays for control of Brown Rot should be applied through swelling bud stage to popcorn stage. Do not apply after 50% bloom or when trees are in leaf.

APRICOTS:

Brown Rot and Shot Hole - use 1 to 1.5 gallons. As a dormant spray for the control of Shot Hole disease apply Nov. 15 to Jan. 15. Bud sprays for control of Brown Rot should be applied through swelling bud stage to popcorn stage. Do not spray when trees are in leaf.

For the following crops begin application when the disease normally appears and repeat at 7 to 10 day intervals unless stated otherwise.

SUGAR BEETS:

Cercospora Leaf Spot - use 4 gallons.

BEEETS:

Downy Mildew - use 3 gallons.

BROCCOLI, CABBAGE, CAULIFLOWER:

Downy Mildew, Leaf Spot - use 3 gallons.



**TOP COP
with sulfur**

ACTIVE INGREDIENTS:

Sulfur as S 50.0%
Copper as Metallic 4.4%

INERT INGREDIENTS 45.6%


TOTAL 100.0%

Derived from Elemental Sulfur and Tribasic Copper Sulfate, 8.4%

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid breathing spray mist. May cause irritation of nose, throat or skin. Avoid contact with eyes, skin and clothing. Do not contaminate feed and food stuff. Do not apply to any area not specified on label. Do not re-use empty container. Wash thoroughly with water and detergent. Puncture, make unusable. Dispose of in a safe place.



NET WEIGHT OF CONTENTS: 55 POUNDS

MANUFACTURED BY:

STOLLER CHEMICAL COMPANY, INC.
8705 KATY FREEWAY, SUITE 400
HOUSTON, TEXAS 77024

CARROTS:

Alternaria Leaf Blight, Cercospora Blight, Downy Mildew - use 3 gallons.

EGGPLANTS:

Alternaria Blight, Anthracnose, Phomopsis - use 3 gallons.

PEPPERS:

Anthraco-nose, Bacteria Spot, Cercospora Leaf Spot (Frog-eye) - use 3 gallons.

POTATOES:

Early and Late Blight - use 4 gallons.

SPINACH:

Downy Mildew (Blue Mold), White Rust - use 3 gallons.

CAUTION

Some plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur on any crop unless local use has proven that sulfur is safe in that locality.

During periods of high temperature, sulfur may burn foliage and fruit. Do not make sulfur applications at excessively high temperatures. Do not use Top Cop with Sulfur with oil or within 4 weeks of an oil application.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS FOR MIXING

BEFORE USING, STIR UNTIL SMOOTH

Pour recommended amount into partially filled spray tank. Keep agitator running during filling and spraying operations. Do not allow mixture to stand. Failure to maintain agitation will cause Top Cop with Sulfur to settle and may necessitate manual stirring to redisperse.

The strong adhesive properties of Top Cop with Sulfur act as a sticker on the plant, and the sticking characteristic necessitates the flushing of equipment with water after each day's use. Sulfur, in any form, is corrosive material. To reduce the effect, equipment should be flushed daily.

WEIGHT: 11.0 POUNDS PER GALLON

CONDITIONS OF SALE

1. Seller warrants that this product consists of the ingredients specified. No one, other than an officer of Seller, is authorized to make any warranty, guarantee or direction concerning this product.

2. Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's Liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price.

EPA Reg No 22555-3

SC 84

ACCEPTED
23 APR 1975
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER
ED UNDER NO. 22555-3

Job# 574