

Begin when plants are small. Repeat at 5 to 7 day intervals. Do not apply within four days of harvest of succulent beans.

CABBAGE: Alternaria Leaf Spot, Downy Mildew—Begin 7 to 10 days after planting. Repeat at 7 to 5 day intervals in plant bed and field. Do not apply within 7 days of harvest.

CORN (Sweet): Helminthosporium Blight—Begin when plants are 6 to 8 inches high. Repeat at 4 to 7 day intervals until harvest. Do not feed treated forage to livestock.

CUCUMBERS: Alternaria (Microsporium) Leaf Spot, Angular Leaf Spot, Anthracnose, Downy Mildew, Pythium Fruit Rot—Begin when diseases threaten or plants begin to run. Repeat at 7 to 10 day intervals under severe disease conditions. Do not use within 5 days of harvest.

EGGPLANT: Anthracnose, Early Blight (Alternaria), Phomopsis Blight or Fruit Rot—Begin when fruit starts to form. Repeat at 7 to 10 day intervals.

ENDIVE, LETTUCE: Downy Mildew—Begin when disease appears. Repeat at 7 to 10 day intervals. Do not apply within 10 days of harvest. Remove residues from head lettuce by stripping and trimming, and from leaf lettuce and endives by washing or other effective means.

PEPPERS: Anthracnose, Cercospora Leaf Spot (Fragaria Spot), Ripe Rot—Begin when disease threatens. Repeat at 7 to 10 day intervals.

SQUASH (Summer and Winter): Anthracnose, Downy Mildew—Begin when plant starts to run. Repeat at 7 to 10 day intervals. Do not use within 5 days of harvest.

TOMATOES (Greenhouse and Field): Cladosporium Leaf Mold, Early and Late Blight, Gray Leaf Spot (Stemphylium), Septoria Leaf Spot—Begin at first fruit cluster and repeat at 7 to 10 day intervals. Do not use within 5 days of harvest.

APPLY 50 TO 100 GALLONS OF DILUTED SPRAY PER ACRE (1.1—2.2 Gallons Per 1000 Sq. Ft.)

CAUTION: May cause irritation of eyes, nose, throat and skin. Avoid breathing spray mist. May be absorbed through skin. Avoid contact with eyes, skin or clothing. In case of contact, flush with plenty of water. For eyes get medical attention. Keep away from fire and sparks.

This product is toxic to fish. Keep out of any body of water. Apply this product only as specified on label. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not reuse container. Destroy when empty. Wrap in newspaper and place in trash can for disposal.

**PACKED BY B. G. PRATT DIVISION
GABRIEL CHEMICALS LTD., PATERSON, N.J. 07509**

PRATT

Liquid MANEB

PLANT FUNGICIDE

CONTROLS COMMON DISEASES ON TURF, ORNAMENTALS AND VEGETABLES

(*As listed on back panel)

CONTAINS 4 POUNDS MANEB PER GALLON WITH ZINCADED

CAUTION: KEEP OUT OF REACH OF CHILDREN

SHAKE WELL BEFORE USING

GENERAL DIRECTIONS
Mix with water according to directions. Apply thoroughly to cover all plant surfaces.

- FLOWER AND ORNAMENTAL USES**
GENERAL DILUTION: 1-400, 2 teaspoonsful per gallon, 1 pint per 50 gallons.
- CARNATIONS, DAHLIAS:** Alternaria Leaf Spot, Anthracnose, Botrytis Blight—Begin when new growth starts. Repeat weekly.
 - DOGWOOD:** Anthracnose—Begin when bud opens. Repeat when bracts fall four weeks later and in late summer.
 - GLADIOLUS:** Botrytis Blight, Curculiaris and Stemphylium Leaf Spots—Begin when flower spikes are developing. Repeat 2 to 3 times at weekly intervals.
 - LILLIES:** Botrytis Blight—Begin with new growth. Repeat weekly.
 - PARSIES:** Anthracnose—Begin with new growth. Repeat weekly.
 - PEONIES:** Alternaria Leaf Spot, Botrytis Blight, Phytophthora Blight—Apply to foliage in early spring and early fall.
 - ROSES:** Black Spot, Cercospora Leaf Spot (Terra), Rust (California)—Begin when first leaves unfold. Repeat at 7 to 10 day intervals.
 - SNAP DRAGONS:** Rust—Begin with emergence. Repeat weekly.
 - ZINNIAS:** Alternaria Leaf Spot, Leaf Blights—Begin with emergence. Repeat weekly.

- TURF AND GRASS USES**
- TURF GRASSES (Mown Areas):** Mix prescribed volume of fungicide with sufficient water to cover 1000 square feet. Use only 5 or 10 gallons of water are required depending upon spray equipment used. Keep pets and children off treated area until spray thoroughly dries.
 - BROWN PATCH:** 4.8 fluid ounces. Begin at first sign of disease. Repeat 7 to 14 day intervals.
 - DOLLAR SPOT:** 9.6 to 12.8 fluid ounces. Apply as for brown patch above.
 - MELTING-OUT:** 4.8 to 6.4 fluid ounces. Apply as for brown patch above.
 - LEAF, STEM, AND STRIPE RUSTS:** 2 fluid ounces. Begin when rust pustules are first seen. Repeat at 7 to 14 day intervals. Do not graze treated areas. Do not feed clippings to livestock.

VEGETABLE USES
GENERAL DILUTION: 2 teaspoonsful to 1 tablespoonful per gallon, or 1.2 to 1.6 quarts per 100 gallons of spray.

BEANS (Snap or String): Anthracnose, Downy Mildew, Rust—

WP10162.5-11-73

904-190