



PRECAUTIONARY STATEMENTS
Hazards To Humans and Domestic Animals
Caution

Harmful if swallowed. Avoid breathing spray mist. May cause irritation of nose, throat or skin. Avoid contact with eyes, skin and clothing.

Environmental Hazards

Do not apply directly to lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

USE PRECAUTIONS

Some fruits and other plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur on any crop unless local use has proved that sulfur is safe in that locality. When crops are intended for processing, consult with processor before applying sulfur.

During periods of high temperature sulfur may burn foliage and fruit. Do not make sulfur applications at excessively high temperatures. Do not use sulfur with oil or within 4 weeks of an oil application unless in a dormant application.

When sulfur is used with arsenicals, lime should be added to prevent plant injury.

DIRECTIONS FOR USE

GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DIRECTIONS FOR MIXING
BEFORE USING, STIR UNTIL SMOOTH

Pour recommended amount in partially filled spray tank. Keep agitator running during filling and spraying operation. DO NOT ALLOW MIXTURE TO STAND. Failure to maintain agitation will cause BIG 8 Flowable Sulfur to settle and may necessitate manual stirring to redispense.

The strong adhesive properties of BIG 8 Flowable Sulfur act as a sticker on the plant, and the sticking characteristic necessitates the flushing of equipment with water after each day's use. Sulfur in any form is corrosive material. TO REDUCE THE EFFECT, EQUIPMENT SHOULD BE FLUSHED DAILY.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Pesticide, spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of according to procedures approved by Federal, state and local disposal authorities.

Triple rinse (or equivalent) and dispose of in an approved land-fill or bury in a safe place.

Consult Federal, state or local disposal authorities for approved alternative procedures.

Tree Fruits

Amount per 100 gallons (Dilute)

Apple, Pear
Scab

Pre-bloom through: Calyx sprays 1 to 2-1/3 pints
Cover sprays 2/3 to 1-1/3 pints

Powdery Mildew

Pre-bloom through: Calyx sprays 1-1/2 to 2 pints
Cover sprays 1-1/3 pints

That Big 8

FLOWABLE SULFUR

CONTAINS 8 POUNDS SULFUR PER GALLON

ACTIVE INGREDIENT(S):

Sulfur 64%

INERT INGREDIENT(S): 36%

Total 100%

CAUTION:

Keep Out of Reach of Children

If Swallowed: Induce vomiting by sticking finger down throat. Never give anything by mouth to an unconscious person. Get medical attention. If On Skin: Wash thoroughly with soap and water. Get medical attention. If in Eyes: Wash eyes with plenty of clear water for at least 15 minutes. Get medical attention. See left side panel for additional precautionary statements.

THIS PRODUCT FREEZES AT 30° F.

A PRODUCT OF:

Stoller Chemical Company, Inc.

8582 Katy Freeway, Suite 200

Houston, Texas 77024

EPA EST. NO. 22555-CA-01

EPA REG. NO. 22555-6

NET CONTENTS: 5 Gallons



Tree Fruits

Amount per 100 gallons (Dilute)

Peach

Powdery Mildew, Brown Rot, Leafspot:

Pink and bloom sprays 1 to 2-1/2 pints

Petal fall, shuck and cover sprays 1/2 to 1 pint

Cherry, Plums & Prunes

Brown Rot, Leafspot:

Pink and bloom sprays 1 pint

Petal fall, shuck and cover sprays 1/2 to 1 pint

Citrus (grapefruit, lemons, limes,

Oranges, tangelos and tangerines)

Rust Mite: 1 to 3 pints

DO NOT apply within 4 weeks of an oil application.

Vines (grapes)

Powdery Mildew: Use 2/3 to 1-1/3 pints per 100 gallons water; maximum of 5 pints BIG 8 Flowable Sulfur per acre in spring and summer. Make first application when shoots are 6 to 8 inches long, second when shoots are 12 to 15 inches long, and third at about bloom stage.

The timing, rate and number of applications will vary with local conditions. Consult State Agricultural Service or State Agricultural Experiment Stations in your region for specific information. Follow the USE PRECAUTIONS given on this label.

Row Crops

Peanuts

Rust: Apply 1-1/3 to 2-1/2 pints per acre in sufficient water for thorough coverage. Begin application approximately 6 weeks after planting. Repeat at 7 to 10 day intervals as long as rust control is required.

Cotton

Atlantic Mite: Apply 1-1/3 pints for early season control and up to 5 pints for mid to late season control.

Potatoes

Powdery Mildew: Use 2 to 2-1/2 pints in 8 gallons water per acre by air. Apply at the first sign of infection and repeat at 3 week intervals.

Strawberries

Powdery Mildew: Use 2/3 pint. Apply at first sign of infection and repeat at 3 week intervals. For strawberries that will be canned, discontinue application of sulfur well before harvest, in accordance with local canners' recommendations.

Tomatoes

Tomato Russet Mite: Use 2-1/2 to 5 pints per acre depending upon size of plants. Use full coverage sprays.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use; but neither this warranty or any other warranty of merchantability of fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions not reasonably foreseeable to the seller, the buyer assumes the risk of any such use.

SC-722-C



12