

3 tablespoonsful	1 gallon water
1 pound to	5 gallons water
1 pound to	7 1/2 gallons water
3 pounds to	30 gallons water

APPLES: South Slope fruit and vegetable market has a large selection of apples for sale. The apples are mostly from the local orchards and are of excellent quality. The apples are available in a variety of sizes and quantities. The prices are very reasonable and the apples are always fresh. The market is open from 10 a.m. to 6 p.m. daily. The address is 1234 Main St., South Slope, Seattle, WA 98108. The phone number is 555-1234.

CHERRIES: In the early spring, when the cherries are in flower, spray with a solution of 100 parts of water to 1 part of lime sulfur. Repeat this treatment in May and June. In the fall, spray with a solution of 100 parts of water to 1 part of lime sulfur. Repeat this treatment in the fall.

PEACHES 4PR.COTS
PLUM PEAR

STRAWBERRIES: (*Botrytis* leaf spot, aphids, spider mite, strawberry leafroller) Apply Spray A thoroughly over spray when insects first appear. For disease control begin applications when new growth appears. Repeat applications at 7 to 10 day intervals as insects are warrant. Do not apply within 7 days of harvest.

GRAPES - Black rot, under mild, early-type grape varieties, the grape leaf-roller, Japanese beetle, Aphid, and the European vine weevil spread their eggs on the leaves. The eggs hatch and the young insects feed on the leaves. For black rot, 2 applications should be made before blossoming and 3 or 4 more applications at 10 to 15 day intervals. For grape leaf-roller Moths apply as a post-blossom spray and repeat 10 days later, make third and fourth applications at 10 day intervals if necessary. Apply with 10 days of harvest.

BLACKBERRY, RASPBERRY, DEWBERRY: To control Anthracnose use a pre-bloom spray (when new growth starts) to 1 in. in leaf and first young spray right after petals fall. **RASPBERRY:** Spray with Bordeaux first. Repeat applications at 10 to 14 day intervals until 1 in. in leaf and new growth starts. Make thorough coverage. For Anthracnose use same fungistat application as for Blackberry. Spray to point of drip. Do not spray during rain.

Apple Orange Peaches Apricots Pear Plum

1. The first step in the process is to identify the problem or issue that needs to be addressed. This involves gathering information and understanding the context of the problem.

Keep new growth covered in the spring by frequent applications. Such a program is the best insurance against disease and insect damage. To control Alternaria Leaf Spot, Rust and Carnation Bud Mite on CARNATIONS, Botrytis Flower Blight and Septoria Leaf Spot on CAULIFLOWERS, and Black Spot Aphids and Leafhoppers on ROSE, use 1 cup in one 1 1/2 gallon of water. Spray underneath and top of foliage thoroughly. Repeat every 7 to 10 days.

EVERGREENS: For Aphids, Mealybugs, Spider Mites, mix 1/2 cup to one gallon of water. Make thorough full-coverage applications and repeat as necessary. **NOTE:** Injury may occur on some species of cacti and conifers.

To prevent damping-off on **AZALEAS, CARNATIONS**
and **MUMS**, use one (1) cupful per one (1) gallon
of water as a root dip treatment before planting.

Hazards: If swallowed. Avoid breathing dust. Wear eye protection. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. May be absorbed through the skin. Avoid contamination of feed and foodstuffs. When not in use keep container tightly closed. Apply this product only as specified on the label.

DOI: 10.1002/eqm2.1201, Published online 10 June 2018 in Wiley Online Library (wileyonlinelibrary.com). © 2018 John Wiley & Sons, Inc.

This contact will kill fish and midlife keepers.

contaminate water by cleaning of equipment in the
vicinity of wastes.

The following table shows the results of the regression analysis for the dependent variable "Number of children in the household" (N = 1,000). The independent variables are "Age of the head of household" and "Gender of the head of household". The results are presented in the following table:

1044

3776-267
Multi-Purpose
HOME ORCHARD
3486-60
SPRAY

ACTIVE INGREDIENTS	
Captan (N-Trichloromethylthio-4-cyclohexene- 2-dicarboximide)	10.00%
Malathion (O,O-Dimethyl dithiophosphate of diethyl mercaptosuccinate)	3.00%
2-O-Diethyl 2-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate*	5.00%
INERT INGREDIENTS	80.00%
Total:	100.00%
*Diazinon	

• **Discharge**

KEEP OUT OF REACH OF CHILDREN
See back panel for additional caution statement.

NET WEIGHT - POUND

Manufactured by:

SHENANDOAH IOWA 5160.

~~SECRET~~