

super

\$11.98

NET WT. 13 LBS.
TREATS 2000 SQ. FT.

KELLY

ACCEPTED

WESTERN

Oct 7, 1974

UNDER THE FEDERAL PESTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 247-6

**dichondra
weed 'n feed**

**dichondra
weed 'n feed**

Kills Oxalis, Dutch Clover, Black-

ERN

dichondra weed 'n feed

dichon
weed 'n

**Kills Oxalis, Dutch Clover, Black-
Medic and Annual Bluegrass** (See rear panel for
additional weeds)

Prevents Crabgrass (Guaranteed)

Contains Long-lasting Dichondra Food 11-0-0

Supplies Iron, Sulfur, Manganese, Zinc

ACTIVE INGREDIENTS:

S-(O,O-diisopropyl phosphorodithioate) ester of
N-(2-mercaptoethyl) benzene sulfonamide 4.42%
Monuron (3-[p-chlorophenyl]-1,1-dimethylurea) 0.35%

INERT INGREDIENTS 95.23%
100.00%

EPA Reg. No. 24232-6
EPA Est. No. 24232-UT-01

GUARANTEED FERTILIZER ANALYSIS:

Total Nitrogen (N) 11.00%
3.30% Water Insoluble Nitrogen
11.00% Urea Nitrogen (5.5% from Ureaforn)
Sulfur as S* 1.20%
Iron as Fe* 1.00%
Manganese as Mn* 0.10%
Zinc as Zn* 0.10%

INERT WEIGHT 86.60%
*Derived from specially processed natural
minerals containing a mixture of sulfides
which gradually become soluble.

CAUTION:

KEEP OUT OF REACH OF CHILDREN

Avoid contact with skin, eyes, or clothing. Harmful
if swallowed. Avoid contamination of feed and feed-
stuffs. Wash thoroughly after use. Do not breathe dust.
Keep children and pets off treated areas until material
is watered in lawn and allowed to dry.
This product is toxic to fish.
Keep out of lakes, streams or ponds. Do not apply
where runoff is likely to occur. Do not contaminate water
by cleaning of equipment, or disposal of wastes. Apply
this product only as specified on this label.
Do not store near fertilizers, seeds, insecticides, or
fungicides. Do not reuse container. Destroy when empty.


KELLY WESTERN SEED DIVISION of THE PAX COMPANY
580 West 13th South, Salt Lake City, Utah 84115

super



dichondra weed 'n feed

super



super

KELLY **KW** WESTERN

weed 'n feed

GENERAL WEED CONTROL

Winter weeds can be controlled by a fall application. Spring and summer weeds can be controlled by applications made in March, April or May. Treatments may be repeated the year round at regular four month intervals to control weeds and prevent germination of many weed seeds. Do not repeat treatments more than three times a year. On heavy or highly organic soils reduce coverage to 1500 sq. ft. for best results. **APPLY ONLY TO MATURE DICHONDRA LAWNS** (at least one year old)

SPOT TREATING

1. Apply to weedy area at recommended rate. Avoid over-application in spot treatments. Use spreader if possible. If discoloration of weeds has not started in three weeks, repeat at recommended rates. Final result at effective rate may take up to four weeks.
2. Immediately wash product off the dichondra leaves as indicated in step 3 below.

CRABGRASS

For best control of CRABGRASS, apply this product at the recommended rate (see "How to Apply"). Between fall and late spring (May 31). Applications made after crabgrass sprouts will kill seedling crabgrass plants plus prevent additional infestation.

OXALIS — CLOVER

Make application according to directions below at any time during the year. Effects of application are usually slow to appear and may not become evident before three or four weeks. A heavy infestation may require a second application. (See spot treating.)

BLACK MEDIC & CHICKWEED


Medic — make normal application. Results should be evident in 7 days. Chickweed — double application necessary in severe infestations (see spot treating).

ANNUAL BLUEGRASS

Applications made at any time of the year will kill annual blue grass, winter grass at any stage of growth plus prevent further infestation. (See spot treating.)

dichondra
weed 'n feed

HOW TO APPLY FOR A BETTER DICHONDRA LAWN



1 Measure area to be covered (for example 40' x 50'). One 13 lb. bag covers 2000 sq. ft. Dichondra should be dry at time of application. Remove grass clippings, leaves and other objects so that product may evenly contact soil surface.

2 Set fertilizer spreader as suggested below. For best control follow crisscross pattern as illustrated. Recommended spreader settings given below are for crisscross patterns. **CAUTION** Never make sharp turns with spreader open. Always close spreader while changing directions.


3 Immediately after spreading, wash product off the dichondra with the medium spray from a hand-held nozzle. Sprinkling systems and stationary sprinklers are less satisfactory. After application, do not water dichondra for 24 hours, then resume normal watering practice. Do not allow water to puddle or flood.

TERN

dichondra weed 'n' feed

WESTERN
KELLY PAX
SUI

HOW TO APPLY FOR A BETTER DICHONDRA LAWN



1 Measure area to be covered (for example 40' x 50'). One 13 lb. bag covers 2000 sq. ft. Dichondra should be dry at time of application. Remove grass clippings, leaves and other objects so that product may evenly contact soil surface.

2 Set fertilizer spreader as suggested below. For best control follow crisscross pattern as illustrated. Recommended spreader settings given below are for crisscross patterns. CAUTION: Never make sharp turns with spreader open. Always close spreader while changing directions.

3 Immediately after spreading, wash product off the dichondra with the medium spray from a hand-held nozzle. Sprinkling systems and stationary sprinklers are less satisfactory. After application, do not water dichondra for 24 hours, then resume normal watering practice. Do not allow water to puddle or flood.

COMMON WEEDS CONTROLLED KILLS PREVENTS

Oxalis	crabgrass	redroot pigweed
crabgrass (round)	watergrass	goosegrass
black medic	annual bluegrass	shepherd's purse
chickweed	deadnettle	and many other
annual bluegrass	grass buttons	spring germinat-
annual ryegrass	lambsquarters	ing annuals
Dutch clover		

SUGGESTED SPREADER SETTINGS

Spreading rate may vary with the age, model, and condition of the spreader. Check your spreader over an area of known size. (One 13 lb. bag covers 2000 sq. ft.) Recommended settings below are for crisscross applications. If your spreader is not included below, set spreader opening at lowest setting which gives an even flow, usually about one-third open. Spread in one direction and then crosswise. Continue crisscrossing until proper amount is applied.

Bandini		Lawn Beauty		Scott		Modern		Central		Deet		Red Star		Germans	
MODEL	SETTING	MODEL	SETTING	MODEL	SETTING	MODEL	SETTING	MODEL	SETTING	MODEL	SETTING	MODEL	SETTING	MODEL	SETTING
20 In.	8½	18-57	8½	35-8	9½	18 In.	7	20 In.	8½	20 In.	8½	20 In.	8½	20 In.	8½

WATERING

Do not water for 24 hours after the material has been washed off dichondra foliage. Then follow normal watering practice. White granules may be seen on the soil surface. These granules are the carriers and are not harmful to dichondra.

SHADED AREAS:

Not recommended for densely shaded areas.

RESEEDING

Dichondra lawns cannot be reseeded for 120 days after application. Materials that provide effective control of weeds may affect seedlings which germinate after treatment. Proper fertilization and watering will encourage the existing dichondra to fill in the thin or bare spots formerly occupied by weeds.

TREES & SHRUBS:

Do not apply on or near young trees or young shrubs.

BLADEGRASS LAWNS:

Do not use on blade grass lawns.

SUMMER APPLICATIONS:

1. Water dichondra lawn thoroughly the day before PAX application.
2. Do not apply on days when temperature is 85 degrees or above.

GUARANTEE: Manufacturer guarantees this product to give results as outlined on this bag or the purchaser may get his money refunded by the manufacturer. No other express warranties of any kind are made, except to conform to the promises and affirmations of facts made on the label. The warranty of merchantability, all other implied warranties and liability of the manufacturer without fault are expressly disclaimed. Buyer assumes all risk of personal injury or damage to property, except for the great mislabeling of the product. All claims must be presented to manufacturer in writing, within thirty days from the date a breach is discovered and an action thereon must be commenced within one year after the cause of action has accrued.

KELLY

WESTERN

dichondra weed 'n feed

dichondra
weed 'n feed

Kills Galva, Dutch Clover, Black-
Medic and Annual Bluegrass (Guaranteed)

Prevents Crabgrass (Guaranteed)

Contains Long-Lasting Dichondra Food 11-0-0

Contains Iron, Sulfur, Manganese, Zinc

super



WESTERN DIVISION of THE PAX COMPANY
Salt Lake City, Utah 84115

dichondra weed 'n feed