USE DIRECTIONS FOR CLEANING

Lightly soiled areas - surfaces to be cleaned should be mopped or brushed with a solution containing I ounce of Steri-Kleen Disinfectant per gallon of water. Rinse cleaned area with fresh water.

Heavily soiled areas - mop or brush soiled surface with a solution containing 2-3 ounces of Steri-Kleen Disinfectant per gallon of water. Rinse cleaned surface with fresh water.

GENERAL PURPOSE DISINFECTION

To disinfect walls, floors and other hard surfaces in homes, schools, restaurants, rest rooms and other public areas, mop or brush surface to be treated with a solution containing I ounce of Steri-Kleen Disinfectant per gallon of water. Surfaces that are heavily soiled should be cleaned with I-2 ounces of Steri-Kleen Disinfectant per gallon of water and then disinfected. Those surfaces which may later come in contact with food must be rinsed with potable water prior to reuse.

To disinfect locker rooms, shower stalls, walkways around pools and other areas against pathogenic fungi, mop area with a solution containing 3 4 ounce in 1 gallon water. This solution will destroy **Trichophyton interdigitale**, a causative agent of athlete's foot.

HOSPITAL DISINFECTION

To disinfect walls, floors, furniture and other hard surfaces in hospitals and nursing homes, mop or brush surface to be treated with a solution containing 2 ounces of Steri-Kleen Disinfectant per gallon of water. Those surfaces that are heavily soiled with blood, fecal material or exudates should be thoroughly scrubbed with a solution containing 2 ounces per gallon of water and then rinsed with a fresh solution containing 2 ounces per gallon of water before disinfection.

SANITIZATION

To sanitize walls, floors and equipment in food processing plants, first clean thoroughly with a solution containing 3 4 ounce of Steri-Kleen Disinfectant per gallon of water. If surface is heavily soiled, use Steri-Kleen Disinfectant at a rate of 2 or 3 ounces per gallon to insure thorough cleaning. Sanitize cleaned surface with a rinse solution containing 1 2 ounce of Steri-Kleen Disinfectant per gallon of water. Food contact surfaces must be rinsed with potable water prior to reuse.



DISINFECTANT - CLEANER - SANITIZER > FUNGICIDE - DEODORANT

ACTIVE INGREDIENTS

n-Alkyl (50% C14, 40% C12, 10% C16) dimethyl benzyl	• •
ammonium Chloride	5.0 %
Sodium metasilicate	3.0 %
Tetrasodium salt of ethylene diamine tetraacetic acid	1.8%
INERT INGREDIENTS	90.2 %

Phenol Coefficient: S. aureus - 32

S. typhosa - 42


EPA Reg. No. 24909-1-AA

ATEL AAM

DANGER: KEEP OUT OF REACH OF CHILDREN!

See first aid and other precautions on side panel.

NET CONTENTS: ONE U.S. GALLON



MANUFACTURED BY

KNAPP CHEMICAL COMPANY

5227 East-Pine Avenue • Fresno California 93727

DEODORIZATION

To deodorize surfaces during containing 2 ounces of Stem Steri-Kleen Disinfectant is a in household, industrial an Steri-Kleen Disinfectant we counter tops and furniture areas, garbage pails, food sobnoxious odors may devel antibacterial-antifungal denursing homes, public rooms efficient cleaning and anti-

Corrosive. Causes eye dam or on clothing. Protect ey swallowed. Avoid contami Container Disposal: When destroy it by crushing or

FIRST AID

In case of contact, immed at least 15 minutes. For nated clothing before reu If swallowed, drink promp solution or if these are not alcohol. Call a physician

NOTE TO PHYSICIAN

Probable mucosal damage Measures against circulate may be ne**gded.**



UNDER TO FUNGICIE FUN ECO LD UNDE be mopped or brushed infectant per gallon of

h a solution containing water. Rinse cleaned

homes, schools, restauh surface to be treated Disinfectant per gallon leaned with 1-2 ounces then disinfected. Those ust be rinsed with pot-

round pools and other n containing 3 4 ounce phyton interdigitale, a

rfaces in hospitals and the asolution containing er. Those surfaces that as should be thoroughly lon of water and then gallon of water before

sing plants, first clean teri-Kleen Disinfectant ri-Kleen Disinfectant at ugh cleaning. Sanitize ounce of Steri-Kleen as must be rinsed with



DISINFECTANT - CLEANER - SANITIZER FUNGICIDE - DEODORANT

ACTIVE INGREDIENTS	.,,,	TEI
n-Alkyl (50% Ci4, 40% Ci2, 10% Ci6) dimethyl benzyl		**
ammonium Chloride		5.0 %
Sodium metasilicate		3.0 %

Phenol Coefficient: S. Lureus - 32 S. typhosa - 42

EPA Reg. No. 24909-1-AA

DANGER: KEEP OUT OF REACH OF CHILDREN!

See first aid and other precautions on side panel.

NET CONTENTS: ONE U.S. GALLON



MANUFACTURED BY

KNAPP CHEMICAL COMPANY

5227 East Pine Avenue • Frisno California 93727

DEODORIZATION

"ivicion

ATOT AAM &

To deodorize surfaces during cleaning, mop or brush surfaces with a solution containing 2 ounces of Steri-Kleen Disinfectant per gallon of water.

Steri-Kleen Disinfectant is an effective anti-hacterial cleaner designed for use in household, industrial and institutional cleaning. When used as directed, Steri-Kleen Disinfectant will disinfect hard surfaces, such as floors, walls, counter tops and furniture. It may be relied upon to deodorize bathroom areas, garbage pails, food storage areas, kennel areas and other areas where obnoxious odors may develop. This product may be used as a general purpose antibacterial-antifungal detergent in homes, schools, restaurants, hospitals, nursing homes, public rooms, food processing plants and in other places where efficient cleaning and antimicrobial action is desired.

DANGER:

Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling. Harmful or fatal if swallowed. Avoid contamination of food.

Container Disposal: When empty, rinse container thoroughly with water and destroy it by crushing or burying in a safe place.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be negded.

ACCEPTED

置至高四 374

UNDER THE FEDERAL DESECTICIDE
FUNCIONE AND RODENTICIDE ACT
FUN ECONOMIC POROM REGISAL...
LD UNDER NO. 224 909