

DANGER:
KEEP OUT OF REACH OF CHILDREN

Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

Rinse container thoroughly with water and discard it.

Manufactured by
R.P. ECKERT CO
 1898 Kingsway Court
 Cincinnati, Ohio 45230
 Phone: 513 231-7740



EX-O-DINE

★ IODINE ★

DETERGENT SANITIZE

DISINFECTANT

RECOMMENDED FOR:

CLEANING, DISINFECTING, AND SANITIZING IN RESTAURANTS, FOOD SERVICE AND FOOD PROCESSING PLANTS, HOSPITALS AND COMMERCIAL BUILDINGS

ACTIVE INGREDIENTS:

Alpha-(p-Nonylphenyl)-omega-hydroxypoly (oxyethylene) iodine complex
 (Providing 1.75% titratable iodine)18.00

Phosphoric Acid16.00

INERT INGREDIENTS:

Manufactured under the following U.S.

Patents: (3,028,299) (2,977,315)
 (2,931,777)

TOTAL:

65.95
 100.00

DANGER:

Keep Out of Reach of Children

(SEE SIDE PANEL FOR FIRST AID AND ADDITIONAL PRECAUTIONS)

EPA REGISTRATION NO. 25293 1

Net Contents One U.S. Gallon

EX-O-DINE

IODINE ★

URGENT SANITIZER

DISINFECTANT

RECOMMENDED FOR:

DISINFECTING, AND SANITIZING IN RESTAURANTS, FOOD SERVICE PROCESSING PLANTS, HOSPITALS AND COMMERCIAL BUILDINGS

CONTENTS:

..... (2,977,315) (777) (yl)-omega-hydroxypoly (oxyethylene)-iodine complex (titratable iodine)	18.05%
.....	16.00%
NETS:	<u>65.95%</u>
following U.S. (2,977,315) (777)	TOTAL: 100.00%

DANGER:

Out of Reach of Children

(SEE SIDE PANEL FOR FIRST AID AND ADDITIONAL PRECAUTIONS)

NO. 25293 1

Net Contents One U.S. Gallon

DIRECTIONS FOR USE

For general cleaning and disinfecting in one operation, use 3 ounces of EX-O-DINE to 5 gallons of water.

For porous or extremely dirty surfaces, use 6 ounces to 5 gallons of water.

For sanitizing previously-cleaned surfaces, use 1 ounce of EX-O-DINE in 5 gallons of water.

RESTAURANTS, FOOD SERVICE & FOOD PROCESSING PLANTS:

Brush-type low temperature glass washer - Adjust machine in accordance with machine manufacturer's directions, if necessary. A solution of 1/2 ounce of EX-O-DINE to 2 1/2 gallons of water provides 25 ppm titratable iodine.

Third Sink sanitizing - For those operations in which no mechanical washing is possible, or for pots and pans. Wash thoroughly with a suitable detergent and rinse completely before immersing in the sanitizing solution. This is critical, since soil or detergent reduces the effectiveness of iodophors. A solution of 1/2 ounce of EX-O-DINE to 2 1/2 gallons of water provides 25 ppm titratable iodine.

To sanitize previously-cleaned food processing equipment and utensils, as above, use 1/2 ounce of EX-O-DINE to 2 1/2 gallons of water. Allow to drain but do not rinse. When used in this manner, EX-O-DINE does not require a potable water rinse, in accordance with the Federal Food Additive Regulation 121.2547.

HOSPITALS:

Use 3 ounces EX-O-DINE to 5 gallons of water for disinfecting in animal rooms and cages, cafeterias, morgues, laboratories, operating areas, patient rooms, bedframes, walls and floors, thermometers, instruments, rubber gloves, tubing & catheters. Allow at least 10 minute contact time. Rinse all food handling surfaces with potable water before reuse. Before disinfecting, instruments should be cleaned to remove all adhering blood and serous exudates.

GENERAL HOUSEKEEPING

Use 3 ounces of EX-O-DINE to 5 gallons of water for cleaning and disinfecting floors, walls, lavatories, shower rooms, refuse containers, kitchen areas and lunch rooms. Rinse all food handling equipment with potable water before reuse.