

PM 31

5/12/75

25891-5

PM 181

When Lispar Algacide is used as directed, it will kill and prevent development of green algae (Cocconeis pyrenoidosa (green), Phormidium inundatum (black algae), Phormidium (Square D algae), Lyngbya varicolor (blue-green) and other swimming pool

Lispar Algacide is compatible with most commonly used swimming pool chemicals, but should not be mixed with soap or other cleaners, or premixed with other water-treating chemicals. Lispar Algacide works best when added by itself directly to pool water. Lispar Algacide at use dilution is non-corrosive to inanimate surfaces.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, drink milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Call a physician.

Rinse empty container thoroughly with water and discard.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

CHILD and/or WILDLIFE PRECAUTIONS

This product is toxic to fish. Treated effluent should not be discharged where it will drain into lakes, streams, ponds or public waters. Apply this product only as specified on this label.

Mfg. by
LISPAR, LTD.

3236 N. 11th St.
Phila, Pa. 19140

ACCEPTED
MAY 12 1975
FEDERAL BUREAU OF INVESTIGATION
FOR ECONOMIC POISON REGISTRATION
UNDER NO. 25891-5

LISPAR

ALGAECIDE 10%

Concentrated Swimming Pool Algacide
Keeps Pool Water Clear and Free of Visible Algae
Controls Algae Slimes

ACTIVE INGREDIENTS:

n-Alkyl (80% C ₁₄ , 30% C ₁₆ , 5% C ₁₂ , 5% C ₁₈)	
dimethyl benzyl ammonium chloride	8.4%
n-di-alkyl (80% C ₁₄ , 30% C ₁₆ , 5% C ₁₂ , 5% C ₁₈)	
monethyl benzyl ammonium chloride	1.6%
INERT INGREDIENTS	90%

DANGER

KEEP OUT OF REACH OF CHILDREN

CORROSIVE CAUSES SEVERE EYE AND SKIN DAMAGE. DO NOT GET IN EYES, ON SKIN OR ON CLOTHING. WEAR GOGGLES OR FACE SHIELD AND RUBBER GLOVES WHEN HANDLING. HARMFUL OR FATAL IF SWALLOWED. AVOID CONTAMINATION OF FOOD.

SEE OTHER PRECAUTIONS AND FIRST AID ON SIDE PANEL

DIRECTIONS FOR USE

Initial Dose: 1 gallon Lispar Algacide to 50,000 gallons of water.
Maintenance Dose: 1 quart Lispar Algacide to 50,000 gallons of water.

Initial dose is used when filling pool at beginning of swimming season. If pool has visible algae growth, treat with initial dose. Vacuum pool after 24 hours to remove algae debris. If visible algae are still present or re-appear, repeat this treatment procedure as necessary. When pool is free of visible algae, use the recommended maintenance dose.

Maintenance dose should be added at 3-5 day intervals. If high temperatures prevail or pool has unusually heavy use, add maintenance dose more frequently.

If make-up water is added at the rate of more than 10% per week of pool capacity, add 1 quart of Lispar Algacide for each 10,000 gallons of fresh make-up water added per week.

When swimming season is over, add 1 gallon of Lispar Algacide to 50,000 gallons of water in treating water left in pool. This dose helps provide a measure of control of algae growth during the winter months. This treatment will also help the servicing of the pool before it is returned to use the following season.

NET ONE GALLON (3.785 LITERS)

EPA EST. 25891-PA-1

EPA REG. 25891-16