U.S. ENVIRONMENTAL PROTECTION AGENCY OFFICE, OF PESTICIDES PROGRAMS REGISTRATION DIVISION (WH 567) WASHINGTON, IAC, 2040)	26884- 10	DEC 14 1984	
NOTICE OF PESTICIDE: THE GISTRATION THE REGISTRATION	NAME OF PESTICIDE PRODUCT		
(Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)	Algenerican	. ••	
NAME AND ADDRESS OF REGISTRANT (Include ZIP code)		<u> </u>	
Г	٦		
Rorco Mfg. Corp. P.O. Pox 521 Gefferson, GA 30549	٦		
NOTE: Changes in labeling formula differing in substance f submitted to and accepted by the Registration Division prio product always refer to the above U.S. EPA registration num	r to use of the label in commerce	=	
On the basis of information furnished by the registrant, the the Federal Insecticide, Fungicide, and Rodenticide Act.	above named pesticide is hereby	Registered Reregistered under	
A copy of the labeling accepted in connection with this Re	eistration/Reregistration is retur	ned herewith.	
Registration is in no way to be construed as an indorsement			
icide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as gwing the registrant a right to exclusive use of the name or to its use if it has been covered by others his product is conditionally registered in accordance with FIFPA sec.  3(c)(7)(A) provided that you:  1. Submit and/or cire all data required for registration/reregistration of your product under FIFPA sec. 3(c)(5) when the Acompany requires all registrants of a milar products to submit such data.  2. Add the phrase "FPPA Registration No. 200944-10" to your label before you release the product for shipment.  3. Submit five (5) copies of your final printo abbeling before you release the product for shipment. Pefor to the X-70 Ecclosure to a further description of final printed labeling.  If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFA sec. 6(e). Your remains for shipment of the product constitutes acceptance of these conditions.  A stamped copy of the label is enclosed for your remains.  Findictures  Findictures			
ATTACHMENT IS APPLICABLE			
IGNATURE OF APPROVING OFFICIAL		DATE	
THE STATE OF ALL HOTHING OF FIGURE		J DATE	

Algaecide No. 230

ATTETED
who COMMENSA
in EPA Letter Dated:

DEC 20 1984

Funglife and Redendede Act as and red, for the posticide registered under EFA Peg. Na.

## 

#### **KEEP OUT OF REACH OF CHILDREN**

# CAUTION

# PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid breathing vapors. Avoid contact with skin, eyes, or clothing.

FIRST AID: If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water Avoid alcohol. Call a physician immediately:

ENVIRONMENTAL HAZARDS: This pesticide is trivic to fish. Keep out of takes, streams, or por is. Permits may be required for discharges containing this pesticide into lakes, streams, ponds, or public water. For guidance, contact the regional office of the Environmental Projection Agency.

## DIRECTIONS FOR USE GENERAL CLASSIFICATION

Algaecide No. 230  and fungi in recirculating commercial and industrial witto its use, systems must be cleaned to remove algal slime, and other deposits. An initial slug addition of fluid ounces of Algaecide No. 230  of water to provide a concentration of 16 to 7  Algaecide No. 230 hased on in the system, is recommended. Repeat initial do age Subsequent slug additions of 0.47 to 4. Algaecide No. 230 per 1000 gallons 40 parts per million of Algaecide No. employed every 2 to 5 days, or as needed. The freque upon the relative amount of bisedoff and the severity of lem. Slug additions should be made in the sump of we Algaecide No. 230 is used to consir washing systems that maintain effective mist eliminate its use, systems should be cleaned to remove be deposits. An initial slug dose of 7.1 to 11 Algaecide No. 230 per 1000 generous of 4.8 to 11.9 fluid ounces of Algaer 1000 gallons of water should be employed each 1 to frequency of addition depends upon the relative amount of the bacterial problem. Slug additions may be made to collection trays of the airwash system.	It is a violation of Federal law to use this product in a
and fungi in recirculating commercial and industrial way to its use, systems must be cleaned to remove algal slime, and other deposits. An initial slug addition of fluid ounces of Algaecide No. 230 of water to provide a concentration of 16 to 7 Algaecide No. 230 based on in the system, is recommended. Repeat initial do age Subsequent slug additions of 0.47 to 4. Algaecide No. 230 per 1000 gallons 40 parts per million of Algaecide No. employed every 2 to 5 days, or as needed. The freque upon the relative amount of bleedoff and the severity of lem. Slug additions should be made in the sump of we Algaecide No. 230 is used to consir washing systems that maintain effective mist eliminate its use, systems should be cleaned to remove be deposits. An initial slug dose of 7.1 to 11 Algaecide No. 230 per 1000 generated. Repeat initial dosege until control is evident tions of 4.8 to 11.9 fluid ounces of Algaecide no 11.9 fluid ounces of Algaecide no frequency of addition depends upon the relative amount of the bacterial problem. Slug additions may be made to	its labeling.
to its use, systems must be cleaned to remove algal slime, and other deposits. An initial slug addition of fluid ounces of Algaecide No. 230 of water to provide a concentration of 16 to 7 Algaecide No. 230 passed on in the system, is recommended. Repeat initial do age Subsequent slug additions of 0.47 to 4. Algaecide No. 230 per 1000 gallons 40 parts per million of Algaecide No. employed every 2 to 5 days, or as needed. The freque upon the relative amount of bleedoff and the severity of lem. Slug additions should be made in the sump of we Algaecide No. 230 is used to consir washing systems that maintain effective mist eliminate its use, systems should be cleaned to remove be deposits. An initial slug dose of 7.1 to 11 Algaecide No. 230 per 1000 generated. Repeat initial dosege until control is evident tions of 4.8 to 11.9 fluid ounces of Algaer 1000 gallons of water should be employed each 1 to frequency of addition depends upon the relative amount of the bacterial problem. Slug additions may be made to	Algaecide No. 230 is used to
slime, and other deposits. An initial slug addition of fluid ounces of Algaecide No. 230 of water to provide a concentration of 16 to 7 Algaecide No. 230 , based on in the system, is recommended. Repeat initial do age Subsequent slug additions of 0.47 to 4. Algaecide No. 230 per 1000 gallons 40 parts per million of Algaecide No. employed every 2 to 5 days, or as needed. The freque upon the relative amount of bleedoff and the severity of lem. Slug additions should be made in the sump of we Algaecide No. 230 is used to consir washing systems that maintain effective mist eliminate its use, systems should be cleaned to remove be deposits. An initial slug dose of 7.1 to 11 Algaecide No. 230 per 1000 generated. Repeat initial dosege until control is evident tions of 4.8 to 11.9 fluid ounces of Algaecide to frequency of addition depends upon the relative amount of the bacterial problem. Slug additions may be made to	
fluid ounces of Algaecide No. 230 of water to provide a concentration of 16 to 7 Algaecide No. 230 , based on in the system, is recommended. Repeat initial do age Subsequent slug additions (* 0.47	
Algaecide No. 230	slime, and other deposits. An initial slug addition of
Algaecide No. 230	fluid ounces of Algaecide No. 230
in the system, is recommended. Repeat initial do lage Subsequent slug additions (* 0.47 to 4. Algaecide No. 230 per 1000 gailons 40 parts per million of Algaecide No. employed every 2 to 5 days, or as needed. The freque upon the relative amount of blaedoff and the saverity of lam. Slug additions should be made in the sump of we Algaecide No. 230 is used to consir washing systems that maintain effective mist eliminate its use, systems should be cleaned to remove be deposits. An initial slug dose of 7.1 to 11 Algaecide No. 230 per 1000 generated. Repeat initial dosage until control is evident tions of 4.8 to 11.9 fluid ounces of Algaecide no frequency of addition depends upon the relative amount of the bacterial problem. Slug additions may be made to	of water to provide a concentration of to
Subsequent slug additions (* 0.47 to 4. Algaecide No. 230 per 1000 gallons 40 parts per million of Algaecide No. employed every 2 to 5 days, or as needed. The freque upon the relative amount of bleedoff and the severity of lam. Slug additions should be made in the sump of we Algaecide No. 230 is used to consir washing systems that maintain effective mist eliminate its use, systems should be cleaned to remove be deposits. An initial slug dose of 7.1 to 11 Algaecide No. 230 per 1000 generated. Repeat initial dosage until control is evident tions of 4.8 to 11.9 fluid ounces of Algaecide No. 230 per 1000 gallons of water should be employed each 1 to frequency of addition depends upon the relative amount of the bacterial problem. Slug additions may be made to	Algaecide No. 230, based on
Algaecide No. 230 per 1000 gallons 40 parts per million of Algaecide No. employed every 2 to 5 days, or as needed. The freque upon the relative amount of bleedoff and the severity of lem. Slug additions should be made in the sump of we Algaecide No. 230 is used to consir washing systems that maintain effective mist eliminate its use, systems should be cleaned to remove be deposits. An initial slug dose of 7.1 to 11 Algaecide No. 230 per 1000 ger mended. Repeat initial dosege until control is evidentions of 4.8 to 11.9 fluid ounces of Algae per 1000 gallons of water should be employed each 1 to frequency of addition depends upon the relative amount of the bacterial problem. Slug additions may be made to	in the system, is recommended. Repeat initial do age
employed every 2 to 5 days, or as needed. The freque upon the relative amount of bleedoff and the severity of lem. Slug additions should be made in the sump of we Algaecide No. 230 is used to consider washing systems that maintain effective mist eliminate its use, systems should be cleaned to remove be deposits. An initial slug dose of 7.1 to 11 Algaecide No. 230 per 1000 gerended. Repeat initial dosege until control is evidentions of 4.8 to 11.9 fluid ounces of Algaer per 1000 galions of water should be employed each 1 to frequency of addition depends upon the relative amount of the bacterial problem. Slug additions may be made to	Subsequent slug additions (* 0.47 to 4
employed every 2 to 5 days, or as needed. The freque upon the relative amount of bleedoff and the severity of lem. Slug additions should be made in the sump of we Algaecide No. 230 is used to consider washing systems that maintain effective mist eliminate its use, systems should be cleaned to remove be deposits. An initial slug dose of 7.1 to 11 Algaecide No. 230 per 1000 gerended. Repeat initial dosege until control is evidentions of 4.8 to 11.9 fluid ounces of Algaer per 1000 galions of water should be employed each 1 to frequency of addition depends upon the relative amount of the bacterial problem. Slug additions may be made to	Algaecide No. 230 per 1000 gailon
upon the relative amount of bleedoff and the severity of lem. Slug additions should be made in the sump of we Algaecide No. 230 is used to consider washing systems that maintain effective mist eliminate its use, systems should be cleaned to remove be deposits. An initial slug dose of 7.1 to 11 Algaecide No. 230 per 1000 generated. Repeat initial dosege until control is evidentions of 4.8 to 11.9 fluid ounces of Algae per 1000 gallons of water should be employed each 1 to frequency of addition depends upon the relative amount of the bacterial problem. Slug additions may be made to	40 parts per million of Algaecide No.
lem. Slug additions should be made in the sump of we Algaecide No. 230 is used to consider washing systems that maintain effective mist eliminate its use, systems should be cleaned to remove be deposits. An initial slug dose of 7.1 to 11 Algaecide No. 230 per 1000 generated. Repeat initial dosege until control is evidentions of 4.8 to 11.9 fluid ounces of Algae per 1000 gallons of water should be employed each 1 to frequency of addition depends upon the relative amount of the bacterial problem. Slug additions may be made to	
Algaectide No. 230 is used to consir washing systems that maintain effective mist elimin to its use, systems should be cleaned to remove be deposits. An initial stug dose of 7.1 to 11 Algaectide No. 230 per 1000 gerended. Repeat initial dosage until control is evidentions of 4.8 to 11.9 fluid ounces of Algae per 1000 gallons of water should be employed each 1 to frequency of addition depends upon the relative amount of the bacterial problem. Slug additions may be made to	
to its use, systems that maintain effective mist eliminate its use, systems should be cleaned to remove be deposits. An initial slug dose of7.1	lem. Slug additions should be made in the sump of w
to its use, systems that maintain effective mist eliminate its use, systems should be cleaned to remove be deposits. An initial slug dose of7.1	Algaecade No. 230 is used to co
deposits. An initial strip dose of7.1	air washing systems that maintain effective mist elimi
Algaecide No. 230 per 1000 g mended. Repeat initial dosage until control is evidentions of <u>A. 8</u> to <u>11.9</u> fluid ounces of <u>Algae</u> per 1000 gallons of water should be employed each 1 to frequency of addition depends upon the relative amount of the bacterial problem. Slug additions may be made to	to its use, syst ms should be cleaned to remove t
mended. Repeat initial dosage until control is evidentions of <u>A. 8</u> to <u>11.9</u> fluid ounces of <u>Algar</u> per 1000 gallons of water should be employed each 1 to frequency of addition depends upon the relative amount of the bacterial problem. Slug additions may be made to	deposits. An initial slug dose of/.1to1
mended. Repeat initial dosage until control is evidentions of <u>A. 8</u> to <u>11.9</u> fluid ounces of <u>Algar</u> per 1000 gallons of water should be employed each 1 to frequency of addition depends upon the relative amount of the bacterial problem. Slug additions may be made to	Aigaecide No. 230 per 1000 (
per 1000 gallons of water should be employed each 1 to frequency of addition depends upon the relative amount of the bacterial problem. Slug additions may be made to	- mended. Repeat initial dosage until control is eviden
frequency of addition depends upon the relative amount of the bacterial problem. Slug additions may be made to	tions of 4 8 to 11 9 fluid ounces of Alga
of the bacterial problem. Slug additions may be made to	
	frequency of addition depends upon the relative amount
collection trays of the airwash system.	of the bacterial problem. Slug additions may be made to
	collection trays of the airwash system.

### STORAGE AND DISPOS

The continuous materials food or field by storage or de justal

STORAGE. On not stack more than four drums high. Leaking or damaged drums for disposel. Spills should be absorbed in sawdust or sand and discretizance closed when not in use.

PESTICIDE DISPOSA() whates resulting from the use of this product in priving waste disprisal facility.

CONTAINER DISFOSAL

 P(A)(10) Triple rinse to enquivalent. Theo offer for my ling minimization is partially landful, by incineration, or in allowed by state and local authoritics.
 xmc/m.

RAETAL Triple rose on represent. Then offer for recycling or recondition carriery landfor on by other procedures approved by state and local author

Manufac		tured by:	
Bonco Mfg. Corp.			
P. O. Box 521	· · · · · · · · · · · · · · · · · · ·		
<u> Jefferson, GA</u>	30549		
EPA REG. NO.		EPA EST.	

NET CONTENTS .