



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

January 9, 2017

Dr. Matthew Brooks
Authorized Representative for Diversified Waterscapes
c/o Ag-Chem Consulting
12208 Quinque Lane
Clifton, VA 20124

Subject: Label Amendment – Addition of me-too molluscicide use
Product Name: Formula F-30 Algae Control
EPA Registration Number: 27588-2
Application Date: 4/26/2016
Decision Number: 516968

Dear Mr. Brooks:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Page 2 of 2
EPA Reg. No. 27588-2
Decision No. 516968


Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. If you have any questions, please contact Lindsay Roe by phone at 703-347-0506, or via email at roe.lindsay@epa.gov.

Sincerely,

A handwritten signature in black ink that reads "Tony Kish". The signature is written in a cursive style with a large, stylized initial "T".

Tony Kish, Product Manager 22 Fungicide Branch
Registration Division (7505P)
Office of Pesticide Programs

Enclosure



MASTER LABEL



Certified to NSF/ANSI 60

FORMULA F-30™

ALGAE CONTROL

ALGAECIDE/BACTERICIDE†

FOR USE IN LAKES, PONDS, RESERVOIRS, CANALS, LAGOONS, SWIMMING POOLS, SPAS, HOT TUBS AND OTHER LISTED WATER SYSTEMS (*Non public Health)

ACTIVE INGREDIENT:	
Copper Sulfate Pentahydrate*	10.9%
CAS No. 7758-99-8	
OTHER INGREDIENTS:	89.1%
TOTAL:	100.0%

*Metallic Copper Equivalent: 2.7%

EPA Reg. No. 27588-2

EPA Est. No. 27588-CA-01

Net Contents: 1 Quart, 1 Gallon, 2.5 Gallons, 5 Gallons, 30 Gallons, 55 Gallons, 275 gallon tote

Manufactured by:
 Diversified Waterscapes, Inc.
 LAGUNA NIGUEL, CA 92677

KEEP OUT OF REACH OF CHILDREN
 CAUTION

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing.

FIRST AID	
IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do NOT induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact INFOTRAC at 1-800-535-5053, 24 hours a day seven days a week for emergency medical treatment information.	

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Mixers, loaders and applicators must wear the following: Long-sleeved shirt, long pants, and shoes and socks.

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove clothing/PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Waters treated with this product may be hazardous to aquatic organisms. Treatment of aquatic weeds and algae can result in oxygen loss from decomposition of dead algae and weeds. This oxygen loss can cause fish and invertebrate suffocation. To minimize this hazard, do not treat more than ½ of the water body to avoid depletion of oxygen due to decaying vegetation. Wait at least 14 days between treatments. Begin treatment along the shore and proceed outwards in bands to allow fish to move into untreated areas. Consult with the State or local agency with primary responsibility for regulating pesticides before applying to public waters, to determine if a permit is required.

Certain water conditions including low pH (≤ 6.5), low dissolved organic carbon (DOC) levels (3.0 mg/L or lower), and "soft" waters (i.e., alkalinity less than 50 mg/L), increases the potential acute toxicity to non-target aquatic organisms.

NOTE: Trout, koi and other species of fish may be killed at the application rates stated on this label. However, fish toxicity generally decreases when the hardness of the water increases. Consult your state fish and game agency before applying this product to public waters.

PRODUCT INFORMATION

Formula F-30 when properly applied, will control a broad spectrum of algae and suppress non-public health bacteria and bacteria that cause taste and odor problems in impounded waters, lakes, ponds, lagoons, wastewater lagoons, reservoirs, livestock watering systems, potable water supplies*, sedimentation basins, swimming pools, spas, hot tubs and other water systems as described in this label. Formula F-30 may be also be used for suppression of non-public health bacteria and toxic gases resulting from bacterial action in sewage lagoons, feedlot run off pits, animal confinement facilities and other ponds containing organic matter or algae.

*Potable Water Supplies = Water destined to be used as drinking water.

Especially effective results have been obtained against spirogyra, cladophora, vaucheria, chara, microcystis, ulathrix and oscillatoria.

If treated water is to be used as a source of water for human consumption, the metallic copper residue must not exceed 1 part per million (ppm). This water must receive additional and separate potable water treatment.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

The maximum application rate per application is 1 ppm of metallic copper for all use sites.

The minimum retreatment interval is 14 days. No more than ½ of any body of water that may contain fish may be treated at one time.

SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity), and the method of application (e.g., ground, aerial, airblast, chemigation) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

DROPLET SIZE: Apply only as a medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

WIND SPEED: Do not apply at wind speeds greater than 15 mph. Only apply this product if the wind direction favors on-target deposition (approximately 3 to 10 mph), and there are no sensitive areas within 250 feet downwind.

TEMPERATURE INVERSIONS: If applying at wind speeds less than 3 mph, the applicator must determine if a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

OTHER STATE AND LOCAL REQUIREMENTS: Applicators must follow all state and local pesticide drift requirements regarding application of copper compounds. Where states have more stringent regulations, they must be observed.

EQUIPMENT: All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

For aerial applications:

- The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.
- Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the water surface unless a greater height is required for aircraft safety.
- When applications are made with a crosswind, the swath must be displaced downwind. The applicator must compensate for this displacement at the up and downwind edge of the application area by adjusting the path of the aircraft upwind.

For ground boom application:

- Do not apply with a nozzle height greater than 4 feet above the crop canopy.

USE IN CONTROL OF ALGAE, NONPUBLIC HEALTH BACTERIA, AND BACTERIA THAT CAUSE ODOR PROBLEMS

For effective algae control, proper chemical concentration should be maintained for a minimum of three hours contact time. The application rates in the chart are based on a static or minimal flow situation. Where significant dilution or loss of water from unregulated inflows or outflows (raceways) occur within a three hour period, Formula F-30 may have to be metered in.

- Identify the algae growth present as one of the following types: Planktonic (suspended), Filamentous (mat-forming), or Chara/Nitella.
- Determine the surface acreage (1 acre = 43,560 sq. ft.) and average depth of infected area.
- Refer to the **"DOSAGE TABLE"** to determine gallons of Formula F-30 to apply per surface acre.

HOW TO APPLY FORMULA F-30

Formula F-30 may be diluted with 5 parts water to 1 part Formula F-30 and coarse sprayed from shore. Application may also be done by pouring a measured amount of Formula F-30 directly from the container into the lake, pond, reservoir or irrigation canal to be treated at several points in the total water area. Generally, 1 gallon of Formula F-30 will treat 325,851.6 gallons (1 acre foot) of water to 0.1 ppm metallic copper. Apply Formula F-30 when algae growth first appears and the temperature of the water to be treated exceeds 60°F.

Larger bodies of water can be treated from the back of a moving boat by dragging a feeder hose behind the boat across the body of water.

Treatment of aquatic weeds and algae can result in oxygen loss from decomposition of dead algae and weeds. This oxygen loss can cause fish and invertebrate suffocation. To minimize this hazard, do not treat more than ½ of the water body to avoid depletion of oxygen due to decaying vegetation. Wait at least 14 days between treatments. Begin treatment along the shore and proceed outwards in bands to allow fish to move into untreated areas.

HOW TO ESTIMATE GALLONS OF WATER/ACRE FEET

Measure Length (L), Width (W) and average Depth (D) in feet. Rectangular or Square shaped bodies of water: **L x W x D x 7.5 = approximate gallons**. Circular or Elliptical bodies of water: **L x W x D x 5.9 = approximate gallons**. Total gallons divided by 325,851.6 (gallons per acre feet) = Acre Feet.

DOSAGE TABLE

GAL. Formula F-30 Per Acre-foot Water	PPM Copper Sulfate Pentahydrate	PPM Metallic Copper
0.5 gal.	0.20	0.05
1.0 gal.	0.40	0.10
2.0 gal.	0.81	0.20

During periods of heavy algae growth, repeat application may be needed. If algae is still present after two weeks repeat the application as directed.

SPECIFIC INSTRUCTIONS

TO CONTROL ALGAE, NONPUBLIC HEALTH BACTERIA, and BACTERIA THAT CAUSE ODOR PROBLEMS IN Irrigation Reservoirs, Impounded Waters, Lakes, Ponds, Lagoons, Reservoirs, Livestock Watering Systems, Potable Water Supplies*, Sedimentation Basins, and Ornamental Water Features or Fountains: For fish-bearing lakes, ponds, drinking water reservoirs, irrigation canals and other applications, apply at the rate of 1 gallon Formula F-30 per acre foot of water for preventive treatment of algae and nonpublic health bacteria. This will yield a concentration of 0.1 ppm metallic copper.

*Potable Water Supplies = Water destined to be used as drinking water.

If algae are present, treat at the rate of 2 gallons of Formula F-30 per acre foot of water. This will yield a concentration of 0.2 ppm metallic copper.

For applications without fish, apply at the rate of up to 2.5 gallons of Formula F-30 acre foot of water. This will yield a rate of 0.25 ppm metallic copper.

Do not exceed 10 gallons of Formula F-30 per acre foot of water (1.0 ppm metallic copper) under any circumstances for water destined for use as drinking water. Formula F-30 may be poured into the water manually after calculating the volume of water to be treated and measuring the quantity of Formula F-30 necessary to attain the desired concentration or by using an automated dispenser calibrated to release the required amount. For best results, disperse Formula F-30 evenly throughout the body of water on a sunny day when algae are near the surface. Do not apply Formula F-30 to water with less than 50 ppm alkalinity. Do not treat more than half of the water body at a time.

TO CONTROL ALGAE or NONPUBLIC HEALTH BACTERIA and BACTERIA THAT CAUSE ODOR PROBLEMS IN Open Channel Irrigation Conveyance Systems, Ditches, Canals and Similar Irrigation Conveyances: To prevent algae growth, apply 1 gallon of Formula F-30 per acre foot of water to yield a rate of 0.1 ppm metallic copper in the water.

If algae are present, apply 2 gallons of Formula F-30 per acre foot of water to yield 0.2 ppm metallic copper. If this application is not effective make next application at a higher rate. Do not exceed 10 gallons of Formula F-30 per acre foot of water (1.0 ppm metallic copper). Minimum retreatment interval is 2 weeks. See Example Calculation Table below for continuous flow rates.

- Formula F-30 should be applied upstream from or directly on the algae or hydrilla when normal delivery of water (clogging of lateral headgates, suction screens, weed screens, and siphon tubes) begins to be interrupted. Delaying treatment could perpetuate a problem causing massing and compacting of plants. Heavy infestations and low flow may cause poor chemical distribution resulting in unsatisfactory control. Under these conditions, increasing water flow rate during application, may be necessary.
- Prior to treatment it is important to accurately determine water flow rates in the absence of weirs, orifices, or similar devices which give accurate water flow measurements, volume of flow may be estimated by the following formula.
Average Width (W) x Average Depth (D) x Velocity (V)[‡] (feet/second) x 0.9 = Cubic Feet per Second (C.F.S.)

Example: 26 x 5 x 1.7 x 0.9 = 198.0 cubic feet per second.

[‡]Velocity (V) is the time it takes a floating object to travel a given distance. Dividing the distance traveled (feet by the time (second) will yield velocity (feet/second). This measurement should be repeated at least three times at the intended application site and then averaged.

- After accurately determining the water flow rate in C.F.S. or gallons/minute, find the corresponding Formula F-30 rate in the chart below:

Example Calculation Table

WATER FLOW RATE		FORMULA F-30 DRIP RATE		
C.F.S.	Gal./Min.	Qts./Hr.	mL/Min.	Fl. Oz./Min.
1	450	1	16	0.5
2	900	2	32	1.1
3	1350	3	58	1.6
4	1800	4	64	2.1

- Calculate the amount of Formula F-30 needed to maintain the drip rate for a period of 3 hours by multiplying Qts./Hr. x 3: ml/Min. x 180: or Fl. Oz./Min. x 180. Dosage will maintain 0.3 ppm metallic copper in the treated water for the 3 hour period. Introduction of the chemical should be made in the channel at weirs or other turbulence creating structure to promote the dispersion of Formula F-30.
- Pour the required amount of Formula F-30 into a drum or tank equipped with a brass needle valve and constructed to maintain a constant drip rate. Readjust accordingly if flow rate changes during the 3 hour period.
- Distance control obtained down the waterway will vary depending upon density of vegetation growth. Periodic maintenance treatments may be required to maintain seasonal control. Wait at least 14 days between applications.

TO CONTROL ALGAE OR NONPUBLIC HEALTH BACTERIA AND BACTERIA THAT CAUSE ODOR PROBLEMS IN Sprinkler, Drip, and Other Types of Irrigation Equipment or Chemigation Systems:

Apply a soon as or prior to visible algae bloom or bacteria growth. Agitation is not required. Do not mix with basic substances. Pour the required amount of Formula F-30 Algae Control to be metered in into a drum or tank equipped with a brass needle valve and constructed to maintain a constant drip rate. Readjust accordingly if flow rate changes during the application period. For best results, Formula F-30 should be applied continuously for the duration of the water application.

Apply 1 to 3 quarts per hour per cubic foot per second depending upon amount of algae presence. See Example Calculation Table above. This will yield 0.3 - 0.9 ppm metallic copper.

TO CONTROL ALGAE AND NONPUBLIC HEALTH BACTERIA AND BACTERIA THAT CAUSE ODOR PROBLEMS IN

Swimming Pools, Spas and Hot Tubs: Apply appropriate rate of Formula F-30 Algae Control at various locations around your pool, spa or hot tub for thorough dispersement.

You will first need to estimate the gallon capacity of your pool:

Measure (L) Length, (W) Width, and (D) Depth in feet.

Square or Rectangular Pools: $L \times W \times D \times 7.5 = \text{Gallons}$

Round or Oval Pools: $L \times W \times D \times 5.9 = \text{Gallons}$

For pools with visible algae, apply at the rate of 1 to 4 Fl. Oz. of Formula F-30 Algae Control per 1,000 gallons of water, depending upon amount of algae presence. Use lower rate for less severe algae presence and higher rate for extremely severe presence. This will yield a rate of 0.25 to 1.0 ppm of metallic copper. Do not apply more than 4 Fl. Oz. (1.0 ppm) per 1,000 gallons of water per application. Every 14 days, test the copper level using a standard commercial swimming pool copper test kit.

For maintenance, apply a dosage of 15 mL of Formula F-30 Algae Control per 1,000 gallons of water every two weeks which will yield a rate of 0.125 ppm of metallic copper.

For spas and hot tubs with visible algae, apply at the rate of 3 mL per 100 gallons of water (0.25 ppm metallic copper).

For maintenance, apply 1.5 mL per 100 gallons of water (0.125 ppm metallic copper) every two weeks.

RESTRICTION:

Treated pool effluent should not be discharged where it will drain into lakes, streams, ponds or public water.

BACTERIA ODOR CONTROL In Sewage Lagoons and Pits (except California)

Application rates may vary depending on amounts of organic matter (sewage) in lagoons and pits. Application should be done by pouring FORMULA F-30 directly from the container into the pit, lagoon or transfer line. Several application points speed up dispersal. Use 1 gallon of FORMULA F-30 in 325,851.6 gallons (1 acre foot) to yield a rate of 0.1 ppm of metallic copper. Non-public health bacteria and odors should be noticeably reduced in 1 to 2 weeks. Repeat application when odor reoccurs. Minimum retreatment interval is 14 days. See subsections below for instructions for specific types of sewage lagoons or pits.

For Feedlot Run-Off Lagoons (except California): Add a portion of the required dosage of FORMULA F-30 around the lagoon to speed dispersal of the product. A minimum of two applications per year (spring and fall) is recommended. Additional applications may be required as needed when the lagoon is pumped.

For Animal Confinement Pits (except California): If pits are located under the confinement buildings, add FORMULA F-30 directly to these pits. If the pits are outside, add FORMULA F-30 to the transfer line to the pit.

For Other Organic Sludges (except California): FORMULA F-30 must be thoroughly mixed with sludge.

MOLLUSCIDIDE**OPEN WATERS: To Control Quagga and Zebra Mussels in Lakes, Ponds, Lagoons, Reservoirs, and Potable Water Supplies*:**

In open or slow-moving, quiescent waters use as a curative measure, i.e., when mussels (veliger, juvenile or adult) have been detected. FORMULA F-30 is miscible in water and has ionic diffusion properties that cause it to readily disperse throughout the water column. Apply near the water surface and allow to disperse, or where means exist, deliver via hose and pump to the depths, sites, and surfaces of worst infestation. When applying to large areas, dispense along a route with gaps no greater than 200 feet. When fish are present, do not treat more than one-half of the body of water at a time, starting near one shore and moving outward in bands so as to allow fish to move away. When treating half of a body of water, the second half must not be treated within 14 days from the last treatment.

For effective control of adult and juvenile mussels, apply at the recommended rate of 4.0 to 32.0 parts per million FORMULA F-30 (i.e., 4.0 to 32.0 gallons of FORMULA F-30 per million gallons of water) to yield a rate of 0.120 to 0.960 mg/L (ppm) metallic copper. Do not exceed 1.0 mg/L (equivalent to 33.33 ppm FORMULA F-30) metallic copper in any single application or in the treated water (background +applied). Allow at least 4 days for mortality to occur. Colder water temperature may require longer exposures and doses closer to the high end of the allowable range. Within the half of the water body being treated repeated applications are permissible if needed to maintain lethal concentrations of copper for sufficient time period. When reapplying, do not exceed a resulting concentration of 1.0 mg/L of metallic copper (background +applied copper) in the treated water. Do not treat the second half of the body of water with 14 days of the last treatment of the first half.

Effective control can also be achieved by longer exposures (e.g., 5-30 days) at lower doses (2.0 to 10.0 parts per million FORMULA F-30, to yield a rate 0.06 to 0.30 mg/L (ppm) metallic copper). Repeat doses are permissible and may be required for severe infestations. When reapplying, do not exceed a resulting concentration of 1.0mg/L (ppm) metallic copper in the treated water (background + applied).

* water destined to be used as drinking water (this water must receive additional and separate potable water treatment).

Dose Rate for Molluscicide FORMULA F-30 in Open Waters (LOW DOSES)

Acres	Depth (ft)	Acre-Ft to Treat	Millions Gallons to Treat	Desired ppm, FORMULA F-30	Desired ppm, as Copper	FORMULA F-30 Dose Rate (gals)	Desired ppm, FORMULA F-30	Desired ppm, as Copper	FORMULA F-30 Dose Rate (gals)
0.1	3	0.3	0.1	2.0	0.06	0.2	4.0	0.12	0.4
0.5	3	1.5	0.5	2.0	0.06	1.0	4.0	0.12	2.0
1	3	3.0	1.0	2.0	0.06	2.0	4.0	0.12	4.0
1	6	6.0	2.0	2.0	0.06	4.0	4.0	0.12	8.0
10	3	30	10	2.0	0.06	20	4.0	0.12	40
10	4.5	45	15	2.0	0.06	30	4.0	0.12	60
10	6	60	20	2.0	0.06	40	4.0	0.12	80
20	3	60	20	2.0	0.06	40	4.0	0.12	80
100	3	300	100	2.0	0.06	200	4.0	0.12	400
1000	3	3000	1000	2.0	0.06	2000	4.0	0.12	4000

Dose Rate for Molluscicide FORMULA F-30 in Open Waters (MEDIUM DOSE)

Acres	Depth (ft)	Acre-Ft to Treat	Millions Gallons to Treat	Desired ppm, FORMULA F-30	Desired ppm, as Copper	FORMULA F-30 Dose Rate (gals)	Desired ppm, FORMULA F-30	Desired ppm, as Copper	FORMULA F-30 Dose Rate (gals)
0.1	3	0.3	0.1	8.0	0.240	0.8	20	0.600	2
0.5	3	1.5	0.5	8.0	0.240	4.0	20	0.600	10
1	3	3.0	1.0	8.0	0.240	8.0	20	0.600	20
1	6	6.0	2.0	8.0	0.240	16	20	0.600	40
10	3	30	10	8.0	0.240	80	20	0.600	200
10	4.5	45	15	8.0	0.240	120	20	0.600	300
10	6	60	20	8.0	0.240	160	20	0.600	400
20	3	60	20	8.0	0.240	160	20	0.600	400
100	3	300	100	8.0	0.240	800	20	0.600	2000
1000	3	3000	1000	8.0	0.240	8000	20	0.600	20000

Dose Rate for Molluscicide FORMULA F-30 in Open Waters (MAXIMUM DOSE)

Acres	Depth (ft)	Acre-Ft to Treat	Millions Gallons to Treat	Desired ppm, FORMULA F-30	Desired ppm, as Copper	FORMULA F-30 Dose Rate (gals)
0.1	3	0.3	0.1	33.3	1.0	3.33
0.5	3	1.5	0.5	33.3	1.0	16.65
1	3	3.0	1.0	33.3	1.0	33.3
1	6	6.0	2.0	33.3	1.0	66.6
10	3	30	10	33.3	1.0	333
10	4.5	45	15	33.3	1.0	499.50
10	6	60	20	33.3	1.0	666
20	3	60	20	33.3	1.0	666
100	3	300	100	33.3	1.0	3330
1000	3	3000	1000	33.3	1.0	33300

For references:

1 acre-foot = 325,851 gal

1 million gal = 3.07 acre-feet

1 hectare= 2.47 acres

1 meter = 3.28 feet

1ppm (1 part per million) = 1 mg/L and/or 1 gal per million gallons

1 gal= 3,785 mL

When calculating dose rates for a given volume of water, achieve a desired concentration of metallic copper in the water to be treated by using the following general formula:

$$\frac{\text{Gallons of applied FORMULA F - 30}}{\text{Million Gallons to be treated}} \times 0.03 = \text{parts per million Copper in the Treated Water}$$

For example treating 3 million gallons with 9.0 gallons of FORMULA F-30 will yield a final copper dose of: (9.0 gals/ 3 million gallons) x 0.030= 0.09mg/L as copper = 0.09 ppm as copper

Always use volumetric measurement devices that are calibrated in accordance with manufacturer specifications.

Flowing Waters: To Control the Mollusk Pests Quagga and Zebra Mussels in flowing potable water supplies*; canals; ditches; aqueducts; and equipment/structures that deliver the treated water directly to publicly owned water treatment facilities to include pipes, intake structures, gatehouses, screens, pumping stations, weirs, penstocks: In flowing waters, use when mollusks (veliger, juvenile, or adult) have been detected. May be used as curative measure when adult or juvenile mollusks are present, or as a preventative measure (to inhibit colonization) when adults and/or planktonic larval mollusks have been detected.

* water destined to be used as drinking water (this water must receive additional and separate potable water treatment).

USEPA MASTER LABEL – FORMULA F-30

FORMULA F-30 may be used continuously on flowing waters as a means of preventing further spread and colonization of mollusks. Start the continuous application when mollusks are present and end application when mollusks are no longer present. Use a metering pump to apply a continuous dose so as to achieve a final dilution of 2.0 to 32 ppm FORMULA F-30 (0.06 to 0.96 ppm metallic copper, or mg/L). Do not exceed 1.0 ppm free metallic copper (background+ applied) in the flowing water, equivalent to 33.3 ppm, as FORMULA F-30. If adult mollusk are already present, allow at least 4 days for mortality to occur, or longer for very well-established populations where adults appear in clumps. For most situations satisfactory control will be obtained at a continuous dose of 2.0 to 10.0 ppm FORMULA F-30 (i.e., 0.06 to 0.30 mg/L (ppm) metallic copper). Colder water temperatures may require longer exposure and dose rate closer to the high end of the allowable range.

Once the initial infestation has been cleared from surfaces, a continuous maintenance dose of 1.2 to 4.0 ppm FORMULA F-30 (yielding a metallic copper concentration of 0.036 to 0.120 ppm) can be used to prevent further colonization.

Example Dose Rates for FORMULA F-30 in Flowing Waters (LOW DOSE)

cfs	Gal/min	MGD	Desired ppm, FORMULA F-30	Desired ppm, as Copper	FORMULA F-30 Feed Rate (fl oz/min)	FORMULA F-30 Feed Rate (ml/min)	Desired ppm, FORMULA F-30	Desired ppm, as Copper	FORMULA F-30 Feed Rate (fl oz/min)	FORMULA F-30 Dose Rate (ml/min)
1	449	0.65	2.0	0.06	0.12	3.37	4.0	0.12	0.214	6.74
1.55	696	1.0	2.0	0.06	0.17	5.21	4.0	0.12	0.353	10.44
3	1,346	1.9	2.0	0.06	0.33	10.1	4.0	0.12	0.674	20.21
4	1,795	2.6	2.0	0.06	0.46	13.47	4.0	0.12	0.91	26.95
5	2,244	3.2	2.0	0.06	0.57	16.82	4.0	0.12	1.13	33.68
10	4,488	6.5	2.0	0.06	1.13	33.68	4.0	0.12	2.28	67.37
15.47	6,943	10	2.0	0.06	1.76	52.11	4.0	0.12	3.47	104.23
50	22,442	32	2.0	0.06	5.68	168.23	4.0	0.12	11.38	336.85
100	44,883	65	2.0	0.06	11.37	336.65	4.0	0.12	22.77	673.70
155	69,429	100	2.0	0.06	17.61	521.109	4.0	0.12	35.27	1042.25
1000	448,830	646	2.0	0.06	112.9	3,366.54	4.0	0.12	227.87	6733.07

Example Dose Rates for FORMULA F-30 in Flowing Waters (MEDIUM AND HIGH DOSE)

cfs	Gal/min	MGD	Desired ppm, FORMULA F-30	Desired ppm, as Copper	FORMULA F-30 Feed Rate (fl oz/min)	FORMULA F-30 Feed Rate (ml/min)	Desired ppm, FORMULA F-30	Desired ppm, as Copper	FORMULA F-30 Feed Rate (fl oz/min)	FORMULA F-30 Dose Rate (ml/min)
1	449	0.65	10	0.30	0.54	15.72	30	0.90	1.70	50.37
1.55	696	1.0	10	0.30	0.83	24.44	30	0.90	2.63	77.96
3	1,346	1.9	10	0.30	1.59	47.22	30	0.90	5.11	150.93
4	1,795	2.6	10	0.30	2.13	62.96	30	0.90	6.81	201.85
5	2,244	3.2	10	0.30	2.67	78.70	30	0.90	8.52	251.85
10	4,488	6.5	10	0.30	5.31	157.22	30	0.90	17.02	503.70
15.47	6,943	10	10	0.30	8.22	242.59	30	0.90	26.30	777.78
50	22,442	32	10	0.30	26.67	787.04	30	0.90	85.19	2,516.66
100	44,883	65	10	0.30	53.15	1572.22	30	0.90	170.19	5,033.33
155	69,429	100	10	0.30	82.22	2433.33	30	0.90	262.96	7,787.03
1000	448,830	646	10	0.30	531.48	15,729.61	30	0.90	1,701.85	50,333.28

MGD= Million Gallons per Day
cfs= Cubic Feet per Second

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Keep container closed when not in use. Open dumping is prohibited.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinsate that cannot be used according to label directions or chemically reprocessed should be disposed of according to procedures approved by Federal, State or Local Disposal Authorities.

(FOR NONREFILLABLE CONTAINERS, EQUAL TO OR LESS THAN 5 GALLONS)

CONTAINER DISPOSAL: Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after flow begins to drip. Repeat this procedure two more times. Offer for recycling if available, or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(FOR NONREFILLABLE CONTAINERS GREATER THAN 5 GALLONS)

CONTAINER DISPOSAL: Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(FOR REFILLABLE CONTAINERS)

CONTAINER DISPOSAL: Refillable containers: Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure 2 more times. Offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

NOTICE – To the extent consistent with applicable law, neither the manufacturer nor the seller makes any warranty expressed or implied, concerning the uses of this product other than indicated on the label. Buyer assumes all risk of use of this material when such use is contrary to label instructions. Read and follow the label directions carefully.