

'N F


WEED

WESTERN
KW
 SUPER
 KELLY

WHEN TO APPLY:

MAY BE APPLIED ANY TIME OF YEAR FOR BEST RESULTS APPLY WHEN WEEDS ARE GROWING ACTIVELY.

HOW TO APPLY FOR A BETTER LAWN



1 Measure area to be covered (for example 50 sq. ft. requires 1/2 lb. of product). Best results will be obtained if lawn is cut two to three days before the application. For maximum effectiveness water lawn thoroughly the day before application. Application in the evening or early morning will be most effective.

2 Immediately before spreading water the lawn well. If sections dry before they can be reached, re-wet, but avoid washing off material already applied. Do not water lawn for at least 24 hours after application. Do not apply on days when temperature is over 85°.

3 Set spreader as recommended below. Fill spreader where spilled material will not fall on lawn. Shake spreader to settle the material. Make application in two crisscross coverages as illustrated. Spreader settings below are for crisscross applications.

SUGGESTED SPREADER SETTINGS

Spreading rate may vary with the age, model, and condition of the spreader; check your spreader by a trial over a known-size area. (22 lb. bag covers 5000 sq. ft.)

Lawn Beauty		Scott		Modern		Central		Sears	
MODEL	SETTING	MODEL	SETTING	MODEL	SETTING	MODEL	SETTING	MODEL	SETTING
18-57	7 1/4	35-8	7 1/2	18 in.	6 1/4	20 in.	8	452-19191	6 1/4
		35-3	7						

Above settings are for crisscross applications. If your spreader is not included above, set spreader opening at lowest setting which gives even flow, usually about 1/3 open. Spread in one direction and then crisscross. Continue crisscrossing until proper amount is applied. Broadcast spreaders are not recommended.

WEEDS CONTROLLED

black medic
 buttercup
 chickweed
 cinquefoil
 clover
 dandelion
 docks

English ivy
 heal-all
 henbit
 knotweed
 lambsquarters
 mallow
 morning glory

mustard
 Oxalis
 pennywort
 pepperweed
 pigweed
 plantain

prostrate spurge
 purslane (pusley)
 ragweed
 red sorrel
 sheepsorrel
 smartweeds
 Spanish nettle

spotted spurge
 thistle
 velvet leaf
 vervain
 Yarrow

NOTICES

SEEDING:

Apply to established lawns only. May be used on reseeded lawns six weeks after reseeding.

SHRUBS AND FLOWERS

Avoid making applications on flowers, shrubs, or other ornamental plantings. Do not apply on or near young trees. Prevent drift of product to desirable plants.

May injure carpetgrass, St. Augustine, bentgrass, dichondra, centipede, or newly seeded lawns.

Do not apply to areas underlaid by roots of desirable trees and shrubs.

Repeat applications may be necessary on difficult to kill weeds.

GUARANTEE: Manufacturer guarantees this product to give results as outlined on this bag or the purchaser may get his money refunded by the manufacturer. No other express warranties of any kind are made, except to conform to the promises and affirmation of facts made on the label. The warranty of merchantability, all other implied warranties and liability of the manufacturer without fault are expressly disclaimed.

Buyer assumes all risk of personal injury or damage to property, except for negligent mislabeling of the product. All claims must be presented to manufacturer in writing, within thirty days from the date a breach is discovered and an action thereon must be commenced within one year after the cause of action has occurred.


KW SUPER WEED'N FEED

Proposed Weed and Feed Label for:

GREEN UP
RED ROOSTER
WILSHIRE
ROLLING ACRES
MAGIC
EASY LIFE
TOP'S ALL
TAM-O-SHANTER
MAGIC GREEN
HAPPY PATCH

Net Contents
22 Pounds

Cover 5,000
Square Feet



22 Lbs. New Weight

GREEN UP


WEED AND FEED

10-6-4

Contains 2,4-D

USDA Reg. No. _____

December 27, 1967



GREEN UP
WEED AND FEED
contains 2,4-D.

Controls broadleaf weeds in lawns.

Plant Food

Non-burning Long Lasting

CAUTION: Keep out of reach of children.
See back panel of bag for additional cautions.

ACTIVE INGREDIENTS AS HERBICIDE

Butoxyethanol ester of 2,4-Dichlorophenoxyacetic acid* ... 1.44%

INERT INGREDIENTS:98.56%

*2,4-Dichlorophenoxyacetic acid equivalent 1.0% by weight.

PLANT FOOD ANALYSIS:

Total Nitrogen (N) 10%
10.0% Ammoniacal Nitrogen

Available Phosphoric Acid (P₂O₅) 6%

Soluble Potash (K₂O) 4%

Potential Acidity 918 lbs., Calcium Carbonate Equivalent per ton

USDA Reg. No. _____

A Product of:
CUSTOM FARM SERVICES, INC.
Atlanta, Georgia

DIRECTIONS:

This product will control dandelions, plantain, buckhorn, mustard and many other weeds in lawns. It will help provide healthy grass while controlling weeds. It will also help control clover.

The contents of this bag should be applied evenly with a mechanical spreader on 5,000 square feet (50 feet by 100 feet). For best results use the following suggested spreader settings.

SPREADER MAKE	SETTING
Scott's	6
Modern	5
Vaughan	6
Lawn Beauty	5
Scars	5
Jackson	6
Lawn Master	6
Cyclone Broadcaster	6

Do not reseed for at least two weeks after application. Do not apply to a newly seeded lawn until the seedling grasses have formed crowns (usually after 3 mowings).

Bentgrass and clovers may be injured by this treatment. Do not use on susceptible Southern grasses such as St. Augustine.

Do not water for at least 24 hours after application. If rain occurs during 24 hours after application a second spreading may be necessary.

CAUTION:

Avoid inhaling dust and contact with skin, eyes or clothing.

Use on lawns only. Do not use for gardens or on ground covers such as pachysandra or blue myrtle. Do not apply directly to or allow dust to contact flowers, vegetables, fruits, shrubs, or trees. Do not store near seeds, fertilizers, insecticides or fungicides.

Destroy empty container in a safe place. Do not burn.

WARRANTY

The manufacturer warrants that the composition of this product conforms to the chemical description given in the ingredient statements and that the product is suitable for the use described when used according to directions. The buyer assumes any risk to persons or property arising out of the use or handling and accepts the product under these conditions.