

**UNICORN  
FLEA AND TICK POWDER #4  
FOR  
CATS AND DOGS**

**Kills Fleas, Lice and Ticks.**

**ACTIVE INGREDIENTS:**

Carbaryl (1-naphthyl N-methyl-carbamate .....	5.0%
Pyrethrins. ....	0.1%
Piperonyl Butoxide, Technical* .....	1.0%

**INERT INGREDIENTS:** ..... 93.9%

**Total:** ..... 100.0%

\*Equivalent to 0.8% (butylcarbityl)(6-propylpiperonyl) ether and 0.2% related compounds.

**KEEP OUT OF REACH OF CHILDREN**


**CAUTION**

**See Side/Back Panel for Additional Precautionary Statements.**

NET WEIGHT: 6 OUNCES

**MANUFACTURED BY:**

Unicorn Laboratories  
13535 Feather Sound Drive, Suite 400  
Clearwater, FL 34622-5587  
(813) 578-4545



- ☐ READ THE ENTIRE LABEL BEFORE EACH USE.
- ☐ USE ONLY ON DOGS OR CATS.

**PRECAUTIONARY STATEMENTS**

**Hazards to Humans and Domestic Animals**

**CAUTION:** Harmful if swallowed. Avoid breathing of dust when you apply powder. Keep out of reach of children. Do not treat puppies and kittens under twelve (12) weeks of age. Consult a veterinarian before using this product on debilitated, aged, pregnant or nursing animals or animals on medication. Sensitivities may occur after using **ANY** pesticide product for pets. If signs of sensitivity occur, bathe pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

<b>USER SAFETY REQUIREMENTS</b>
<ul style="list-style-type: none"> <li>■ Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product.</li> <li>■ Change clothing as soon as possible after use.</li> <li>■ Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.</li> </ul>

**Statement of Practical Treatment**

**IF IN EYES:** Flush with plenty of water. If irritation persist, see a physician.

Do not use this product on animals simultaneously or within 30 days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides or chemicals. However, flea and tick collars may be immediately replaced.

**NOTE TO PHYSICIAN and VETERINARIAN:** Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine sulfate is antidotal.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**HOW TO USE:** For control of fleas, ticks and lice - Use product only outside or in a well-ventilated area. Wearing household latex or rubber gloves, dust pet liberally by rubbing dust into animals hair. Begin at tail and work toward the head. Take care not to get dust in animals eyes, nose or mouth. Be sure to dust paws and between toes. This product should not be applied by children. Do not use on pregnant dogs. Repeat as needed, but not more often than once weekly.

**STORAGE AND DISPOSAL**

**Storage:** Store in original container in cool, locked storage area.

**Disposal:** Do not reuse container. Empty completely, wrap in newspaper before discarding in trash.