

AUG 11 1992

Mr. Barry Santerre
Unicorn Laboratories
1000 118th Avenue North
St. Petersburg, FL 33716

Dear Mr. Santerre:

Subject: Labeling Amendment
Unicorn Stirofos Flea & Tick Powder
EPA Registration No. 28293-13
Your Submission Dated May 4, 1992

The amendment referred to above, submitted in connection with registration under FIFRA sec 3(c)(7)(A), is acceptable, provided that you:

1. Submit/cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling:
 - a. Under Precautionary Statements the statement should read: "Wear long-sleeved shirt and pants; chemical resistant gloves; shoes and socks."
 - b. Singularize Active Ingredients.
 - c. Under Directions for Use delete "General Directions."
3. Submit five copies for your final printed labeling before you release the product for shipment.

CONCURRENCES

SYMBOL	SURNAME	DATE						
	<i>W. J. ...</i>	<i>Jan 11 1992</i>						

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the labeling is enclosed for your records.

Sincerely,

George T. LaRocca
Product Manager 13
Insecticide-Rodenticide Branch
Registration Division (H7505C)

UNICORN

STIROFOS FLEA & TICK POWDER

**A Ready-To-Use Insecticide Dust to Control
Horn Flies and Lice and in the control of Face Flies
on Beef Cattle and Dairy Cattle and to control
Lice on Swine**

Kills Fleas & Ticks on Cats and Dogs.

Fast Acting, Lasts Seven Days.

ACTIVE INGREDIENTS:

Tetrachlorvinphos: 2-chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate..... 3.0%

INERT INGREDIENTS:..... 97.0%

100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Side/Back Panel for Additional Precautionary Statements.

NET WEIGHT: 6 OUNCES

MANUFACTURED BY:

**Unicorn Laboratories
1002 118th Avenue North
St. Petersburg, Florida 33716**

**EPA REG. NO. 28293-13-AA(E:4/88) ACCEPTED with COMMENTS EST. NO. 28293-FL-1
in EPA Letter dated**

AUG 11 1992
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 28293-13

PRELIMINARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed. Avoid breathing dust. Do not get in eyes, on skin or on clothing. Wash thoroughly with soap and water after handling and before smoking or eating. Avoid contamination of feed and food stuffs. Wear long-sleeved shirt, and pants, shoes and socks.

NOTE TO PHYSICIAN: This product is a cholinesterase inhibitor. Atropine is antidotal. 2-PAM is also antidotal and may be used in conjunction with atropine.

Environmental Hazards

This pesticide is toxic to fish. Keep out of lakes, streams and ponds. Do not contaminate water by disposal of wastes. Apply this product only as specified on this label. Do not contaminate water when disposing of equipment washwaters.

Statement of Practical Treatment

IF IN EYES: Flush with plenty of water. If irritation persists see physician.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

LIVESTOCK DUST

General Directions: Product can be applied by hand or through the use of self-treating dust bags. There is no withholding period from last application to slaughter.

Animal	Insect	Remarks
Beef	Horn Flies	HAND DUSTING: Apply approximately 2 oz of dust by shaker can, rotary duster or by spoon to the upper portions of the back, neck and poll and to the face as an aid in the control of face flies. Rub in lightly to carry the dust beneath the hair. Repeat as necessary. SELF TREATING DUST BAG FORCED USE: Put the product in cotton cloth or double bag or use pre-packed weather proof Cattle dust bag and hand in barn door exits or alleyways leading from animal buildings, salt or mineral blocks, or watering holes. Protect cloth or burlap bags from weather. FREE CHOICE USE: Use the same dust bags as above but place in loafing sheds, holding pens, feedlots, near watering holes or other areas where cattle gather. The free choice aids in the control of lice.
Cattle	Face Flies	
Dairy	Lice	
Cattle		

to each animal with special attention given to the neck and around the ears. Repeat as necessary but not more often than once every 14 days. In severe infestations, both individual animals and bedding may should be applied per 100 sq. ft. of bedding.

DOG AND CAT USE

To Kill Fleas, Ticks and thereby reducing the itching and scratching caused by the bites of these pests: Dust entire animal beginning at back and working toward the head. Make sure to dust the legs, paws and between the toes. Dust animal's bedding and sleeping quarters regularly. Repeat at weekly intervals if necessary.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Do not store in areas of extreme heat or cold. Protect from moisture or extreme humidity.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. Wastes resulting from the use of this product may be disposed of on site or at approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty can into application equipment, then dispose of empty can in a sanitary landfill or by incineration or, if allowed by State and local authorities, by burning. If burned, stay out of smoke. **For Containers one gallon or less** - Do not reuse empty container. Wrap container in several layers of newspaper and discard in trash.