

UNICORN

FLEA AND TICK SPRAY

Kills Fleas & Ticks on Dogs
Repels Flies, Mosquitoes and Gnats
Lanolizes Animals Coat

ACTIVE INGREDIENTS:

1-Naphthyl-N-Methylcarbamate.....	0.50%
Pyrethrin.....	0.06%
Piperonyl Butoxide, Technical*.....	0.12%
N-octyl bicycloheptene dicarboximide.....	0.20%

INERT INGREDIENTS:.....	99.12%
	100.00%

* Consists of 0.094% (Butylcarbityl) (6-Propyl piperonyl) ether and 0.024% related compounds.

Keep Out of Reach of Children


CAUTION

See Back Panel for Additional Caution

NET CONTENTS: 16 OUNCES

MANUFACTURED BY:

Unicorn Laboratories
1002 118th Avenue North
St. Petersburg, Florida 33716



EPA REG. NO. 28293-16(E:8/13/90)

EPA EST. NO. 28293-FL-1

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin and eyes. Wash hands with soap and water after using or handling. Do not use on cats or puppies less than four weeks of age. Keep out of reach of children.

Notice to Physician: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine Sulfate is antidotal.

Statement of Practical Treatment

IF INHALED: Remove victim to fresh air. Apply artificial respiration is indicated.

IF ON SKIN: Remove contaminated clothing and wash affected areas with soap and water.

IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists.

Physical or Chemical Hazard

FLAMMABLE. Contents under pressure. Do not use near fire, sparks or flame. Do not puncture or incinerate. Exposure to temperatures above 130°F may cause bursting.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Remove cap and press down button with hole away from you. Hold container approximately 6 inches from animal. Work from rear of animal towards the head, fluffing the hair to permit the spray to penetrate to the skin. Spray only enough to slightly dampen the hair and skin. Wet ticks with the spray. Do not get spray in eyes or on scrotum.

Use spray on the basis of 4 to 6 seconds for a 5 pound cat and 20 to 30 seconds for a 25 pound dog. Repeat treatment if necessary; but not more than once a week. (The hissing sound and the cool sensation caused by the spray may temporarily frighten nervous pets; however, they are expected to become accustomed to the treatment.) Treat bedding, interior of kennels and other places where ticks and fleas congregate to help prevent reinfestation.

STORAGE AND DISPOSAL

Store in original container in cool, locked storage area.

Replace cap and discard container in trash. Do not incinerate or puncture.