

UNICORN

FLEA & TICK POWDER

**READY TO USE ON:
VEGETABLES ■ GARDENS ■ LAWNS
POULTRY ■ DOGS ■ CATS**

Kills Fleas, Lice and Ticks on Dogs and Cats

ACTIVE INGREDIENTS:

Carbaryl (1-Naphthyl N-methylcarbamate)..... 10.00%

INERT INGREDIENTS:..... 90.00%

100.00%

Keep Out of Reach of Children

CAUTIONS


See Back of Panel for Additional Cautions.

NET WEIGHT: 10 OUNCES

MANUFACTURED BY:

Unicorn Laboratories
1002 118th Avenue North
St. Petersburg, Florida 33716

ACCEPTED
MAY 19 1992
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
this product is registered for the pesticide
use specified under
EPA Reg. No. 28293-18



EPA REG. NO. 28293-18

EPA EST. NO. 28293-FL-1

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Avoid breathing of dust when applying powder. May be harmful if swallowed. Wash hands with soap and water after using.

Statement of Practical Treatment

IF SWALLOWED: Drink 1 or 2 glasses of water and induce vomiting by sticking finger down throat. Never give anything by mouth to an unconscious person. Call Physician.

IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated.

IF ON SKIN: Wash with soap and water.

IF IN EYES: Flush with plenty of water. If irritation persists see Physician.

Notice to Physician: Carbaryl is a cholinesterase inhibitor. Atropine is antidotal.

Environmental Hazards

This product is highly toxic to bees exposed to direct treatment or residual on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

This pesticide is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes.


DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

APPLICATION: Begin application when insect damage is first observed and repeat on a regular 7 to 10 day schedule or as necessary for good insect control.

VEGETABLES:

Corn, Eggplant, Pepper, Tomato: To control Corn Earworms, European Corn Borers, Sap Beetles, Armyworms, Tomato Fruitworms, Flea Beetles, Leafhoppers, Tomato Hornworms, and Stink Bugs - dust the



upper and lower leaf surfaces thoroughly. For better control of Corn Earworms, apply to silks as soon as they appear. Make 3 to 4 applications at 2 to 3 day intervals. Reapply immediately after rain.

Beans: To control Mexican Bean Beetles, Bean Leaf Beetles, Cucumber Beetles, Leafhoppers, Flea Beetles, Japanese Beetles, Corn Earworms, Lygus Bugs, and Lima Bean Pod Borers - dust the upper and lower leaf surfaces thoroughly.

Cucumber, Summer Squash: To control Pickleworms, Melonworms, Cucumber Beetles, Squash Bugs and Flea Beetles - dust the upper and lower leaf surfaces thoroughly. Apply to cucumber and summer squash only when plants are dry.

Potato: To control Colorado Potato Beetles, Leafhoppers, Flea Beetles, and Lygus Bugs - dust the upper and lower leaf surface thoroughly.

APPLICATION TO ABOVE VEGETABLES CAN BE MADE UP TO DAY OF HARVEST. SUBSEQUENT PLANTING OF FOOD AND FEED CROPS IN TREATED SOIL MUST BE MADE WITH CROPS ON THIS OR OTHER CARBARYL LABELS.

LAWNS: To control Chinch Bugs, Armyworms, Ants, Fleas, Sod Webworms, Lawn Moths, Millipedes, and Earwigs - apply 1/2 pound to 200 sq. ft. For best results apply after rain or watering and do not water for at least two days after application. Repeat in two or three weeks if necessary.

POULTRY: Litter Treatment - apply 1/2 pound to 40 sq. ft. to control Northern Fowl Mites, Chicken Mites, Lice, Fleas and Bed Bugs. Use a scoop and scatter by hand, or apply by hand duster. Avoid contamination of nests, eggs, feeding and watering troughs. Do not house birds in treated houses within 7 days of slaughter.

Hand Application to Birds: Use 1/2 pound to treat 100 birds by means of shaker can, or hand duster to control Northern Fowl Mites, Lice, and Fleas. Direct dust to vent and fluff areas. Do not contaminate feed or drinking water. The application of this material cannot be repeated within 4 weeks of each treatment. Hand application to birds is a supplement to litter treatment for control of these pests. Do not apply to poultry within 7 days of slaughter.

DOGS, CATS: How to Use: For best results to kill Fleas, Ticks, and Lice on animals, dust powder liberally over the entire body. Rub dust into hair and around the feet, pads and between the toes so that they are well treated. Take care not to get the powder in animal's eyes, nose or mouth. Dust bedding and kennel and sleeping area regularly. Repeat as needed, but not more often than once weekly. Do not treat kittens less than four weeks old. Do not use on nursing puppies. Do not use on pregnant dogs.

STORAGE AND DISPOSAL

Do not contaminate water, feed or food by storage or disposal.

Storage: Store in original container in cool, locked storage area.

Disposal: Do not reuse empty container. Empty completely, then dispose of in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

