



**PRECAUTIONARY STATEMENT
HAZARDS TO HUMANS & DOMESTIC ANIMALS
CAUTION**

HARMFUL IF SWALLOWED If swallowed, do not induce vomiting. Wash face, hands and arms with soap and water after handling and before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse. Do not breathe spray mist. Wear a pesticide respirator jointly approved by the Mining Enforcement and Safety Administration (formerly the U.S. Bureau of Mines) and by the National Institute for Occupational Safety and Health.

Do not remain in treated area. Ventilate the area thoroughly before re-entry. Do not apply directly on very young animals. Do not use in poultry houses. Remove pets, birds, and cover fish aquariums before spraying. Do not apply in a food processing plant while plant is in operation. When applied in a food processing plant, remove or protect food during treatment. After treatment, wash all equipment, benching, shelving, etc. with potable water where exposed food will be handled.

ENVIRONMENTAL HAZARD

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated. Do not contaminate any body of water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label. Do not reuse empty containers. Destroy by perforating or crushing and burying in a safe place away from water supplies. Return drums to reconditioners.

PHYSICAL HAZARD

Keep container closed. Do not use, pour, spill or store near heat or open flame.

STORAGE AND DISPOSAL

Do not contaminate water, food and feed by storage or disposal. Open dumping is prohibited. Pesticide, spray mixture or rinsate that cannot be used or chemically processed should be disposed of in a land fill approved for pesticides, or buried in a safe place away from water supplies. Triple rinse, or equivalent and offer for recycling, reconditioning or disposal in approved landfill, or bury in a safe place. Consult Federal, State or Local disposal authorities for approved alternative procedures.

Directions For Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

OUTDOOR SPRAY

TO REDUCE THE ANNOYANCE OF FLIES, ADULT MOSQUITOES, GNATS AND SMALL FLYING MOTHS around patios, swimming pools, home yards, and outdoor areas in parks and playgrounds: When the air is still, spray as a fine mist or fog to penetrate hiding places such as grasses, shrubs and ornamentals around homes, picnic areas, stadiums and open air theaters. Treat as large an area as possible surrounding the patio, pool or grounds where insect protection is desired. Repeated applications may be necessary when infestation is heavy. Apply at the rate of 1.5 gallons per acre (4.4 fl. oz. per 1,000 sq. ft.). Do not spray flowers and ornamentals at close range. Allow only the fine mist or "dry" fog to contact desirable plants. Direct spray away from the swimming pool. Application to any body of water is prohibited.



Unicorn

BLITZ RTU SPRAY

Containing
PYRETHRIN
FOR HOME, FARM AND INDUSTRY
USDA ACCEPTED

For Application by Trained Personnel Only

A contact spray for control of listed flying insects in home, restaurants, hotels, food storage warehouses, dairies, meat packing plants and similar establishments with electric mist spraying or fogging equipment. Also outdoor adult mosquito control.

ACTIVE INGREDIENTS:

Pyrethrins	0.10%
Piperonyl butoxide, technical*	1.00%
Isoparaffinic Hydrocarbons	98.90%

* Consists of 0.80% (butylcarbitol) (6-propylpiperonyl) ether and 0.20% related compounds.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Harmful or fatal if swallowed

Read Side Panel For Additional Precautionary Statements

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED Call a physician immediately. If gastric lavage is deemed necessary, undertake if material was taken internally. Do not induce vomiting unless no other treatment is available. Vomiting may cause aspiration pneumonia. If necessary to induce vomiting, give 1 or 2 glasses of water and insert finger in back of throat. Repeat until vomit fluid is clear.
IF INHALED Remove victim to fresh air and apply respiration if indicated.
IF ON SKIN Wash promptly with soap and water. Rinse thoroughly.
IF IN EYES Rinse eyes for at least 15 minutes with water and call a physician immediately.

Unicorn Laboratories
P.O. Box 90014
Houston, Texas 77070

NET CONTENTS GALLONS

E.P.A. No. 28293-X
E.P.A. Est. No. 28293-TX-1

ACCEPTED
With Comments in
EPA Notice of
SEP 09 1982
Unicorn Laboratories
Fungicide, Insecticide, Fec.
as amended, 28293-21
Registered under
EPA Reg. No. 28293-21

**DIRECTIONS (continued)
INDOOR SPRAY**

FOR CONTROL OF FLIES, ADULT MOSQUITOES, GNATS AND SMALL FLYING MOTHS in homes, food establishments, farm buildings and warehouses. Close all windows and stop all ventilators. Pilot lights should be extinguished. Remove pets, birds and cover fish aquariums before spraying. Allow only fine mist or "dry" fog to contact fabric or surfaces that might be affected by the oil base. Direct fine mist or fog upward toward the center of the room and around the sides until the area is filled with fog or mist particles. Keep area closed for 10 to 15 minutes. Do not remain in treated areas. Ventilate the area thoroughly before re-entry. Clean up and destroy fallen insects.

CONTACT SPRAY-FOR USE WITH SPRAYERS THAT GIVE COARSE WETTING SPRAY PARTICLES TO KILL COCKROACHES AND ACCESSIBLE STAGES OF GRANARY INSECTS: GRANARY WEEVILS, CONFUSED FLOUR BEETLES, RED FLOUR BEETLES, INDIAN MEAL MOTH LARVAE AND ADULTS, CADELLES, GRAIN MITES AND SAW-TOOTHED GRAIN BEETLES: Cover or remove exposed food. Spray outside walls of grain bins, floorings, about the base of milling machinery, walls, baseboards, areas behind cabinets, around pipes, and inside surfaces of boxcars and trucks using 1 gallon of spray per 1,000 sq. ft. of space. Apply in small amounts directly into cracks and crevices using equipment capable of delivering a pin stream in points between equipment and floors, openings leading to voids and hollow spaces in walls, equipment legs and bases, conduits, motor housings and junction or switch boxes. Prior to application, de-energize electrical equipment. After application, allow time for oil to evaporate before starting motors or placing switches in "on" position. Repeat applications as necessary. Do not remain in treated areas. Ventilate the area thoroughly before re-entry. Infested stored food products should be fumigated or treated by other effective methods using an approved product intended for this purpose.

CONTACT AND SPACE SPRAY-TO KILL COCKROACHES, SILVERFISH, EARWIGS, ANTS, SPIDERS IN SHELTERED SITUATIONS: Adjust spraying device to deliver a coarse wet spray. Apply as a contact spray into all cracks and crevices in woodwork, walls, floors, underneath sinks, behind pipes and in all places where these insects shelter. Contact as many insects as possible. Do not allow spray to contact stored food. Then adjust sprayer to deliver aerosol size particles. Apply as a space spray at the rate of 3 to 6 fl. oz. per 1,000 cu. ft. of room space. Repeat dosage as necessary for control. Leave the room closed for 1 hour. Do not remain in treated area. Ventilate the area thoroughly before re-entry.

FOR WASPS AND SPIDERS: Spray insects directly and apply heavy spray to nests and webs. The most effective wasp control is obtained by spraying after dark when all insects have returned to the nest.

FOR CONTROL OF CARPET BEETLES: Clean all rugs, carpets and fabrics to be treated. Spray both surfaces of carpets and floor space beneath evenly with a fine mist spray. Particular attention should be paid to cracks and crevices around the baseboards.

ANIMAL SPRAY FOR HORSES AND DAIRY AND BEEF CATTLE: For protection against stableflies, hornflies and houseflies attacking animals, spray the animal against the grain of the hair, directing a fine mist around the base of the horns, on the legs and along the sides and belly and over the back. Do not spray udders. Apply just enough to wet the ends of the hair but not enough to wet the hide. One or two ounces of spray is sufficient to treat an adult cow. Apply once daily or as necessary.

BUYER ASSUMES ALL RISKS OF USE, STORAGE OR HANDLING OF THIS MATERIAL NOT IN STRICT ACCORDANCE WITH THE DIRECTIONS GIVEN HERewith