

UNICORN

DAIRY & BEEF CATTLE DUST

Controls Hornflies on Dairy Cattle  
and  
Hornflies, Lice and Ticks on Beef Cattle

**ACCEPTED**  
JUL 20 1989  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under  
EPA Reg. No. 28293-106

ACTIVE INGREDIENTS:

- X Malathion : 0,0-Dimethyl dithiophosphate of  
diethyl mercaptosuccinate ..... 4.0%
- Methoxychlor, Technical ..... 5.0%
- (equivalent to 4.4% 2,2-bis(p-methoxyphenyl)  
1,1,1-trichloroethane and 0.6% other  
isomers and reaction products

INERT INGREDIENTS: ..... 91.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION


See Back Panel for Additional Precautions

Net Weight:                      Pounds

Manufactured By

UNICORN LABORATORIES

ST. PETERSBURG, FLORIDA 33716



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid breathing dust. Avoid contact with skin; wash thoroughly after using. Avoid contamination of feed and food stuff. Do not apply within of 5 hours before milking or during milking. Do not use on poultry or in poultry houses. Do not treat animals less than one month of age.

STATEMENT OF PRACTICAL TREATMENT

- IF SWALLOWED: Call physician at once. Drink 2 glasses of water and induce vomiting by touching back of throat with finger.
- IF IN EYES: Flush with plenty of water. If irritation persists, get medical attention.
- IF INHALED: Remove victim to fresh air.
- IF ON SKIN: Remove contaminated clothing. Wash areas of contact with soap and water.

ENVIRONMENTAL HAZARDS

"This pesticide is toxic to fish, aquatic invertebrates, and aquatic life stages of amphibians. Do not apply directly to water or wetlands (i.e., swamps, bogs, marshes, and potholes). Drift and runoff may be hazardous to aquatic organisms in areas near the application site. Do not contaminate water when disposing of equipment washwaters."

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

**BEST AVAILABLE COPY**

APPLICATION BY HAND

- DAIRY AND BEEF CATTLE: For control of Hornflies, apply 2 heaping table-spoonsfull (2 ounces) to the back and flanks of infested animals. Rub lightly into hair. Repeat application no more than once every 3 weeks.
- BEEF CATTLE: For control of Lice and Ticks on Beef Cattle, use as above. Treat animals thoroughly. Repeat treatment as necessary.
- FACE FLIES: As an aid in control of face flies on Beef Cattle, dust the faces of cattle thoroughly. Frequent repeated applications may be necessary.
- HOGS: For control of Hog Lice, apply to Hogs, pens and litter. One treatment maybe sufficient. Treat animals thoroughly. Repeat in 10 days or thereafter as needed.
- SHEEP & GOATS: For the control of Lice and Keds on Sheep and Goats, treat animals thoroughly, preferably following shearing. Repeat as necessary.

DUST BAG APPLICATION

BEEF CATTLE: To aid in control of Hornflies and Lice, hang dust bag where cattle tend to congregate. For Beef Cattle a good location is near salt blocks, mineral blocks, or water supply. Do not hang dust bags directly over mineral, salt, feed or water supply. Dust bags should be hung so that the bottom of the bag is 4" to 6" below the topline of the cattle. Allow one dust bag for each 10 to 15 animals. Do not use dust bag application for Dairy Cattle.

STORAGE & DISPOSAL

Do not contaminate food, feedstuffs or water supply by storage or disposal. Store in original container in locked storage area. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary land fill or by incineration, or if allowed by State and Local authorities, by burning. If burned, stay out of smoke.