


UNICORN
MALATHION
LIVESTOCK SPRAY

Controls Hornflies, Ticks, Lice

ON

Beef Cattle, Sheep, Goats, Hogs, Horses



ACTIVE INGREDIENTS:

Malathion*	57%
INERT INGREDIENTS:**	43%

*0,0-dimethyl phosphorodithioate of diethyl
mercaptosuccinate

**Contains Aromatic Petroleum Solvent

Contains 5 lb. Malathion per gallon


KEEP OUT OF REACH OF CHILDREN

WARNING

See Back Panel for Additional Precautions

Net Contents: 1 Gallon

Manufactured By
UNICORN LABORATORIES
HOUSTON, TX. 77090



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing spray mist. Causes eye irritation. Do not get in eyes, skin or clothing. Do not treat animal under one month of age. Do not treat lactating dairy animals or non-lactating dairy animals within two weeks of freshening. Do not contaminate feed, food or watering troughs.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call Physician or Poison Control Center at once. Do not induce vomiting, unless directed by physician.
IF ON SKIN: Wash with soap and water. Remove contaminated clothing. Wash clothes be for reuse.
IF IN EYES: Flush with plenty of water. Get medical attention if irritation persists.
IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated.

Note to Physician: Malathion is a cholinesterase inhibitor. Atropine Sulfate is antidotal. 2-PAN may be effective as an adjunct to atropine.

PHYSICAL & CHEMICAL HAZARDS


Do not use, pour, spill or store near heat or open flame.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to lakes, streams or ponds. This product is highly toxic to bees exposed to direct treatment or to residues remaining in treated area.

DIRECTION FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.




Livestock Pest Control

Animal	Pest	Rate	Remarks
Hogs	Lice	1 gal per 100 gals of water OR 6½ ozs per 5 gals of water.	Apply complete coverage spray to animals, pens and litter. One treatment may be sufficient. Repeat only if needed.
	Sarcoptic mange	1 gal per 100 gals of water OR 6½ ozs per 5 gals of water.	Apply complete coverage spray to all animals in herd, bedding and walls. Use extreme care to thoroughly cover all body surfaces of the animal, including inside of ears. One thorough spraying will usually control sarcoptic mange of swine, however, extensive cases will require second treatment about 10 days after first application.
			After spraying, swine should be kept out of sun and wind for a few hours.
			Residue tolerance: 4 ppm in meat, fat, and meat by-products.
Sheep and Goats	Lice Ticks Keds	1 gal per 100 gals of water OR 6½ ozs per 5 gals of water.	Spray animals thoroughly. Repeat application after 2 or 3 weeks if needed. Do not apply to milk goats. Do not treat animals under one month of age.
			Residue tolerance: 4 ppm in meat, fat and meat by-products.
Horses, beef and non-milking cattle	Lice	1 gal per 100 gals of water OR 6½ ozs per 5 gals of water.	Apply complete coverage spray. One treatment may be sufficient. Repeat only if needed.
		BACK-RUBBING DEVICES: To reduce lice apply a mixture of 2% CYTHION (using 57% Emulsifiable Concentrate) in fuel oil. There may also be a reduction in hornflies. These devices should be made continuously accessible, one to each 35-45 head of cattle. Back-rubbing devices must be retreated every 2 to 3 weeks.	
	Ticks	1-2 gals per 100 gals of water OR 6½-13 ozs per 5 gals of water.	Apply complete coverage spray. Repeat at 2 week intervals if needed.
	Hornflies	1-1½ gals per 100 gals of water OR 6½-10 ozs per 5 gals of water.	Apply complete coverage spray. Repeat at 2 week intervals if needed.
Poultry: Chickens, Ducks Geese, Turkeys			Do not apply to lactating dairy animals, or nonlactating dairy animals within 2 weeks of freshening.
			Do not treat animals under one month of age.
			Residue tolerance: 4 ppm in meat, fat and meat by-products.
	Northern fowl mite Poultry lice Chicken red mite	2 lbs per 1 gal water per 100-150 birds	Repeat application in 4-8 weeks or when necessary. As a supplement to premise treatment for chicken red mite.

Northern fowl mite Chicken body lice Shall lice	Tail dipping	8½ ozs per 15 gals water per 400 birds	Hold bird by wings and dip 3 to 4 inches of tail into solution. Treat vent and surrounding areas. Repeat in 7-10 days if necessary.
Chicken red mite Poultry lice	Roost paint	2-7 ozs per 1 gal water	Brush on at rate of 1 pt per 150 ft of roost.
Northern fowl mite Chicken red mite Poultry lice Flies	Premise treatments	4 lbs per 1 gal water	Apply liberally to litter, walls, ceilings, roost nests and adjacent areas. Force spray into cracks and crevices.
Poultry ticks	Premise treatments	6-7 ozs per 1 gal water	Apply liberally to walls, ceilings and adjacent areas. Force spray into cracks and crevices.
Chiggers	Range treatments	1-1½ pt per acre	Treat range thoroughly the day before placing poultry on range. Repeat every 2-3 weeks.

Residue tolerance: 4 ppm in or on meat and meat by-products
Eggs 0.1 ppm (from application to poultry)



STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage: Store in original container away from heat, use precaution to prevent cross contamination of other pesticides, fertilizer, food and feed.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

E.P.A. REG. NO. 28293-123
E.P.A. EST. NO. 28293-TX-1

