

PMB

28293-125

186

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

ost016d1X

JAN - 9 1995


~~34596855~~
17

Mr. Barry J. Santerre
UNICORN LABORATORIES
1000 118th Avenue North
St. Petersburg, Florida 33716

Dear Mr Santerre:

Subjects: Label and CSF Amendment
Delete Use For Mosquito from Labeling and
Replace Claim for Repellent
UNICORN PERMETHRIN RTU Spray
EPA Registration Number 28293-125
Your Amendment Applications Dated
October 13, 1993 and May 10, 1994

The amended labeling referred to above, submitted under a cover letter dated May 10, 1994, in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. Submit three (3) copies of the final printed labeling. A stamped copy is enclosed for your records.

Sincerely,

George T. LaRocca
Product Manager (13)
Insecticide-Rodenticide Branch
Registration Division (7505C)

CC: Paul Lewis
EPA/ARB

CONCURRENCES

SYMBOL							
SURNAME							
DATE							

UNICORN

PERMETHRIN RTU SPRAY

For Commercial/Agricultural Use

A Concentrated PERMANONE Stable Emulsion INSECTICIDE

For Use Outdoors and in Homes, Kennels and Non-Food Areas of Commercial Buildings: Warehouse, Theaters, Office Buildings, Schools, Motels, Hotels and Food/Feed Processing Establishments.

For Control of Fleas, Brown Dog Ticks and Lice on Premises and on Dogs.

For Control of Ticks which may carry Lyme Disease

ACTIVE INGREDIENT:

Permethrin*..... 0.50%

INERT INGREDIENTS:..... 99.50%

100.00%

* (3-phenoxyphenyl) methyl (±) cis/trans 3-(2,2-dichloroethenyl) 2,2-dimethylcyclopropanecarboxylate. Cis/trans ratio: Min. 35% (±) cis and Max. 65% (±) trans

PERMANONE - Registered Trademark of Fairfield American Corporation.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Rear Panel for Use Cautions.

NET CONTENTS: ONE GALLON

Contains 8.29 lb. permethrin per gallon

ACCEPTED
JAN - 9 1995
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 28293-125

MANUFACTURED BY:

Unicorn Laboratories
1000 118th Avenue North
St. Petersburg, Florida 33716

EPA REG. NO. 28293-125(E:9/22/92)

EPA EST. NO. 28293-FL-1

30/6

(

PRECAUTIONARY STATEMENTS

Hazardous to Humans and Domestic Animals

CAUTION: Harmful if swallowed. May be absorbed through skin. Avoid inhalation of spray mist. Avoid contact with skin, eyes, or clothing. Wash thoroughly after handling and before smoking or eating. Avoid contamination of feed or foodstuffs. Remove pets and birds and cover fish aquaria before space spraying or surface applications. Do not use on humans. Vacate room after treatment and ventilate before reoccupying. Do not allow children or pets to contact treated areas until surfaces are dry.

Environmental Hazards

This product is extremely toxic to fish and other aquatic organisms. Do not apply directly to water, to areas where surface water is present or to intertidal or to intertidal areas below the high water mark.

Statement of Practical Treatment

- IF SWALLOWED:** Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Call a physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person.
- IF INHALED:** Remove affected person to fresh air. Apply artificial respiration if indicated.
- IF ON SKIN:** Remove contaminated clothing and wash affected areas with soap and water.
- IF IN EYES:** Flush eyes with plenty of water. Call a physician if irritation persists.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SHAKE WELL BEFORE USING

INDOOR APPLICATION

Do not use in food/feed areas of food/feed handling establishments, restaurants or other areas where food/feed is commercially prepared or processed. Do not use in serving areas while food/feed is exposed or facility is in operation. Serving areas are areas where prepared food/feed are served such as dining rooms, but excluding areas where food/feed may be prepared or held. In the home all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food/feed should be covered or removed.

Non-food/feed areas are areas such as garbage rooms, lavatories, floor drains (to sewers, entries and vestibules), offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning and bottling).

Not for use in USDA Meat and Poultry Plants.

Vacate treated areas and ventilate before reoccupying. For maximum effectiveness, a combination of spot, surface treatment and fogging is recommended.

Do not apply to classrooms while in use.

SURFACE SPRAYING: To control cockroaches, waterbugs, palmetto bugs, scorpions, millipedes, centipedes, sowbugs, pillbugs, ants, silverfish, firebrats, spiders, crickets, clover mites, cheese mites, granary weevils, rice weevils, confused flour beetles, rust red flour beetles, drugstore beetles, meal worms, grain mites and cadelles: Use a good sprayer adjusted to a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery, and behind shelves and drawers. For silverfish, spray bookcases. For ants, spray trails, nests and points of entry. If surface application only is to be used, spray floors and other surfaces applying at a rate of 1 gallon to 750 square feet of surface.

To control Carpet Beetles: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, crevices and infested areas of shelving. Repeat treatment as needed. Allow all sprayed articles to dry thoroughly before use.

To control Bedbugs: Spray mattresses lightly, particularly around tufts and seams. Take keds apart and spray in all joints. Treat baseboards, moldings and floors. Repeat treatment as needed. Allow all sprayed articles to dry thoroughly before use.

To control Fleas (adult and larvae) and Ticks, Including Ticks which may carry Lyme disease (adult and larvae): Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and entire areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Repeat treatment as needed. Allow to dry thoroughly before allowing pets to contact treated areas.

To control Clothes Moths: Remove any infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly, keeping nozzle at least three feet from fabric to avoid staining. Allow all sprayed articles to dry thoroughly before use.

To kill Lice and Louse Eggs: Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Spray from a distance of 8 to 10 inches. Treat only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned. Allow all treated articles to dry thoroughly before use.

SPACE SPRAYING: Close doors and windows and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. For rapid control of house flies, fruit flies, gnats, mosquitoes, skipper flies, wasps, hornets, bees, black flies, Angoumois grain moths, Mediterranean flour moths, Indian meal moths and tobacco moths: Direct spray at an upward angle, distributing it uniformly throughout the entire area at a rate of 1 ounce per 1000 cubic feet of space. Keep area closed for two hours. Ventilate before reoccupying. Repels house flies.

196

FOGGING: To kill exposed or accessible states of the above named crawling insects, use a rate of 1 gallon to 20,000 to 40,000 cubic feet of space. Contact insects directly when possible. Keep area closed for two hours. Ventilate before reoccupying.

ON LIVESTOCK: To protect cattle (beef and dairy), goats, sheep, hogs and horses from horn flies, face flies, house flies, mosquitoes and gnats: Apply a light mist sufficient to wet the surface of the hair. For face flies, spray face and head, but do not spray into eyes. Kills and repels horn flies up to 4 days.

To control stable flies, horse flies and deer flies: Apply at a rate of 2 ounces per adult animal, sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

To control blood-sucking lice: Apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every three weeks if required.

To control poultry lice: Spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

To control bedbugs and mites in poultry houses: Spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide.

ON DOGS: To control fleas, ticks and lice: Start spraying at the tail, moving the dispenser rapidly and making sure that the animal's entire body is covered, including the legs and underbody. While spraying, fluff the hair so that the spray will penetrate to the skin. Make sure spray wets thoroughly. Do not spray into eyes or face. Avoid contact with genitalia. Repeat as necessary.

NOTE: Not recommended for use on puppies less than four weeks old. Do not use on old, sick or debilitated animals.

OUTDOOR APPLICATION

TO CONTROL TICKS, INCLUDING TICKS WHICH MAY CARRY LYME DISEASE (PERIMETER TREATMENT):

Use Unicorn Permethrin RTU Spray as a residual treatment to control Ticks including the Deer Tick (*Ixodes dammini*), Lone Star Tick (*Amblyomma americanum*), Brown Dog Tick (*Rhipicephalus sanguineus*), American Dog Tick (*Dermacentor variabilis*) and Western Black-legged Tick (*Ixodes pacificus*).

For each areas listed below, apply until surface is wet (approximately 1 gal/750-1000 ft²).

Treat around windows and doors, porches, screens, eaves, patios, garages, under stairways and in crawl spaces where these pests may enter.

Treat a band of vegetation and leaf litter 6 to 10 feet wide around the outside of buildings to help control ticks occurring there.

To help control ticks in outdoor activity areas, treat a band of vegetation and leaf litter 6 - 10 feet wide and adjacent to activity areas, especially where dense vegetation occurs.

676

FOR CONTROL OF GARDEN INSECTS: For use on roses, dahlias, asters and other ornamentals to kill Japanese beetles hit by spray as well as whiteflies, aphids, exposed thrips, spiders, red mites and leafminers.

May be used on plants as a multi-purpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas.

May also be used on other plants such as African violets, asters, azaleas, begonias, camellias, carnations, chrysanthemums, dahlias, delphiniums, dogwoods, English ivy, eunonymus, fuschia, geraniums, crassula, grape vines, Kentia palms, laurels, marigolds, rhododendrons, roses, rubber plants, snapdragons, stocks, wandering Jews and zinnias.

Effective when used as directed against whiteflies, aphids, spider and red mites, exposed thrips, lace bugs, Japanese beetles, pavement ants, clover mites and armyworms. When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90°F. Contact insects directly whenever possible. Repeat treatment as infestations occur.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE AND SPILL PROCEDURES: Store upright at room temperature. Do not expose to freezing temperatures. Avoid exposure to extreme temperatures. In case of spill or leakage, soak up with an absorbent material such as sand, sawdust, fuller's earth, etc. Dispose of with chemical waste.

PESTICIDE DISPOSAL: Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of at or by an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent), then offer to recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved State and local procedures.

CONTAINERS ONE GALLON AND SMALLER: Do not transport or store below 32°F. Do not reuse container. Wrap container in several layers of newspaper and discard in trash.

BUYER ASSUMES ALL RISKS OF USE, STORAGE OR HANDLING OF THIS PRODUCT NOT IN STRICT ACCORDANCE WITH DIRECTIONS GIVEN HEREWITH.